

Tetsuya Ohira

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9351643/publications.pdf>

Version: 2024-02-01

192
papers

5,504
citations

101535

36
h-index

102480

66
g-index

203
all docs

203
docs citations

203
times ranked

5799
citing authors

#	ARTICLE	IF	CITATIONS
1	Dietary fiber intake and risk of incident disabling dementia: the Circulatory Risk in Communities Study. <i>Nutritional Neuroscience</i> , 2023, 26, 148-155.	3.1	13
2	Impact of Major Cardiovascular Risk Factors on the Incidence of Cardiovascular Disease among Overweight and Non-Overweight Individuals: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2022, 29, 422-437.	2.0	6
3	Association Between Lifestyle Habits and the Prevalence of Abdominal Obesity After the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Journal of Epidemiology</i> , 2022, 32, 496-501.	2.4	4
4	Associations Between Occupational Status, Support at Work, and Salivary Cortisol Levels. <i>International Journal of Behavioral Medicine</i> , 2022, 29, 299-307.	1.7	3
5	Accurate endoscopic identification of the afferent limb at the Y anastomosis using the fold disruption sign after gastric resection with Roux-Y reconstruction. <i>Digestive Endoscopy</i> , 2022, 34, 238-243.	2.3	5
6	The Association between Parenting Confidence and Later Child Mental Health in the Area Affected by the Fukushima Nuclear Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 476.	2.6	0
7	Does laughing with others lower the risk of functional disability among older Japanese adults? The JAGES prospective cohort study. <i>Preventive Medicine</i> , 2022, 155, 106945.	3.4	6
8	Lifestyle Factors Associated with Undernutrition in Older People after the Great East Japan Earthquake: A Prospective Study in the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 3399.	2.6	1
9	Effects of a laughter program on body weight and mental health among Japanese people with metabolic syndrome risk factors: a randomized controlled trial. <i>BMC Geriatrics</i> , 2022, 22, 361.	2.7	3
10	The Associations between Evacuation Status and Lifestyle-Related Diseases in Fukushima after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 5661.	2.6	1
11	Lifestyle-related diseases caused by evacuation: Results of the comprehensive health check in the Fukushima health management survey. , 2022, , 99-121.		0
12	Development of an Application for Sustainable Support of Returning Residents Displaced by the Fukushima Nuclear Accident. <i>Studies in Health Technology and Informatics</i> , 2022, , .	0.3	0
13	Changes in the proportion of anemia among young women after the Great East Japan Earthquake: the Fukushima health management survey. <i>Scientific Reports</i> , 2022, 12, .	3.3	2
14	Impact of lifestyle and psychosocial factors on the onset of hypertension after the Great East Japan earthquake: a 7-year follow-up of the Fukushima Health Management Survey. <i>Hypertension Research</i> , 2022, 45, 1609-1621.	2.7	2
15	Does Laughter Predict Onset of Functional Disability and Mortality Among Older Japanese Adults? The JAGES Prospective Cohort Study. <i>Journal of Epidemiology</i> , 2021, 31, 301-307.	2.4	20
16	Dietary pattern changes in Fukushima residents after the Great East Japan Earthquake: the Fukushima Health Management Survey 2011-2013. <i>Public Health Nutrition</i> , 2021, 24, 2195-2204.	2.2	6
17	Longitudinal Trends in Blood Pressure Associated With the Frequency of Laughter: The Circulatory Risk in Communities Study (CIRCS), a Longitudinal Study of the Japanese General Population. <i>Journal of Epidemiology</i> , 2021, 31, 125-131.	2.4	6
18	Association of Admission Hyperglycemia with Clinical Outcomes in Japanese Patients with Acute Large Vessel Occlusion Stroke: A post hoc Analysis of the Recovery by Endovascular Salvage for Cerebral Ultra-Acute Embolism Japan Registry 2. <i>Cerebrovascular Diseases</i> , 2021, 50, 12-19.	1.7	8

#	ARTICLE	IF	CITATIONS
19	Higher psychological distress experienced by evacuees relocating outside Fukushima after the nuclear accident: The Fukushima Health Management Survey. <i>International Journal of Disaster Risk Reduction</i> , 2021, 52, 101962.	3.9	9
20	Salt taste perception and blood pressure levels in population-based samples: the Circulatory Risk in Communities Study (CIRCS). <i>British Journal of Nutrition</i> , 2021, 125, 203-211.	2.3	2
21	Impact of Perceived Social Support on the Association Between Anger Expression and the Risk of Stroke: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2021, , .	2.4	2
22	Psychosocial support for the examinees and their families during the secondary confirmatory examination: Analyses of support records at first visit. <i>Fukushima Journal of Medical Sciences</i> , 2021, 67, 53-63.	0.4	1
23	Socioeconomic status, damage-related conditions, and PTSD following the Fukushima-daiichi nuclear power plant accident: The Fukushima Health Management Survey. <i>Fukushima Journal of Medical Sciences</i> , 2021, 67, 71-82.	0.4	4
24	Dietary Patterns and Progression of Impaired Kidney Function in Japanese Adults: A Longitudinal Analysis for the Fukushima Health Management Survey, 2011â€”2015. <i>Nutrients</i> , 2021, 13, 168.	4.1	7
25	Relationship Between Risk of Hyper-Low-density Lipoprotein Cholesterolemia and Evacuation After the Great East Japan Earthquake. <i>Journal of Epidemiology</i> , 2021, , .	2.4	4
26	Serum High-Sensitivity C-Reactive Protein Levels and the Risk of Atrial Fibrillation in Japanese Population: the Circulatory Risk in Communities Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 194-202.	2.0	11
27	Serum Albumin and Risks of Stroke and Its Subtypesâ€”The Circulatory Risk in Communities Study (CIRCS) â€”. <i>Circulation Journal</i> , 2021, 85, 385-392.	1.6	8
28	The association between body mass index and recovery from post-traumatic stress disorder after the nuclear accident in Fukushima. <i>Scientific Reports</i> , 2021, 11, 5330.	3.3	2
29	Response to the Letter to the Editor: â€”Absorbed radiation doses in the thyroid as estimated by UNSCEAR and subsequent risk of childhood thyroid cancer following the Great East Japan Earthquakeâ€”™, by Ohira et al.. <i>Journal of Radiation Research</i> , 2021, 62, 425-426.	1.6	1
30	Association of habitual exercise with adultsâ€™ mental health following the Fukushima Daiichi nuclear power plant accident: the Fukushima Health Management Survey. <i>Mental Health and Physical Activity</i> , 2021, 20, 100388.	1.8	1
31	Evacuation after the Great East Japan Earthquake is an independent factor associated with hyperuricemia: The Fukushima Health Management Survey. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1177-1188.	2.6	9
32	Trajectories of peer relationship problems and emotional symptoms in children 5Â½years after a nuclear disaster: Fukushima Health Management Survey. <i>Journal of Radiation Research</i> , 2021, 62, i114-i121.	1.6	2
33	Factors Hindering Social Participation among Older Residents from Evacuation Zones after the Nuclear Power Plant Accident in Fukushima: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 4426.	2.6	3
34	Relationship between physical activity/exercise habits and the frequency of new onset of lifestyle-related diseases after the Great East Japan Earthquake among residents in Fukushima: the Fukushima Health Management Survey. <i>Journal of Radiation Research</i> , 2021, 62, i129-i139.	1.6	5
35	Needs Survey for Health Support Application Development Project for Residents Returning from Evacuation After the Fukushima Nuclear Accident. <i>Studies in Health Technology and Informatics</i> , 2021, 281, 1091-1092.	0.3	0
36	Association between Psychosocial Factors and Oral Symptoms among Residents in Fukushima after the Great East Japan Earthquake: A Cross-Sectional Study from the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 6054.	2.6	1

#	ARTICLE	IF	CITATIONS
37	Relationship between Endothelial Dysfunction and Prevalence of Chronic Kidney Disease: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2021, 28, 622-629.	2.0	8
38	Associations of the COVID-19 pandemic with the economic status and mental health of people affected by the Fukushima disaster using the difference-in-differences method: The Fukushima Health Management Survey. <i>SSM - Population Health</i> , 2021, 14, 100801.	2.7	6
39	Synergistic Effect of History of Cardiovascular Disease and Mental Distress on Post-Traumatic Stress Disorder after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 10283.	2.6	0
40	Association between post-traumatic stress disorder symptoms and bone fractures after the Great East Japan Earthquake in older adults: a prospective cohort study from the Fukushima Health Management Survey. <i>BMC Geriatrics</i> , 2021, 21, 18.	2.7	7
41	Association between frequency of laughter and oral health among community-dwelling older adults: a population-based cross-sectional study in Japan. <i>Quality of Life Research</i> , 2021, 30, 1561-1569.	3.1	6
42	Fine Needle Aspiration Cytology Implementation and Malignancy Rates in Children and Adolescents Based on Japanese Guidelines: The Fukushima Health Management Survey. <i>Thyroid</i> , 2021, 31, 1683-1692.	4.5	4
43	Association between Laughter and Lifestyle Diseases after the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 12699.	2.6	1
44	Factors Influencing the Proportion of Non-examinees in the Fukushima Health Management Survey for Childhood and Adolescent Thyroid Cancer: Results From the Baseline Survey. <i>Journal of Epidemiology</i> , 2020, 30, 301-308.	2.4	3
45	Communication between Health Professionals and Community Residents in Fukushima: A Focus on the Feedback Loop. <i>Health Communication</i> , 2020, 35, 1274-1282.	3.1	17
46	Changes in the mental health status of adolescents following the Fukushima Daiichi nuclear accident and related factors: Fukushima Health Management Survey. <i>Journal of Affective Disorders</i> , 2020, 260, 432-439.	4.1	12
47	The association between self-reported sleep dissatisfaction after the Great East Japan Earthquake, and a deteriorated socioeconomic status in the evacuation area: the Fukushima Health Management Survey. <i>Sleep Medicine</i> , 2020, 68, 63-70.	1.6	11
48	AGE DEPENDENCE OF INDIVIDUAL EXTERNAL DOSES IN AN EARLY STAGE AFTER THE FUKUSHIMA NUCLEAR ACCIDENT. <i>Radiation Protection Dosimetry</i> , 2020, 188, 238-245.	0.8	4
49	Associations between Dietary Patterns and Cardiometabolic Risks in Japan: A Cross-Sectional Study from the Fukushima Health Management Survey, 2011â€“2015. <i>Nutrients</i> , 2020, 12, 129.	4.1	13
50	Blood concentration of tacrolimus and age predict tacrolimus-induced left ventricular dysfunction after bone marrow transplantation in adults. <i>Journal of Medical Ultrasonics (2001)</i> , 2020, 47, 97-105.	1.3	2
51	Anger Expression and the Risk of Cardiovascular Disease Among Urban and Rural Japanese Residents: The Circulatory Risk in Communities Study. <i>Psychosomatic Medicine</i> , 2020, 82, 215-223.	2.0	10
52	The Impact of Valvuloarterial Impedance on Left Ventricular Geometrical Change after Transcatheter Aortic Valve Replacement: A Comparison between Valvuloarterial Impedance and Mean Pressure Gradient. <i>Journal of Clinical Medicine</i> , 2020, 9, 3143.	2.4	0
53	Nested matched case control study for the Japan Fukushima Health Management Survey's first full-scale (second-round) thyroid examination. <i>Medicine (United States)</i> , 2020, 99, e20440.	1.0	6
54	Influence of post-disaster evacuation on incidence of hyperuricemia in residents of Fukushima Prefecture: the Fukushima Health Management Survey. <i>Clinical and Experimental Nephrology</i> , 2020, 24, 1025-1032.	1.6	7

#	ARTICLE	IF	CITATIONS
55	All-Cause Mortality After the Great East Japan Earthquake in Fukushima Prefecture: Trends From 2009 to 2016 and Variation by Displacement. <i>Disaster Medicine and Public Health Preparedness</i> , 2020, , 1-4.	1.3	1
56	Association between Psychological Factors and Evacuation Status and the Incidence of Cardiovascular Diseases after the Great East Japan Earthquake: A Prospective Study of the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7832.	2.6	6
57	Joint impact of muscle mass and waist circumference on type 2 diabetes in Japanese middle-aged adults: The Circulatory Risk in Communities Study (<scp>CIRCS</scp>). <i>Journal of Diabetes</i> , 2020, 12, 677-685.	1.8	5
58	Lifestyle factors associated with prevalent and exacerbated musculoskeletal pain after the Great East Japan Earthquake: a cross-sectional study from the Fukushima Health Management Survey. <i>BMC Public Health</i> , 2020, 20, 677.	2.9	7
59	Urinary 8-Hydroxy-2'-Deoxyguanosine Levels and Cardiovascular Disease Incidence in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1086-1096.	2.0	4
60	Relationship between the prevalence of polycythemia and factors observed in the mental health and lifestyle survey after the Great East Japan Earthquake. <i>Medicine (United States)</i> , 2020, 99, e18486.	1.0	6
61	Cardiovascular reactivity to acute stress associated with sickness absence among Japanese men and women: A prospective study. <i>Brain and Behavior</i> , 2020, 10, e01541.	2.2	3
62	Influence of post-disaster evacuation on childhood obesity and hyperlipidemia. <i>Pediatrics International</i> , 2020, 62, 669-676.	0.5	12
63	Serum uric acid and risk of stroke and its types: the Circulatory Risk in Communities Study (CIRCS). <i>Hypertension Research</i> , 2020, 43, 313-321.	2.7	30
64	Absorbed radiation doses in the thyroid as estimated by UNSCEAR and subsequent risk of childhood thyroid cancer following the Great East Japan Earthquake. <i>Journal of Radiation Research</i> , 2020, 61, 243-248.	1.6	29
65	Effects of Psychological and Lifestyle Factors on Metabolic Syndrome Following the Fukushima Daiichi Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1010-1018.	2.0	11
66	Changes in drinking behavior among evacuees after the Fukushima Daiichi Nuclear Power Plant accident: the Fukushima Health Management Survey. <i>Fukushima Journal of Medical Sciences</i> , 2020, 66, 133-142.	0.4	4
67	Analysis of direction of association between radiation risk perception and relocation using a random-intercept and cross lagged panel model: The Fukushima Health Management Survey. <i>SSM - Population Health</i> , 2020, 12, 100706.	2.7	5
68	Job stress factors measured by Brief Job Stress Questionnaire and sickness absence among Japanese workers: A longitudinal study. <i>Fukushima Journal of Medical Sciences</i> , 2020, 66, 88-96.	0.4	5
69	Cytological examination of the thyroid in children and adolescents after the Fukushima Nuclear Power Plant accident: the Fukushima Health Management Survey. <i>Endocrine Journal</i> , 2020, 67, 1233-1238.	1.6	1
70	Parental Recognition of Bullying and Associated Factors Among Children After the Fukushima Nuclear Disaster: A 3-Year Follow-Up Study From the Fukushima Health Management Survey. <i>Frontiers in Psychiatry</i> , 2019, 10, 283.	2.6	6
71	Effects of laughter therapy on quality of life in patients with cancer: An open-label, randomized controlled trial. <i>PLoS ONE</i> , 2019, 14, e0219065.	2.5	29
72	Risk Factors for Problem Drinking among Evacuees in Fukushima following the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Tohoku Journal of Experimental Medicine</i> , 2019, 248, 239-252.	1.2	14

#	ARTICLE	IF	CITATIONS
73	Unreliable Information as a Risk Factor for Worse Mental Fatigue among Residents in Fukushima after the Nuclear Power Station Accident. <i>Tohoku Journal of Experimental Medicine</i> , 2019, 248, 261-272.	1.2	7
74	Who needs care? - The long-term trends and geographical distribution of deaths due to acute myocardial infarction in Fukushima Prefecture following the Great East Japan Earthquake. <i>International Journal of Disaster Risk Reduction</i> , 2019, 41, 101318.	3.9	3
75	Development and Implementation of an Internet Survey to Assess Community Health in the Face of a Health Crisis: Data from the Pregnancy and Birth Survey of the Fukushima Health Management Survey, 2016. <i>International Journal of Environmental Research and Public Health</i> , 2019, 16, 1946.	2.6	2
76	Trends in the Incidence of Sudden Deaths and Heart Diseases in Fukushima After the Great East Japan Earthquake. <i>International Heart Journal</i> , 2019, 60, 1253-1258.	1.0	7
77	The latest update on individual external doses in an early stage after the Fukushima nuclear accident. <i>Radiation Protection Dosimetry</i> , 2019, 187, 402-406.	0.8	7
78	Common carotid artery kinking is a predictor of cardiovascular events: A long-term follow-up study using carotid ultrasonography. <i>Echocardiography</i> , 2019, 36, 2227-2233.	0.9	5
79	External Radiation Dose, Obesity, and Risk of Childhood Thyroid Cancer After the Fukushima Daiichi Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Epidemiology</i> , 2019, 30, 853-860.	2.7	22
80	The Authors Respond. <i>Epidemiology</i> , 2019, 30, e11.	2.7	2
81	Psychometric evaluation of the simplified Japanese version of the Athens Insomnia Scale: The Fukushima Health Management Survey. <i>Journal of Sleep Research</i> , 2019, 28, e12771.	3.2	17
82	Obesity and mental health improvement following nutritional education focusing on gut microbiota composition in Japanese women: a randomised controlled trial. <i>European Journal of Nutrition</i> , 2019, 58, 3291-3302.	3.9	31
83	Incidence of Thyroid Cancer Among Children and Young Adults in Fukushima, Japan, Screened With 2 Rounds of Ultrasonography Within 5 Years of the 2011 Fukushima Daiichi Nuclear Power Station Accident. <i>JAMA Otolaryngology - Head and Neck Surgery</i> , 2019, 145, 4.	2.2	44
84	Mental health promotion at work associated with utilization of the "Guide of Mental Health Professionals for Workers in Osaka". <i>Journal of Occupational Safety and Health</i> , 2019, 12, 145-151.	0.0	0
85	Findings of Thyroid Ultrasound Examination Within 3 Years After the Fukushima Nuclear Power Plant Accident: The Fukushima Health Management Survey. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 861-869.	3.6	28
86	Impact of evacuation on trends in the prevalence, treatment, and control of hypertension before and after a disaster. <i>Journal of Hypertension</i> , 2018, 36, 924-932.	0.5	15
87	Associations of Tobacco Smoking with Impaired Endothelial Function: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 836-845.	2.0	20
88	Association between poor psychosocial conditions and diabetic nephropathy in Japanese type 2 diabetes patients: A cross-sectional study. <i>Journal of Diabetes Investigation</i> , 2018, 9, 162-172.	2.4	9
89	Lifestyle factors and social ties associated with the frequency of laughter after the Great East Japan Earthquake: Fukushima Health Management Survey. <i>Quality of Life Research</i> , 2018, 27, 639-650.	3.1	18
90	The Authors Respond. <i>Epidemiology</i> , 2018, 29, e57-e58.	2.7	0

#	ARTICLE	IF	CITATIONS
91	Associations of disaster-related and psychosocial factors with changes in smoking status after a disaster: a cross-sectional survey after the Great East Japan Earthquake. <i>BMJ Open</i> , 2018, 8, e018943.	1.9	14
92	Effects of lifestyle on hepatobiliary enzyme abnormalities following the Fukushima Daiichi nuclear power plant accident. <i>Medicine (United States)</i> , 2018, 97, e12890.	1.0	7
93	Spatial analysis of the geographical distribution of thyroid cancer cases from the first-round thyroid ultrasound examination in Fukushima Prefecture. <i>Scientific Reports</i> , 2018, 8, 17661.	3.3	13
94	Nonfasting Glucose and Incident Stroke and Its Types—The Circulatory Risk in Communities Study (CIRCS) —. <i>Circulation Journal</i> , 2018, 82, 1598-1604.	1.6	8
95	Reduction of radiation-related anxiety promoted wellbeing after the 2011 disaster: “Fukushima Health Management Survey”. <i>Journal of Radiological Protection</i> , 2018, 38, 1428-1440.	1.1	19
96	Changes in Risk Perception of the Health Effects of Radiation and Mental Health Status: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 1219.	2.6	39
97	Serum Fatty Acid and Risk of Coronary Artery Disease—The Circulatory Risk in Communities Study (CIRCS) —. <i>Circulation Journal</i> , 2018, 82, 3013-3020.	1.6	26
98	The Relationship between Sleep Time and Mental Health Problems According to the Strengths and Difficulties Questionnaire in Children after an Earthquake Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 633.	2.6	12
99	Association between Serum Uric Acid Level and Ventricular Tachyarrhythmia in Heart Failure Patients with Implantable Cardioverter-Defibrillator. <i>Cardiology</i> , 2018, 140, 47-51.	1.4	7
100	Trajectories of Emotional Symptoms and Peer Relationship Problems in Children after Nuclear Disaster: Evidence from the Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 82.	2.6	15
101	Effect of Radiological Countermeasures on Subjective Well-Being and Radiation Anxiety after the 2011 Disaster: The Fukushima Health Management Survey. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 124.	2.6	20
102	Impact of social relationships on income—laughter relationships among older people: the JAGES cross-sectional study. <i>BMJ Open</i> , 2018, 8, e019104.	1.9	7
103	Associations Between Childhood Thyroid Cancer and External Radiation Dose After the Fukushima Daiichi Nuclear Power Plant Accident. <i>Epidemiology</i> , 2018, 29, e32-e34.	2.7	23
104	External Dose Estimation in an Early Stage after the Fukushima Daiichi Nuclear Power Plant Accident. <i>Japanese Journal of Health Physics</i> , 2018, 53, 100-110.	0.1	4
105	Serum α -linolenic and other $n-3$ fatty acids, and risk of disabling dementia: Community-based nested case—control study. <i>Clinical Nutrition</i> , 2017, 36, 793-797.	5.0	30
106	Effect of evacuation on liver function after the Fukushima Daiichi Nuclear Power Plant accident: The Fukushima Health Management Survey. <i>Journal of Epidemiology</i> , 2017, 27, 180-185.	2.4	25
107	Persistent prevalence of polycythemia among evacuees 4 years after the Great East Japan Earthquake: A follow-up study. <i>Preventive Medicine Reports</i> , 2017, 5, 251-256.	1.8	8
108	Associations of central aortic pressure and brachial blood pressure with flow mediated dilatation in apparently healthy Japanese men: The Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2017, 259, 46-50.	0.8	8

#	ARTICLE	IF	CITATIONS
109	Changes of Posttraumatic Stress Responses in Evacuated Residents and Their Related Factors. Asia-Pacific Journal of Public Health, 2017, 29, 182S-192S.	1.0	21
110	Exercise Habits Are Important for the Mental Health of Children in Fukushima After the Fukushima Daiichi Disaster. Asia-Pacific Journal of Public Health, 2017, 29, 171S-181S.	1.0	17
111	Changes in Cardiovascular Risk Factors After the Great East Japan Earthquake. Asia-Pacific Journal of Public Health, 2017, 29, 47S-55S.	1.0	64
112	The impact of evacuation on the incidence of chronic kidney disease after the Great East Japan Earthquake: The Fukushima Health Management Survey. Clinical and Experimental Nephrology, 2017, 21, 995-1002.	1.6	26
113	Effects of socioeconomic factors on cardiovascular-related symptoms among residents in Fukushima after the Great East Japan Earthquake: a cross-sectional study using data from the Fukushima Health Management Survey. BMJ Open, 2017, 7, e014077.	1.9	13
114	Comparative Analysis of the Growth Pattern of Thyroid Cancer in Young Patients Screened by Ultrasonography in Japan After a Nuclear Accident. JAMA Otolaryngology - Head and Neck Surgery, 2017, 144, 57-63.	2.2	17
115	Matched-pair analysis of a multi-institutional cohort reveals that epidermal growth factor receptor mutation is not a risk factor for postoperative recurrence of lung adenocarcinoma. Lung Cancer, 2017, 114, 23-30.	2.0	12
116	Association between markers of arterial stiffness and atrial fibrillation in the Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2017, 263, 244-248.	0.8	14
117	Changes in Hepatobiliary Enzyme Abnormality After the Great East Japan Earthquake: The Fukushima Health Management Survey. Scientific Reports, 2017, 7, 710.	3.3	8
118	Lifestyle-related factors that explain disaster-induced changes in socioeconomic status and poor subjective health: a cross-sectional study from the Fukushima health management survey. BMC Public Health, 2017, 17, 340.	2.9	19
119	Prevalence and Characterization of Thyroid Hemiagenesis in Japan: The Fukushima Health Management Survey. Thyroid, 2017, 27, 1011-1016.	4.5	21
120	Evacuation after the Great East Japan Earthquake was associated with poor dietary intake: The Fukushima Health Management Survey. Journal of Epidemiology, 2017, 27, 14-23.	2.4	47
121	Simulation of expected childhood and adolescent thyroid cancer cases in Japan using a cancer-progression model based on the National Cancer Registry. Medicine (United States), 2017, 96, e8631.	1.0	19
122	Influence of Post-disaster Evacuation on Incidence of Metabolic Syndrome. Journal of Atherosclerosis and Thrombosis, 2017, 24, 327-337.	2.0	48
123	Perception of Radiation Risk as a Predictor of Mid-Term Mental Health after a Nuclear Disaster: The Fukushima Health Management Survey. International Journal of Environmental Research and Public Health, 2017, 14, 1067.	2.6	32
124	The Relationship between Starting to Drink and Psychological Distress, Sleep Disturbance after the Great East Japan Earthquake and Nuclear Disaster: The Fukushima Health Management Survey. International Journal of Environmental Research and Public Health, 2017, 14, 1281.	2.6	32
125	Fifty-year Time Trends in Blood Pressures, Body Mass Index and their Relations in a Japanese Community: The Circulatory Risk in Communities Study (CIRCS). Journal of Atherosclerosis and Thrombosis, 2017, 24, 518-529.	2.0	15
126	Effects of Changes in Living Place and Communities on Health and Welfare After the Great East Japan Earthquake: A Lesson from a Case of Fukushima. Trends in the Sciences, 2017, 22, 6_10-6_17.	0.0	0

#	ARTICLE	IF	CITATIONS
127	Three-year trend survey of psychological distress, post-traumatic stress, and problem drinking among residents in the evacuation zone after the Fukushima Daiichi Nuclear Power Plant accident [The Fukushima Health Management Survey]. <i>Psychiatry and Clinical Neurosciences</i> , 2016, 70, 245-252.	1.8	74
128	Explanatory Meetings on Thyroid Examination for the "Fukushima Health Management Survey" after the Great East Japan Earthquake: Reduction of Anxiety and Improvement of Comprehension. <i>Tohoku Journal of Experimental Medicine</i> , 2016, 239, 333-343.	1.2	26
129	Prevalence of Renal Dysfunction among Evacuees and Non-evacuees after the Great East Earthquake: Results from the Fukushima Health Management Survey. <i>Internal Medicine</i> , 2016, 55, 2563-2569.	0.7	12
130	Hypo-high-density Lipoprotein Cholesterolemia Caused by Evacuation after the Fukushima Daiichi Nuclear Power Plant Accident: Results from the Fukushima Health Management Survey. <i>Internal Medicine</i> , 2016, 55, 1967-1976.	0.7	34
131	Association between psychological distress and dietary intake among evacuees after the Great East Japan Earthquake in a cross-sectional study: the Fukushima Health Management Survey. <i>BMJ Open</i> , 2016, 6, e011534.	1.9	37
132	Comparison of childhood thyroid cancer prevalence among 3 areas based on external radiation dose after the Fukushima Daiichi nuclear power plant accident. <i>Medicine (United States)</i> , 2016, 95, e4472.	1.0	46
133	Inappropriate Suppression of Thyrotropin Concentrations in Young Patients with Thyroid Nodules Including Thyroid Cancer: The Fukushima Health Management Survey. <i>Thyroid</i> , 2016, 26, 717-725.	4.5	8
134	Relationship between HbA1c and risk of retinal hemorrhage in the Japanese general population: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 834-838.	2.3	2
135	Occupational status and job stress in relation to cardiovascular stress reactivity in Japanese workers. <i>Preventive Medicine Reports</i> , 2016, 4, 61-67.	1.8	19
136	Comprehensive Survey Results of Childhood Thyroid Ultrasound Examinations in Fukushima in the First Four Years After the Fukushima Daiichi Nuclear Power Plant Accident. <i>Thyroid</i> , 2016, 26, 843-851.	4.5	65
137	Dehydroepiandrosterone-sulfate is associated with cardiovascular reactivity to stress in women. <i>Psychoneuroendocrinology</i> , 2016, 69, 116-122.	2.7	9
138	Evacuation and Risk of Hypertension After the Great East Japan Earthquake. <i>Hypertension</i> , 2016, 68, 558-564.	2.7	83
139	Drinking Behavior and Mental Illness Among Evacuees in Fukushima Following the Great East Japan Earthquake: The Fukushima Health Management Survey. <i>Alcoholism: Clinical and Experimental Research</i> , 2016, 40, 623-630.	2.4	33
140	Laughter is the Best Medicine? A Cross-Sectional Study of Cardiovascular Disease Among Older Japanese Adults. <i>Journal of Epidemiology</i> , 2016, 26, 546-552.	2.4	95
141	Predictors of severe psychological distress trajectory after nuclear disaster: evidence from the Fukushima Health Management Survey. <i>BMJ Open</i> , 2016, 6, e013400.	1.9	42
142	Epidermal growth factor receptor gene mutation as risk factor for recurrence in patients with surgically resected lung adenocarcinoma: a matched-pair analysis. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2016, 23, 216-222.	1.1	13
143	Psychometric Evaluation of the Japanese Version of the Posttraumatic Stress Disorder Checklist in Community Dwellers Following the Fukushima Daiichi Nuclear Power Plant Incident. <i>SAGE Open</i> , 2016, 6, 215824401665244.	1.7	24
144	The relationship between sodium concentrations in spot urine and blood pressure increases: a prospective study of Japanese general population: the Circulatory Risk in Communities Study (CIRCS). <i>BMC Cardiovascular Disorders</i> , 2016, 16, 55.	1.7	10

#	ARTICLE	IF	CITATIONS
145	Effect of Evacuation on Body Weight After the Great East Japan Earthquake. American Journal of Preventive Medicine, 2016, 50, 553-560.	3.0	73
146	A survey on psychiatric services for an occupational mental health program in Osaka prefecture. Journal of Occupational Safety and Health, 2016, 9, 9-15.	0.0	0
147	Laughter and Subjective Health Among Community-Dwelling Older People in Japan. Journal of Nervous and Mental Disease, 2015, 203, 934-942.	1.0	25
148	Evacuation after the Fukushima Daiichi Nuclear Power Plant Accident Is a Cause of Diabetes: Results from the Fukushima Health Management Survey. Journal of Diabetes Research, 2015, 2015, 1-9.	2.3	63
149	Health effects of radiation and other health problems in the aftermath of nuclear accidents, with an emphasis on Fukushima. Lancet, The, 2015, 386, 479-488.	13.7	195
150	Increased prevalence of atrial fibrillation after the Great East Japan Earthquake: Results from the Fukushima Health Management Survey. International Journal of Cardiology, 2015, 198, 102-105.	1.7	44
151	Effects of a traditional herbal medicine on peripheral blood flow in women experiencing peripheral coldness: a randomized controlled trial. BMC Complementary and Alternative Medicine, 2015, 15, 105.	3.7	23
152	Psychological distress and the perception of radiation risks: the Fukushima health management survey. Bulletin of the World Health Organization, 2015, 93, 598-605.	3.3	160
153	HOW LIFESTYLE AFFECTS HEALTH—CHANGES IN HEALTH STATUS BEFORE AND AFTER THE EARTHQUAKE. Fukushima Journal of Medical Sciences, 2014, 60, 211-212.	0.4	10
154	Life as an evacuee after the Fukushima Daiichi nuclear power plant accident is a cause of polycythemia: the Fukushima Health Management Survey. BMC Public Health, 2014, 14, 1318.	2.9	30
155	Menopausal Status in Relation to Cardiovascular Stress Reactivity in Healthy Japanese Participants. Psychosomatic Medicine, 2014, 76, 701-708.	2.0	11
156	An association between central aortic pressure and subclinical organ damage of the heart among a general Japanese cohort: Circulatory Risk in Communities Study (CIRCS). Atherosclerosis, 2014, 232, 94-98.	0.8	9
157	Adult Height and Body Mass Index in Relation to Risk of Total Stroke and its Subtypes: The Circulatory Risk in Communities Study. Journal of Stroke and Cerebrovascular Diseases, 2014, 23, 667-674.	1.6	29
158	Effects of a laughter and exercise program on physiological and psychological health among community-dwelling elderly in Japan: Randomized controlled trial. Geriatrics and Gerontology International, 2013, 13, 152-160.	1.5	50
159	Alkaline Phosphatase and Risk of Stroke Among Japanese: The Circulatory Risk in Communities Study (CIRCS). Journal of Stroke and Cerebrovascular Diseases, 2013, 22, 1046-1055.	1.6	36
160	Total and high molecular weight adiponectin levels and risk of cardiovascular disease in individuals with high blood glucose levels. Atherosclerosis, 2013, 229, 222-227.	0.8	21
161	Cardiovascular Disease Epidemiology in Asia. Circulation Journal, 2013, 77, 1646-1652.	1.6	217
162	Insulin secretion capacity in normal glucose tolerance, impaired glucose tolerance, and diabetes in obese and non-obese Japanese patients. Journal of Diabetes Investigation, 2012, 3, 271-275.	2.4	24

#	ARTICLE	IF	CITATIONS
163	Associations of Anger, Anxiety, and Depressive Symptoms With Carotid Arterial Wall Thickness. <i>Psychosomatic Medicine</i> , 2012, 74, 517-525.	2.0	30
164	Risk factors for sudden cardiac death among Japanese. <i>Journal of Hypertension</i> , 2012, 30, 1137-1143.	0.5	19
165	Non-fasting blood glucose and risk of incident coronary heart disease in middle-aged general population: The Circulatory Risk in Communities Study (CIRCS). <i>Preventive Medicine</i> , 2012, 55, 603-607.	3.4	22
166	Associations between alcohol consumption and sleep-disordered breathing among Japanese women. <i>Respiratory Medicine</i> , 2011, 105, 796-800.	2.9	9
167	C-reactive protein levels and risk of stroke and its subtype in Japanese: The Circulatory Risk in Communities Study (CIRCS). <i>Atherosclerosis</i> , 2011, 217, 187-193.	0.8	35
168	Carotid Artery Wall Thickness and Risk of Stroke Subtypes. <i>Stroke</i> , 2011, 42, 397-403.	2.0	44
169	Chronic Kidney Disease and Drinking Status in Relation to Risks of Stroke and Its Subtypes. <i>Stroke</i> , 2011, 42, 2531-2537.	2.0	53
170	Psychological Distress and Cardiovascular Disease: The Circulatory Risk in Communities Study (CIRCS). <i>Journal of Epidemiology</i> , 2010, 20, 185-191.	2.4	38
171	Reproductive history, hormone replacement, and incidence of venous thromboembolism: the Longitudinal Investigation of Thromboembolism Etiology. <i>British Journal of Haematology</i> , 2010, 149, 606-612.	2.5	18
172	Î³-Glutamyltranspeptidase and Incident Stroke Among Japanese Men and Women. <i>Stroke</i> , 2010, 41, 385-388.	2.0	40
173	Effects of Habitual Alcohol Intake on Ambulatory Blood Pressure, Heart Rate, and Its Variability Among Japanese Men. <i>Hypertension</i> , 2009, 53, 13-19.	2.7	97
174	Serum and Dietary Magnesium and Risk of Ischemic Stroke: The Atherosclerosis Risk in Communities Study. <i>American Journal of Epidemiology</i> , 2009, 169, 1437-1444.	3.4	118
175	Perceived Level of Life Enjoyment and Risks of Cardiovascular Disease Incidence and Mortality. <i>Circulation</i> , 2009, 120, 956-963.	1.6	72
176	Trends for Blood Pressure and Its Contribution to Stroke Incidence in the Middle-Aged Japanese Population. <i>Stroke</i> , 2009, 40, 1571-1577.	2.0	104
177	Trends in the Incidence of Coronary Heart Disease and Stroke and Their Risk Factors in Japan, 1964 to 2003. <i>Journal of the American College of Cardiology</i> , 2008, 52, 71-79.	2.8	152
178	The joint impact on being overweight of self reported behaviours of eating quickly and eating until full : cross sectional survey. <i>BMJ: British Medical Journal</i> , 2008, 337, a2002-a2002.	2.3	196
179	Associations of Psychosocial Factors With Heart Rate and Its Short-Term Variability: Multi-Ethnic Study of Atherosclerosis. <i>Psychosomatic Medicine</i> , 2008, 70, 141-146.	2.0	44
180	Longitudinal Association of Serum Carotenoids and Tocopherols with Hostility: The CARDIA Study. <i>American Journal of Epidemiology</i> , 2007, 167, 42-50.	3.4	10

#	ARTICLE	IF	CITATIONS
181	Effects of weight training on quality of life in recent breast cancer survivors. Cancer, 2006, 106, 2076-2083.	4.1	179
182	Lipoprotein(a) and Incident Ischemic Stroke. Stroke, 2006, 37, 1407-1412.	2.0	96
183	Risk Factors for Ischemic Stroke Subtypes. Stroke, 2006, 37, 2493-2498.	2.0	329
184	Plasma Fibrinogen Concentrations and Risk of Stroke and Its Subtypes Among Japanese Men and Women. Stroke, 2006, 37, 2488-2492.	2.0	19
185	Carotid Intima-Media Thickness and Plaque Characteristics as a Risk Factor for Stroke in Japanese Elderly Men. Stroke, 2004, 35, 2788-2794.	2.0	262
186	Serum Total Homocysteine Concentrations and Risk of Stroke and Its Subtypes in Japanese. Circulation, 2004, 109, 2766-2772.	1.6	133
187	Prospective Study of Major and Minor ST-T Abnormalities and Risk of Stroke Among Japanese. Stroke, 2003, 34, e250-3.	2.0	29
188	Smoking Raises the Risk of Total and Ischemic Strokes in Hypertensive Men.. Hypertension Research, 2003, 26, 209-217.	2.7	61
189	The relation of anger expression with blood pressure levels and hypertension in rural and urban Japanese communities. Journal of Hypertension, 2002, 20, 21-27.	0.5	32
190	Prospective Study of Depressive Symptoms and Risk of Stroke Among Japanese. Stroke, 2001, 32, 903-908.	2.0	133
191	Validity and Reliability of the Japanese Version of the Selected Anger Expression Scale and Age, Sex, Occupation and Regional Differences in Anger Expression Among Japanese. Journal of Epidemiology, 2000, 10, 118-123.	2.4	8
192	Impact of anger expression on blood pressure levels in white-color workers with low-coping behavior. Environmental Health and Preventive Medicine, 2000, 5, 37-42.	3.4	8