Cyril C Curtain

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/935674/cyril-c-curtain-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

3,169 48 50 22 h-index g-index citations papers 3,328 50 3.4 3.94 avg, IF L-index ext. citations ext. papers

| # | Paper | IF | Citations |
|----|--|-----|-----------|
| 48 | Apolipoprotein C-II Adopts Distinct Structures in Complex with Micellar and Submicellar Forms of the Amyloid-Inhibiting Lipid-Mimetic Dodecylphosphocholine. <i>Biophysical Journal</i> , 2016 , 110, 85-94 | 2.9 | 2 |
| 47 | Stabilization of nontoxic Albligomers: insights into the mechanism of action of hydroxyquinolines in Alzheimer's disease. <i>Journal of Neuroscience</i> , 2015 , 35, 2871-84 | 6.6 | 56 |
| 46 | Small angle X-ray scattering analysis of Cu(2+)-induced oligomers of the Alzheimer's amyloid peptide. <i>Metallomics</i> , 2015 , 7, 536-43 | 4.5 | 19 |
| 45 | Alpha-synuclein oligomers and fibrils originate in two distinct conformer pools: a small angle X-ray scattering and ensemble optimisation modelling study. <i>Molecular BioSystems</i> , 2015 , 11, 190-6 | | 22 |
| 44 | Dopamine-Induced Esynuclein Oligomers 2014 , 291-300 | | |
| 43 | Guanidine hydrochloride denaturation of dopamine-induced Esynuclein oligomers: a small-angle X-ray scattering study. <i>Proteins: Structure, Function and Bioinformatics</i> , 2014 , 82, 10-21 | 4.2 | 7 |
| 42 | Ammonium hydroxide treatment of Alþroduces an aggregate free solution suitable for biophysical and cell culture characterization. <i>PeerJ</i> , 2013 , 1, e73 | 3.1 | 72 |
| 41 | Stereospecific interactions are necessary for Alzheimer disease amyloid-ltoxicity. <i>Neurobiology of Aging</i> , 2011 , 32, 235-48 | 5.6 | 43 |
| 40 | Histidine 14 modulates membrane binding and neurotoxicity of the Alzheimer's disease amyloid-beta peptide. <i>Journal of Alzheimerts Disease</i> , 2010 , 19, 1387-400 | 4.3 | 27 |
| 39 | The structure of dopamine induced alpha-synuclein oligomers. <i>European Biophysics Journal</i> , 2010 , 39, 1407-19 | 1.9 | 76 |
| 38 | Membrane-targeted strategies for modulating APP and Abeta-mediated toxicity. <i>Journal of Cellular and Molecular Medicine</i> , 2009 , 13, 249-61 | 5.6 | 3 |
| 37 | We see what we are trained to see, or must we? Some personal lessons from a brush with kuru research. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2008 , 363, 3633-4 | 5.8 | 1 |
| 36 | Twenty years of metallo-neurobiology: where to now?. European Biophysics Journal, 2008, 37, 241-5 | 1.9 | 49 |
| 35 | Applications of electron paramagnetic resonance to studies of neurological disease. <i>European Biophysics Journal</i> , 2008 , 37, 281-94 | 1.9 | 7 |
| 34 | Structural studies of the Alzheimer's amyloid precursor protein copper-binding domain reveal how it binds copper ions. <i>Journal of Molecular Biology</i> , 2007 , 367, 148-61 | 6.5 | 85 |
| 33 | Copper Coordination by EAmyloid and the Neuropathology of Alzheimer Disease 2007, 125-141 | | |
| 32 | Free Radicals, Metal Ions, and Alaggregation and Neurotoxicity 2007 , 31-47 | | 3 |

(1998-2006)

| 31 | Copper-mediated amyloid-beta toxicity is associated with an intermolecular histidine bridge. <i>Journal of Biological Chemistry</i> , 2006 , 281, 15145-54 | 5.4 | 150 |
|----|---|-----|-----|
| 30 | Copper-dependent inhibition of human cytochrome c oxidase by a dimeric conformer of amyloid-beta1-42. <i>Journal of Neuroscience</i> , 2005 , 25, 672-9 | 6.6 | 291 |
| 29 | Methylation of the imidazole side chains of the Alzheimer disease amyloid-beta peptide results in abolition of superoxide dismutase-like structures and inhibition of neurotoxicity. <i>Journal of Biological Chemistry</i> , 2005 , 280, 13355-63 | 5.4 | 101 |
| 28 | Dopamine promotes alpha-synuclein aggregation into SDS-resistant soluble oligomers via a distinct folding pathway. <i>FASEB Journal</i> , 2005 , 19, 1377-9 | 0.9 | 217 |
| 27 | Enhanced toxicity and cellular binding of a modified amyloid beta peptide with a methionine to valine substitution. <i>Journal of Biological Chemistry</i> , 2004 , 279, 42528-34 | 5.4 | 92 |
| 26 | Tyrosine gated electron transfer is key to the toxic mechanism of Alzheimer's disease beta-amyloid. <i>FASEB Journal</i> , 2004 , 18, 1427-9 | 0.9 | 231 |
| 25 | Neurotoxic, redox-competent Alzheimer's beta-amyloid is released from lipid membrane by methionine oxidation. <i>Journal of Biological Chemistry</i> , 2003 , 278, 42959-65 | 5.4 | 156 |
| 24 | Cu2+-induced modification of the kinetics of A beta(1-42) channels. <i>American Journal of Physiology - Cell Physiology</i> , 2003 , 285, C873-80 | 5.4 | 16 |
| 23 | Methionine oxidation: Implications for the mechanism of toxicity of the Emmyloid peptide from Alzheimer's disease. <i>International Journal of Peptide Research and Therapeutics</i> , 2003 , 10, 413-417 | | 13 |
| 22 | Structure of the Alzheimer's disease amyloid precursor protein copper binding domain. A regulator of neuronal copper homeostasis. <i>Journal of Biological Chemistry</i> , 2003 , 278, 17401-7 | 5.4 | 208 |
| 21 | Methionine oxidation: Implications for the mechanism of toxicity of the Emmyloid peptide from Alzheimer's disease. <i>International Journal of Peptide Research and Therapeutics</i> , 2003 , 10, 413-417 | 2.1 | 3 |
| 20 | Metal ions, pH, and cholesterol regulate the interactions of Alzheimer's disease amyloid-beta peptide with membrane lipid. <i>Journal of Biological Chemistry</i> , 2003 , 278, 2977-82 | 5.4 | 171 |
| 19 | Magnetic Resonance Studies of EAmyloid Peptides. Australian Journal of Chemistry, 2003, 56, 349 | 1.2 | 19 |
| 18 | Copper and zinc binding modulates the aggregation and neurotoxic properties of the prion peptide PrP106-126. <i>Biochemistry</i> , 2001 , 40, 8073-84 | 3.2 | 247 |
| 17 | Alzheimer's disease amyloid-beta binds copper and zinc to generate an allosterically ordered membrane-penetrating structure containing superoxide dismutase-like subunits. <i>Journal of Biological Chemistry</i> , 2001 , 276, 20466-73 | 5.4 | 530 |
| 16 | Selective, high-affinity binding of ferric ions by glycine-extended gastrin(17). <i>Biochemistry</i> , 2001 , 40, 10741-6 | 3.2 | 32 |
| 15 | Residues within the HFRIGC sequence of HIV-1 vpr involved in growth arrest activities. <i>Biochemical and Biophysical Research Communications</i> , 1999 , 264, 287-90 | 3.4 | 17 |
| 14 | Efficacy of fusion peptide homologs in blocking cell lysis and HIV-induced fusion. <i>AIDS Research and Human Retroviruses</i> , 1998 , 14, 385-92 | 1.6 | 9 |

| 13 | Structural requirements for the cytotoxicity of the N-terminal region of HIV type 1 Nef. <i>AIDS Research and Human Retroviruses</i> , 1998 , 14, 1543-51 | 1.6 | 5 |
|----|--|-----|----|
| 12 | Cytotoxicity resulting from addition of HIV-1 Nef N-terminal peptides to yeast and bacterial cells. <i>Biochemical and Biophysical Research Communications</i> , 1997 , 232, 707-11 | 3.4 | 13 |
| 11 | Preparation and characterization of a biologically active spin-labeled sea anemone toxin. <i>The Protein Journal</i> , 1996 , 15, 427-34 | | 4 |
| 10 | Antivirals that target the amino-terminal domain of HIV type 1 glycoprotein 41. <i>AIDS Research and Human Retroviruses</i> , 1995 , 11, 677-86 | 1.6 | 18 |
| 9 | The amino-terminal peptide of HIV-1 glycoprotein 41 fuses human erythrocytes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 1995 , 1271, 304-14 | 6.9 | 28 |
| 8 | Fusogenic activity of amino-terminal region of HIV type 1 Nef protein. <i>AIDS Research and Human Retroviruses</i> , 1994 , 10, 1231-40 | 1.6 | 24 |
| 7 | The amino-terminal peptide of HIV-1 gp41 interacts with human serum albumin. <i>AIDS Research and Human Retroviruses</i> , 1993 , 9, 1145-56 | 1.6 | 22 |
| 6 | Structure and dynamics of microemulsions which mimic the lipid phase of low-density lipoproteins. <i>Lipids and Lipid Metabolism</i> , 1990 , 1042, 42-50 | | 9 |
| 5 | Fatty-acid spin probe interactions with erythrocyte ghosts and liposomes prepared from erythrocyte ghosts. <i>Journal of Membrane Biology</i> , 1989 , 111, 155-68 | 2.3 | 13 |
| 4 | Thermotropic lipid phase separation in the human immunodeficiency virus. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1988 , 943, 331-42 | 3.8 | 20 |
| 3 | Electron spin resonance spectroscopy in the study of lymphoid cell receptors. <i>Methods in Enzymology</i> , 1987 , 150, 418-46 | 1.7 | 8 |
| 2 | Estimation of spin probe clustering in biological membranes. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1987 , 898, 202-13 | 3.8 | 6 |
| 1 | Spin probe clustering in human erythrocyte ghosts. <i>Journal of Membrane Biology</i> , 1985 , 84, 81-95 | 2.3 | 24 |