

Birgitte Fagerlund

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9358534/publications.pdf>

Version: 2024-02-01

129
papers

2,491
citations

185998

28
h-index

253896

43
g-index

137
all docs

137
docs citations

137
times ranked

4167
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | The relative and interactive impact of multiple risk factors in schizophrenia spectrum disorders: a combined register-based and clinical twin study. <i>Psychological Medicine</i> , 2023, 53, 1266-1276. | 2.7 | 8 |
| 2 | Reward Processing as an Indicator of Vulnerability or Compensatory Resilience in Psychoses? Results From a Twin Study. <i>Biological Psychiatry Global Open Science</i> , 2023, 3, 47-55. | 1.0 | 3 |
| 3 | Effects of methylphenidate on subjective sleep parameters in adults with ADHD: a prospective, non-randomized, non-blinded 6-week trial. <i>Nordic Journal of Psychiatry</i> , 2023, 77, 102-107. | 0.7 | 2 |
| 4 | Heritability of specific cognitive functions and associations with schizophrenia spectrum disorders using CANTAB: a nation-wide twin study. <i>Psychological Medicine</i> , 2022, 52, 1101-1114. | 2.7 | 18 |
| 5 | Early Antipsychotic Nonresponse as a Predictor of Nonresponse and Nonremission in Adolescents With Psychosis Treated With Aripiprazole or Quetiapine: Results From the TEA Trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2022, , . | 0.3 | 7 |
| 6 | Effects of prenatal nutrient supplementation and early life exposures on neurodevelopment at age 10: a randomised controlled trial - the COPSYPH study protocol. <i>BMJ Open</i> , 2022, 12, e047706. | 0.8 | 4 |
| 7 | A nation-wide twin study of social cognition in schizophrenia spectrum disorders. <i>NPJ Schizophrenia</i> , 2022, 8, 12. | 2.0 | 2 |
| 8 | Add-On MEmantine to Dopamine Antagonism to Improve Negative Symptoms at First Psychosis- the AMEND Trial Protocol. <i>Frontiers in Psychiatry</i> , 2022, 13, . | 1.3 | 3 |
| 9 | Cortico-cognition coupling in treatment resistant schizophrenia. <i>NeuroImage: Clinical</i> , 2022, 35, 103064. | 1.4 | 4 |
| 10 | Subclinical cognitive deficits are associated with reduced cerebrovascular response to visual stimulation in mid-sixties men. <i>GeroScience</i> , 2022, 44, 1905-1923. | 2.1 | 8 |
| 11 | Differential effects of age at illness onset on verbal memory functions in antipsychotic-naïve schizophrenia patients aged 12-43 years. <i>Psychological Medicine</i> , 2021, 51, 1570-1580. | 2.7 | 17 |
| 12 | Multimodal assessment of white matter microstructure in antipsychotic-naïve schizophrenia patients and confounding effects of recreational drug use. <i>Brain Imaging and Behavior</i> , 2021, 15, 36-48. | 1.1 | 6 |
| 13 | Associations Between Cognitive Function and Levels of Glutamatergic Metabolites and Gamma-Aminobutyric Acid in Antipsychotic-Naïve Patients With Schizophrenia or Psychosis. <i>Biological Psychiatry</i> , 2021, 89, 278-287. | 0.7 | 36 |
| 14 | Generalized neurocognitive impairment in individuals at ultra-high risk for psychosis: The possible key role of slowed processing speed. <i>Brain and Behavior</i> , 2021, 11, e01962. | 1.0 | 10 |
| 15 | Fish Oil Supplementation in Pregnancy and Neurodevelopment in Childhood—A Randomized Clinical Trial. <i>Child Development</i> , 2021, 92, 1624-1635. | 1.7 | 6 |
| 16 | Association of neurocognitive functioning with sleep stage dissociation and REM sleep instability in medicated patients with schizophrenia. <i>Journal of Psychiatric Research</i> , 2021, 136, 198-203. | 1.5 | 3 |
| 17 | Effects of methylphenidate on sensory and sensorimotor gating of initially psychostimulant-naïve adult ADHD patients. <i>European Neuropsychopharmacology</i> , 2021, 46, 83-92. | 0.3 | 4 |
| 18 | Associations between cognition and white matter microstructure in first-episode antipsychotic-naïve patients with schizophrenia and healthy controls: A multivariate pattern analysis. <i>Cortex</i> , 2021, 139, 282-297. | 1.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Effects of methylphenidate on mismatch negativity and P3a amplitude of initially psychostimulant-naïve, adult ADHD patients. <i>Psychological Medicine</i> , 2021, , 1-9. | 2.7 | 3 |
| 20 | Global fractional anisotropy predicts transition to psychosis after 12 months in individuals at ultra-high risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2021, 144, 448-463. | 2.2 | 9 |
| 21 | Changes in negative symptoms are linked to white matter changes in superior longitudinal fasciculus in individuals at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2021, 237, 192-201. | 1.1 | 6 |
| 22 | CYP2D6 Genotyping and Antipsychotic-Associated Extrapyramidal Adverse Effects in a Randomized Trial of Aripiprazole Versus Quetiapine Extended Release in Children and Adolescents, Aged 12–17 Years, With First Episode Psychosis. <i>Journal of Clinical Psychopharmacology</i> , 2021, 41, 667-672. | 0.7 | 2 |
| 23 | Do young adolescents with first-episode psychosis or ADHD show sensorimotor gating deficits?. <i>Psychological Medicine</i> , 2020, 50, 607-615. | 2.7 | 6 |
| 24 | Cognitive reserve attenuates age-related cognitive decline in the context of putatively accelerated brain ageing in schizophrenia-spectrum disorders. <i>Psychological Medicine</i> , 2020, 50, 1475-1489. | 2.7 | 12 |
| 25 | Auditory sensory gating in young adolescents with early-onset psychosis: a comparison with attention deficit/hyperactivity disorder. <i>Neuropsychopharmacology</i> , 2020, 45, 649-655. | 2.8 | 6 |
| 26 | Measuring movements in adolescents with psychosis using the Microsoft Kinect sensor: a pilot study exploring a new tool for assessing aspects of antipsychotic-induced parkinsonism. <i>Child and Adolescent Mental Health</i> , 2020, 25, 79-94. | 1.8 | 0 |
| 27 | Cerebral Glutamate and Gamma-Aminobutyric Acid Levels in Individuals at Ultra-high Risk for Psychosis and the Association With Clinical Symptoms and Cognition. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2020, 5, 569-579. | 1.1 | 12 |
| 28 | T46. THE EFFECT OF COMPREHENSIVE COGNITIVE REMEDIATION IN INDIVIDUALS AT ULTRA-HIGH RISK FOR PSYCHOSIS: A SINGLE-BLIND, RANDOMISED, CLINICAL TRIAL (FOCUS). <i>Schizophrenia Bulletin</i> , 2020, 46, S249-S249. | 2.3 | 0 |
| 29 | T60. GENETIC INFLUENCES ON MEMORY FUNCTIONS AND RELATED BRAIN STRUCTURES AND ASSOCIATIONS WITH SCHIZOPHRENIA SPECTRUM DISORDERS: A NATION-WIDE TWIN STUDY. <i>Schizophrenia Bulletin</i> , 2020, 46, S254-S254. | 2.3 | 0 |
| 30 | Effectiveness of cognitive remediation in the ultra-high risk state for psychosis. <i>World Psychiatry</i> , 2020, 19, 401-402. | 4.8 | 18 |
| 31 | Cognitive remediation plus standard treatment versus standard treatment alone for individuals at ultra-high risk of developing psychosis: Results of the FOCUS randomised clinical trial. <i>Schizophrenia Research</i> , 2020, 224, 151-158. | 1.1 | 23 |
| 32 | No Effects of Cognitive Remediation on Cerebral White Matter in Individuals at Ultra-High Risk for Psychosis—A Randomized Clinical Trial. <i>Frontiers in Psychiatry</i> , 2020, 11, 873. | 1.3 | 9 |
| 33 | Baseline measures of cerebral glutamate and GABA levels in individuals at ultrahigh risk for psychosis: Implications for clinical outcome after 12 months. <i>European Psychiatry</i> , 2020, 63, e83. | 0.1 | 7 |
| 34 | Interview and questionnaire assessment of cognitive impairment in subjects at ultra-high risk for psychosis: Associations with cognitive test performance, psychosocial functioning, and positive symptoms. <i>Psychiatry Research</i> , 2020, 294, 113498. | 1.7 | 3 |
| 35 | A machine-learning framework for robust and reliable prediction of short- and long-term treatment response in initially antipsychotic-naïve schizophrenia patients based on multimodal neuropsychiatric data. <i>Translational Psychiatry</i> , 2020, 10, 276. | 2.4 | 24 |
| 36 | Heritability of Memory Functions and Related Brain Volumes: A Schizophrenia Spectrum Study of 214 Twins. <i>Schizophrenia Bulletin Open</i> , 2020, 1, . | 0.9 | 3 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Identification of a Serotonin 2A Receptor Subtype of Schizophrenia Spectrum Disorders With Pimavanserin: The Sub-Sero Proof-of-Concept Trial Protocol. <i>Frontiers in Pharmacology</i> , 2020, 11, 591. | 1.6 | 8 |
| 38 | Associations between facial affect recognition and neurocognition in subjects at ultra-high risk for psychosis: A case-control study. <i>Psychiatry Research</i> , 2020, 290, 112969. | 1.7 | 6 |
| 39 | Discovering correlates of age-related decline in a healthy late-midlife male birth cohort. <i>Aging</i> , 2020, 12, 16709-16743. | 1.4 | 2 |
| 40 | Cortical structures and their clinical correlates in antipsychotic-naïve schizophrenia patients before and after 6 weeks of dopamine D _{2/3} receptor antagonist treatment. <i>Psychological Medicine</i> , 2019, 49, 754-763. | 2.7 | 19 |
| 41 | Diagnostic stability and long-term symptomatic and functional outcomes in first-episode antipsychotic-naïve patients with schizophrenia. <i>European Psychiatry</i> , 2019, 62, 130-137. | 0.1 | 8 |
| 42 | T88. THE IMPACT OF AGE OF ONSET AND ILLNESS DURATION ON WHITE MATTER AND COGNITION TRAJECTORIES IN SCHIZOPHRENIA: A 7-YEAR FOLLOW-UP STUDY ACROSS MULTIPLE TIME-POINTS. <i>Schizophrenia Bulletin</i> , 2019, 45, S237-S238. | 2.3 | 0 |
| 43 | T97. PATTERNS OF COGNITIVE FUNCTION ARE UNIQUELY ASSOCIATED WITH WHITE MATTER-MICROSTRUCTURE IN INDIVIDUALS AT ULTRA-HIGH RISK FOR PSYCHOSIS. <i>Schizophrenia Bulletin</i> , 2019, 45, S241-S242. | 2.3 | 0 |
| 44 | S15. HERITABILITY AND CORRELATION TO SCHIZOPHRENIA SPECTRUM DISORDERS OF CEREBRAL BLOOD FLOW MEASURED BY PSEUDO-CONTINUOUS ARTERIAL SPIN LABELING IN DANISH TWINS. <i>Schizophrenia Bulletin</i> , 2019, 45, S311-S311. | 2.3 | 0 |
| 45 | O4.5. EXPLORING THE RELATIONSHIP BETWEEN SENSORY FILTERING AND COGNITION: RESULTS OF A LARGE COHORT OF ANTIPSYCHOTIC-NAÏVE FIRST-EPISEODE SCHIZOPHRENIA PATIENTS. <i>Schizophrenia Bulletin</i> , 2019, 45, S170-S171. | 2.3 | 1 |
| 46 | O12.1. CARDIOMETABOLIC ADVERSE EFFECTS AND ITS PREDICTORS IN CHILDREN AND ADOLESCENTS WITH FIRST-EPISEODE PSYCHOSIS DURING TREATMENT WITH QUETIAPINE-ER VERSUS ARIPIPRAZOLE: 12-WEEK RESULTS: FROM THE TEA TRIAL. <i>Schizophrenia Bulletin</i> , 2019, 45, S197-S197. | 2.3 | 0 |
| 47 | Widespread higher fractional anisotropy associates to better cognitive functions in individuals at ultra-high risk for psychosis. <i>Human Brain Mapping</i> , 2019, 40, 5185-5201. | 1.9 | 22 |
| 48 | 4.1 COGNITIVE RESERVE ATTENUATES AGE-RELATED COGNITIVE DECLINE IN THE CONTEXT OF ACCELERATED BRAIN AGEING IN SCHIZOPHRENIA-SPECTRUM DISORDERS: EVIDENCE FOR ACTIVE COMPENSATION. <i>Schizophrenia Bulletin</i> , 2019, 45, S91-S92. | 2.3 | 1 |
| 49 | Attenuated mismatch negativity in patients with first-episode antipsychotic-naive schizophrenia using a source-resolved method. <i>Neurolmage: Clinical</i> , 2019, 22, 101760. | 1.4 | 10 |
| 50 | O4.6. HERITABILITY OF SPECIFIC COGNITIVE FUNCTIONS AND ASSOCIATIONS WITH SCHIZOPHRENIA SPECTRUM DISORDERS USING CANTAB: A NATION-WIDE TWIN STUDY. <i>Schizophrenia Bulletin</i> , 2019, 45, S171-S171. | 2.3 | 1 |
| 51 | Cardiometabolic Adverse Effects and Its Predictors in Children and Adolescents With First-Episode Psychosis During Treatment With Quetiapine-Extended Release Versus Aripiprazole: 12-Week Results From the Tolerance and Effect of Antipsychotics in Children and Adolescents With Psychosis (TEA) Trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2019, 58, 1062-1078. | 0.3 | 16 |
| 52 | Heritability of Cerebral Blood Flow and the Correlation to Schizophrenia Spectrum Disorders: A Pseudo-continuous Arterial Spin Labeling Twin Study. <i>Schizophrenia Bulletin</i> , 2019, 45, 1231-1241. | 2.3 | 16 |
| 53 | Accuracy of diagnostic classification algorithms using cognitive-, electrophysiological-, and neuroanatomical data in antipsychotic-naïve schizophrenia patients. <i>Psychological Medicine</i> , 2019, 49, 2754-2763. | 2.7 | 20 |
| 54 | Visual attention in adults with attention-deficit/hyperactivity disorder before and after stimulant treatment. <i>Psychological Medicine</i> , 2019, 49, 2617-2625. | 2.7 | 8 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Emotion recognition latency, but not accuracy, relates to real life functioning in individuals at ultra-high risk for psychosis. <i>Schizophrenia Research</i> , 2019, 210, 197-202. | 1.1 | 13 |
| 56 | Sleep efficiency and neurophysiological patterns in middle-aged men are associated with cognitive change over their adult life course. <i>Journal of Sleep Research</i> , 2019, 28, e12793. | 1.7 | 8 |
| 57 | Heritability of cerebral glutamate levels and their association with schizophrenia spectrum disorders: a ¹ [H]-spectroscopy twin study. <i>Neuropsychopharmacology</i> , 2019, 44, 581-589. | 2.8 | 28 |
| 58 | The impact of schizophrenia and intelligence on the relationship between age and brain volume. <i>Schizophrenia Research: Cognition</i> , 2019, 15, 1-6. | 0.7 | 8 |
| 59 | Patterns of Cortical Structures and Cognition in Antipsychotic-Naïve Patients With First-Episode Schizophrenia: A Partial Least Squares Correlation Analysis. <i>Biological Psychiatry: Cognitive Neuroscience and Neuroimaging</i> , 2019, 4, 444-453. | 1.1 | 12 |
| 60 | Hyper- and Hypomentalyzing in Patients with First-Episode Schizophrenia: fMRI and Behavioral Studies. <i>Schizophrenia Bulletin</i> , 2019, 45, 377-385. | 2.3 | 38 |
| 61 | Investigation of sleep spindle activity and morphology as predictors of neurocognitive functioning in medicated patients with schizophrenia. <i>Journal of Sleep Research</i> , 2019, 28, e12672. | 1.7 | 13 |
| 62 | Associations between P3a and P3b amplitudes and cognition in antipsychotic-naïve first-episode schizophrenia patients. <i>Psychological Medicine</i> , 2019, 49, 868-875. | 2.7 | 18 |
| 63 | Discovering markers of healthy aging: a prospective study in a Danish male birth cohort. <i>Aging</i> , 2019, 11, 5943-5974. | 1.4 | 11 |
| 64 | Overlapping and disease specific trait, response, and reflection impulsivity in adolescents with first-episode schizophrenia spectrum disorders or attention-deficit/hyperactivity disorder. <i>Psychological Medicine</i> , 2018, 48, 604-616. | 2.7 | 14 |
| 65 | Heritability of Schizophrenia and Schizophrenia Spectrum Based on the Nationwide Danish Twin Register. <i>Biological Psychiatry</i> , 2018, 83, 492-498. | 0.7 | 374 |
| 66 | Change and dispersion of QT interval during treatment with quetiapine extended release versus aripiprazole in children and adolescents with first-episode psychosis: results from the TEA trial. <i>Psychopharmacology</i> , 2018, 235, 681-693. | 1.5 | 7 |
| 67 | Examining speed of processing of facial emotion recognition in individuals at ultra-high risk for psychosis: Associations with symptoms and cognition. <i>Schizophrenia Research</i> , 2018, 195, 562-563. | 1.1 | 5 |
| 68 | O3.3. REWARD PROCESSING AS A VULNERABILITY INDICATOR FOR PSYCHOSIS: RESULTS FROM A TWIN STUDY. <i>Schizophrenia Bulletin</i> , 2018, 44, S80-S80. | 2.3 | 0 |
| 69 | F21. ELECTROPHYSIOLOGICAL PARAMETERS OF SELECTIVE ATTENTION IN ADOLESCENTS WITH A FIRST EPISODE OF PSYCHOSIS: A COMPARISON WITH ADHD. <i>Schizophrenia Bulletin</i> , 2018, 44, S226-S227. | 2.3 | 0 |
| 70 | F34. AUDITORY SENSORY GATING IN YOUNG ADOLESCENTS WITH EARLY-ONSET PSYCHOSIS: A COMPARISON WITH ADHD. <i>Schizophrenia Bulletin</i> , 2018, 44, S232-S232. | 2.3 | 0 |
| 71 | F176. CLINICAL CORRELATES OF CORTICAL STRUCTURE IN ANTIPSYCHOTIC-NAÏVE SCHIZOPHRENIA PATIENTS BEFORE AND AFTER SIX-WEEK TREATMENT WITH A DOPAMINE D2/3 RECEPTOR ANTAGONIST. <i>Schizophrenia Bulletin</i> , 2018, 44, S289-S289. | 2.3 | 0 |
| 72 | Alterations of Intrinsic Connectivity Networks in Antipsychotic-Naïve First-Episode Schizophrenia. <i>Schizophrenia Bulletin</i> , 2018, 44, 1332-1340. | 2.3 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Delay Aversion and Executive Functioning in Adults With Attention-Deficit/Hyperactivity Disorder: Before and After Stimulant Treatment. <i>International Journal of Neuropsychopharmacology</i> , 2018, 21, 997-1006. | 1.0 | 11 |
| 74 | No cognitive-enhancing effect of GLP-1 receptor agonism in antipsychotic-treated, obese patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 52-62. | 2.2 | 36 |
| 75 | Two subgroups of antipsychotic-naive, first-episode schizophrenia patients identified with a Gaussian mixture model on cognition and electrophysiology. <i>Translational Psychiatry</i> , 2017, 7, e1087-e1087. | 2.4 | 32 |
| 76 | Is an Early Age at Illness Onset in Schizophrenia Associated With Increased Genetic Susceptibility? Analysis of Data From the Nationwide Danish Twin Register. <i>EBioMedicine</i> , 2017, 18, 320-326. | 2.7 | 22 |
| 77 | Quetiapine extended release versus aripiprazole in children and adolescents with first-episode psychosis: the multicentre, double-blind, randomised tolerability and efficacy of antipsychotics (TEA) trial. <i>Lancet Psychiatry</i> , 2017, 4, 605-618. | 3.7 | 40 |
| 78 | Mismatch negativity and P3a amplitude in young adolescents with first-episode psychosis: a comparison with ADHD. <i>Psychological Medicine</i> , 2017, 47, 377-388. | 2.7 | 30 |
| 79 | Negative symptoms mediate the relationship between neurocognition and function in individuals at ultrahigh risk for psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2017, 135, 250-258. | 2.2 | 25 |
| 80 | No cognitive-enhancing effect of GLP-1 receptor agonism in antipsychotic-treated, obese patients with schizophrenia: authors' response. <i>Acta Psychiatrica Scandinavica</i> , 2017, 136, 526-527. | 2.2 | 0 |
| 81 | 48. Associations Between Electrophysiological Measures of Selective Attention and Neurocognitive Measures of Working Memory and Attention in Antipsychotic-Naive, First-Episode Schizophrenia Patients. <i>Schizophrenia Bulletin</i> , 2017, 43, S26-S26. | 2.3 | 0 |
| 82 | Testing a decades-old assumption: Are individuals with lower sensory gating indeed more easily distracted?. <i>Psychiatry Research</i> , 2017, 255, 387-393. | 1.7 | 9 |
| 83 | The effect of positive symptoms on social cognition in first-episode schizophrenia is modified by the presence of negative symptoms. <i>Neuropsychology</i> , 2017, 31, 209-219. | 1.0 | 41 |
| 84 | Neurocognitive performance, subjective well-being, and psychosocial functioning after benzodiazepine withdrawal in patients with schizophrenia or bipolar disorder: a randomized clinical trial of add-on melatonin versus placebo. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2017, 267, 163-171. | 1.8 | 35 |
| 85 | Five years of specialised early intervention versus two years of specialised early intervention followed by three years of standard treatment for patients with a first episode psychosis: randomised, superiority, parallel group trial in Denmark (OPUS II). <i>BMJ: British Medical Journal</i> , 2017, 356, i6681. | 2.4 | 79 |
| 86 | SU18. GLP-1 Receptor Agonist Treatment in Schizophrenia Patients With Obesity. <i>Schizophrenia Bulletin</i> , 2017, 43, S167-S167. | 2.3 | 2 |
| 87 | Passive Double-Sensory Evoked Coherence Correlates with Long-Term Memory Capacity. <i>Frontiers in Human Neuroscience</i> , 2017, 11, 598. | 1.0 | 6 |
| 88 | SA37. Cognition and White Matter Integrity in Antipsychotic-Naive First-Episode Schizophrenia Patients. <i>Schizophrenia Bulletin</i> , 2017, 43, S126-S127. | 2.3 | 0 |
| 89 | Pretreatment Cardiometabolic Status in Youth With Early-Onset Psychosis. <i>Journal of Clinical Psychiatry</i> , 2017, 78, e1035-e1046. | 1.1 | 36 |
| 90 | Melatonin and cortisol profiles in late midlife and their association with age-related changes in cognition. <i>Nature and Science of Sleep</i> , 2016, 8, 47. | 1.4 | 28 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Cognitive Change during the Life Course and Leukocyte Telomere Length in Late Middle-Aged Men. <i>Frontiers in Aging Neuroscience</i> , 2016, 8, 300. | 1.7 | 10 |
| 92 | Are Weight Status and Cognition Associated? An Examination of Cognitive Development in Children and Adolescents with Anorexia Nervosa 1â€™%Year after First Hospitalisation. <i>European Eating Disorders Review</i> , 2016, 24, 366-376. | 2.3 | 17 |
| 93 | 4.17 QUETIAPINE EXTENDED RELEASE VERSUS ARIPIPRAZOLE IN CHILDREN AND ADOLESCENTS WITH PSYCHOSIS IN THE RANDOMIZED, BLINDED CLINICAL TOLERABILITY AND EFFICACY OF ANTIPSYCHOTICS (TEA) TRIAL. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2016, 55, S168. | 0.3 | 0 |
| 94 | Predictors and longitudinal course of cognitive functioning in schizophrenia spectrum disorders, 10 years after baseline: The OPUS study. <i>Schizophrenia Research</i> , 2016, 175, 57-63. | 1.1 | 61 |
| 95 | Social cognition in patients at ultra-high risk for psychosis: What is the relation to social skills and functioning?. <i>Schizophrenia Research: Cognition</i> , 2016, 5, 21-27. | 0.7 | 46 |
| 96 | Frontal D2/3Receptor Availability in Schizophrenia Patients Before and After Their First Antipsychotic Treatment: Relation to Cognitive Functions and Psychopathology. <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, pyw006. | 1.0 | 17 |
| 97 | Subjective sleep quality and daytime sleepiness in late midlife and their association with age-related changes in cognition. <i>Sleep Medicine</i> , 2016, 17, 165-173. | 0.8 | 49 |
| 98 | Influence of early life characteristics on psychiatric admissions and impact of psychiatric disease on inflammatory biomarkers and survival: a <scp>D</scp>anish cohort study. <i>World Psychiatry</i> , 2015, 14, 364-365. | 4.8 | 7 |
| 99 | The FOCUS trial: cognitive remediation plus standard treatment versus standard treatment for patients at ultra-high risk for psychosis: study protocol for a randomised controlled trial. <i>Trials</i> , 2015, 16, 25. | 0.7 | 40 |
| 100 | Cognitive Profile of Children and Adolescents with Anorexia Nervosa. <i>European Eating Disorders Review</i> , 2015, 23, 34-42. | 2.3 | 42 |
| 101 | Increased deoxythymidine triphosphate levels is a feature of relative cognitive decline. <i>Mitochondrion</i> , 2015, 25, 34-37. | 1.6 | 8 |
| 102 | Association Study of CHRNA7 Promoter Variants with Sensory and Sensorimotor Gating in Schizophrenia Patients and Healthy Controls: A Danish Caseâ€™Control Study. <i>NeuroMolecular Medicine</i> , 2015, 17, 423-430. | 1.8 | 23 |
| 103 | Cognitive function in idiopathic intracranial hypertension: a prospective caseâ€™control study. <i>BMJ Open</i> , 2014, 4, e004376. | 0.8 | 71 |
| 104 | Association of the CHRNA7 promoter variant â€™86T with Tourette syndrome and comorbid obsessive-compulsive disorder. <i>Psychiatry Research</i> , 2014, 219, 710-711. | 1.7 | 1 |
| 105 | Treatment of antipsychotic-associated obesity with a GLP-1 receptor agonistâ€™ protocol for an investigator-initiated prospective, randomised, placebo-controlled, double-blinded intervention study: the TAO study protocol. <i>BMJ Open</i> , 2014, 4, e004158. | 0.8 | 20 |
| 106 | Subclinical cognitive decline in middleâ€™age is associated with reduced taskâ€™induced deactivation of the brain's default mode network. <i>Human Brain Mapping</i> , 2014, 35, 4488-4498. | 1.9 | 51 |
| 107 | Quetiapine versus aripiprazole in children and adolescents with psychosis - protocol for the randomised, blinded clinical Tolerability and Efficacy of Antipsychotics (TEA) trial. <i>BMC Psychiatry</i> , 2014, 14, 199. | 1.1 | 10 |
| 108 | The Relationship Between Cognitive Ability and Demographic Factors in Late Midlife. <i>Journal of Aging and Health</i> , 2014, 26, 37-53. | 0.9 | 40 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Social cognition and neurocognitive deficits in first-episode schizophrenia. <i>Schizophrenia Research</i> , 2014, 153, 9-17. | 1.1 | 76 |
| 110 | P201: Late midlife sleep pattern and sleep structure and the association to age-related changes in cognition. <i>Clinical Neurophysiology</i> , 2014, 125, S101. | 0.7 | 0 |
| 111 | Cognitive function patients with idiopathic intracranial hypertension. <i>Journal of the Neurological Sciences</i> , 2013, 333, e483. | 0.3 | 0 |
| 112 | The influence of impaired processing speed on cognition in first-episode antipsychotic-naïve schizophrenic patients. <i>European Psychiatry</i> , 2013, 28, 332-339. | 0.1 | 38 |
| 113 | Relationship of frontal D2/3 binding potentials to cognition: a study of antipsychotic-naïve schizophrenia patients. <i>International Journal of Neuropsychopharmacology</i> , 2013, 16, 23-36. | 1.0 | 18 |
| 114 | Profile of cognitive deficits and associations with depressive symptoms and intelligence in chronic early-onset schizophrenia patients. <i>Scandinavian Journal of Psychology</i> , 2013, 54, 363-370. | 0.8 | 4 |
| 115 | Poster #64 RECRUITMENT STATUS OF THE TEA TRIAL: TOLERANCE AND EFFECT OF ANTIPSYCHOTICS IN CHILDREN AND ADOLESCENTS WITH PSYCHOSIS. AN INVESTIGATOR-INITIATED, PHASE IV, RANDOMISED DOUBLE-BLIND MULTI-CENTRE TRIAL OF THE BENEFITS AND HARMS OF ARIPIPRAZOLE VERSUS QUETIAPINE IN CHILDREN AND ADOLESCENTS WITH PSYCHOSIS. <i>Schizophrenia Research</i> , 2012, 136, S208. | 1.1 | 0 |
| 116 | Glucagon-like peptide-1 analogs against antipsychotic-induced weight gain: potential physiological benefits. <i>BMC Medicine</i> , 2012, 10, 92. | 2.3 | 24 |
| 117 | Stability of prepulse inhibition and habituation of the startle reflex in schizophrenia: a 6-year follow-up study of initially antipsychotic-naïve, first-episode schizophrenia patients. <i>International Journal of Neuropsychopharmacology</i> , 2011, 14, 913-925. | 1.0 | 31 |
| 118 | Cognitive effects of six months of treatment with quetiapine in antipsychotic-naïve first-episode schizophrenia. <i>Psychiatry Research</i> , 2011, 187, 49-54. | 1.7 | 18 |
| 119 | Prolonged-release melatonin versus placebo for benzodiazepine discontinuation in patients with schizophrenia: a randomized clinical trial - the SMART trial protocol. <i>BMC Psychiatry</i> , 2011, 11, 160. | 1.1 | 28 |
| 120 | Course of intelligence deficits in early onset, first episode schizophrenia: a controlled, 5-year longitudinal study. <i>European Child and Adolescent Psychiatry</i> , 2010, 19, 341-351. | 2.8 | 18 |
| 121 | Deficient maturation of aspects of attention and executive functions in early onset schizophrenia. <i>European Child and Adolescent Psychiatry</i> , 2010, 19, 773-786. | 2.8 | 25 |
| 122 | THE COURSE OF COGNITIVE DEFICITS IN SCHIZOPHRENIA FROM ILLNESS ONSET TO 6 YEARS POST ONSET: PRELIMINARY RESULTS FROM A PROSPECTIVE LONGITUDINAL STUDY. <i>Schizophrenia Research</i> , 2010, 117, 211. | 1.1 | 1 |
| 123 | Do Attention Deficits Influence IQ Assessment in Children and Adolescents With ADHD?. <i>Journal of Attention Disorders</i> , 2009, 12, 551-562. | 1.5 | 71 |
| 124 | EXTRASTRIATAL DOPAMINE D2 RECEPTOR BINDING POTENTIALS IN ANTIPSYCHOTIC-NAÏVE FIRST-EPISEODE SCHIZOPHRENIC PATIENTS. <i>Schizophrenia Research</i> , 2008, 102, 42. | 1.1 | 0 |
| 125 | Effects of Donepezil Adjunctive Treatment to Ziprasidone on Cognitive Deficits in Schizophrenia. <i>Clinical Neuropharmacology</i> , 2007, 30, 3-12. | 0.2 | 49 |
| 126 | Structural brain abnormalities in early onset first-episode psychosis. <i>Journal of Neural Transmission</i> , 2007, 114, 489-498. | 1.4 | 37 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Cognitive deficits and levels of IQ in adolescent onset schizophrenia and other psychotic disorders. Schizophrenia Research, 2006, 85, 30-39. | 1.1 | 61 |
| 128 | Effects of Low-Dose Risperidone and Low-Dose Zuclopenthixol on Cognitive Functions in First-Episode Drug-Naïve Schizophrenic Patients. CNS Spectrums, 2004, 9, 364-374. | 0.7 | 48 |
| 129 | Structural gray matter brain abnormalities in first-episode early onset psychosis: A voxel based morphometry study in children and adolescents aged ten to eighteen. Schizophrenia Research, 2003, 60, 204. | 1.1 | 0 |