

Mirosław Kozłowski

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9366084/miroslaw-kozlowski-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

55
papers

1,125
citations

16
h-index

31
g-index

60
ext. papers

1,260
ext. citations

3.7
avg, IF

3.79
L-index

#	Paper	IF	Citations
55	MicroRNA-30a inhibits epithelial-to-mesenchymal transition by targeting Snai1 and is downregulated in non-small cell lung cancer. <i>International Journal of Cancer</i> , 2012 , 130, 2044-53	7.5	241
54	An immune response enriched 72-gene prognostic profile for early-stage non-small-cell lung cancer. <i>Clinical Cancer Research</i> , 2009 , 15, 284-90	12.9	123
53	The diagnostic value of the measurement of matrix metalloproteinase 9 (MMP-9), squamous cell cancer antigen (SCC) and carcinoembryonic antigen (CEA) in the sera of esophageal cancer patients. <i>Clinica Chimica Acta</i> , 2008 , 389, 61-6	6.2	66
52	Whole genome sequencing puts forward hypotheses on metastasis evolution and therapy in colorectal cancer. <i>Nature Communications</i> , 2018 , 9, 4782	17.4	51
51	CXCL5 as a potential novel prognostic factor in early stage non-small cell lung cancer: results of a study of expression levels of 23 genes. <i>Tumor Biology</i> , 2014 , 35, 4619-28	2.9	41
50	Identification of the vasodilatory endothelial cannabinoid receptor in the human pulmonary artery. <i>Journal of Hypertension</i> , 2007 , 25, 2240-8	1.9	40
49	Higher importance of interleukin 6 than classic tumor markers (carcinoembryonic antigen and squamous cell cancer antigen) in the diagnosis of esophageal cancer patients. <i>Ecological Management and Restoration</i> , 2012 , 25, 242-9	3	33
48	Virodhamine relaxes the human pulmonary artery through the endothelial cannabinoid receptor and indirectly through a COX product. <i>British Journal of Pharmacology</i> , 2008 , 155, 1034-42	8.6	33
47	Prognostic significance of DAPK and RASSF1A promoter hypermethylation in non-small cell lung cancer (NSCLC). <i>Folia Histochemica Et Cytobiologica</i> , 2009 , 47, 275-80	1.4	30
46	Prognostic value of pretreatment CEA, SCC-Ag and CA 19-9 levels in sera of patients with non-small cell lung cancer. <i>European Journal of Cancer Prevention</i> , 1992 , 1, 401-6	2	28
45	Serum vascular endothelial growth factors C and D in patients with oesophageal cancer. <i>European Journal of Cardio-thoracic Surgery</i> , 2010 , 38, 260-7	3	26
44	The activity of class I, II, III, and IV alcohol dehydrogenase (ADH) isoenzymes and aldehyde dehydrogenase (ALDH) in esophageal cancer. <i>Digestive Diseases and Sciences</i> , 2009 , 54, 725-30	4	24
43	Mechanisms of endothelium-dependent relaxation evoked by anandamide in isolated human pulmonary arteries. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2014 , 387, 477-86	3.4	21
42	The Serum Concentrations of Chemokine CXCL12 and Its Specific Receptor CXCR4 in Patients with Esophageal Cancer. <i>Disease Markers</i> , 2016 , 2016, 7963895	3.2	21
41	Vascular endothelial growth factor C and D expression correlates with lymph node metastasis and poor prognosis in patients with resected esophageal cancer. <i>Neoplasma</i> , 2011 , 58, 311-9	3.3	20
40	Vasodilatory effects of cannabidiol in human pulmonary and rat small mesenteric arteries: modification by hypertension and the potential pharmacological opportunities. <i>Journal of Hypertension</i> , 2020 , 38, 896-911	1.9	20
39	Relaxation of human pulmonary arteries by PPAR α agonists. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2013 , 386, 445-53	3.4	16

38	Serum tissue inhibitor of metalloproteinase 1 (TIMP-1) and vascular endothelial growth factor A (VEGF-A) are associated with prognosis in esophageal cancer patients. <i>Advances in Medical Sciences</i> , 2013 , 58, 227-34	2.8	16
37	Serum soluble Fas ligand (sFasL) in patients with primary squamous cell carcinoma of the esophagus. <i>Folia Histochemica Et Cytobiologica</i> , 2007 , 45, 199-204	1.4	16
36	Comparison between clinical significance of serum CXCL-8 and classical tumor markers in oesophageal cancer (OC) patients. <i>Clinical and Experimental Medicine</i> , 2019 , 19, 191-199	4.9	15
35	The preoperative study of mediastinal lymph nodes metastasis in lung cancer by endoscopic ultrasonography (EUS) and helical computed tomography (CT). <i>Lung Cancer</i> , 2001 , 34 Suppl 2, S123-6	5.9	15
34	EUELC project: a multi-centre, multipurpose study to investigate early stage NSCLC, and to establish a biobank for ongoing collaboration. <i>European Respiratory Journal</i> , 2009 , 34, 1477-86	13.6	14
33	MMP-10, MMP-7, TIMP-1 and TIMP-2 mRNA expression in esophageal cancer. <i>Acta Biochimica Polonica</i> , 2017 , 64, 295-299	2	12
32	Comparative evaluation of serum C-reactive protein (CRP) levels in the different histological subtypes of esophageal cancer (squamous cell carcinoma and adenocarcinoma of esophagus). <i>Journal of Clinical Laboratory Analysis</i> , 2012 , 26, 73-81	3	12
31	Lymphatic vessel invasion detected by the endothelial lymphatic marker D2-40 (podoplanin) is predictive of regional lymph node status and an independent prognostic factor in patients with resected esophageal cancer. <i>Folia Histochemica Et Cytobiologica</i> , 2011 , 49, 90-7	1.4	12
30	Gene Expression Signature Differentiates Histology But Not Progression Status of Early-Stage NSCLC. <i>Translational Oncology</i> , 2017 , 10, 450-458	4.9	11
29	Systematic biobanking, novel imaging techniques, and advanced molecular analysis for precise tumor diagnosis and therapy: The Polish MOBIT project. <i>Advances in Medical Sciences</i> , 2017 , 62, 405-413	2.8	11
28	Ligands at beta2-, beta3-, and the low-affinity state of beta1-adrenoceptors block the alpha1-adrenoceptor-mediated constriction in human pulmonary and rat mesenteric arteries. <i>Journal of Cardiovascular Pharmacology</i> , 2005 , 46, 76-82	3.1	11
27	Validation for histology-driven diagnosis in non-small cell lung cancer using hsa-miR-205 and hsa-miR-21 expression by two different normalization strategies. <i>International Journal of Cancer</i> , 2016 , 138, 689-97	7.5	11
26	Clinical significance of serum macrophage-colony stimulating factor (M-CSF) in esophageal cancer patients and its comparison with classical tumor markers. <i>Clinical Chemistry and Laboratory Medicine</i> , 2010 , 48, 1467-73	5.9	10
25	Alcohol dehydrogenase isoenzymes and aldehyde dehydrogenase activity in the sera of patients with esophageal cancer. <i>Clinical and Experimental Medicine</i> , 2009 , 9, 131-7	4.9	10
24	Modified schrödinger equation for attosecond laser pulse interaction with matter. <i>Lasers in Engineering</i> , 2002 , 12, 53-58		10
23	Development of LC-QTOF-MS method for human lung tissue fingerprinting. A preliminary application to nonsmall cell lung cancer. <i>Electrophoresis</i> , 2017 , 38, 2304-2312	3.6	9
22	Prognostic value of vascular endothelial growth factor-C and podoplanin mRNA expression in esophageal cancer. <i>Oncology Letters</i> , 2015 , 10, 3668-3674	2.6	9
21	Increased MET Gene Copy Number but Not mRNA Level Predicts Postoperative Recurrence in Patients with Non-Small Cell Lung Cancer. <i>Translational Oncology</i> , 2014 , 7, 605-12	4.9	9

20	Squamous cell carcinoma antigen in patients with cancer of the larynx. <i>Clinical Otolaryngology</i> , 1999 , 24, 270-3		9
19	Expression of matrix metalloproteinase-9 in the neoplastic and interstitial inflammatory infiltrate cells in the different histopathological types of esophageal cancer. <i>Folia Histochemica Et Cytobiologica</i> , 2008 , 46, 471-8	1.4	8
18	Prognostic significance of Notch ligands in patients with non-small cell lung cancer. <i>Oncology Letters</i> , 2017 , 13, 506-510	2.6	7
17	Elevated levels of serum metalloproteinase 9 in patients with esophageal squamous cell carcinoma 2009 , 119, 558-63		7
16	New monoallelic combination of KRAS gene mutations in codons 12 and 13 in the lung adenocarcinoma. <i>Advances in Medical Sciences</i> , 2013 , 58, 83-9	2.8	6
15	Relationships between level of lipid peroxidation products and expression of Nrf2 and its activators/ inhibitors in non-small cell lung cancer tissue. <i>Free Radical Biology and Medicine</i> , 2014 , 75 Suppl 1, S31	7.8	6
14	Serum matrix metalloproteinase 2 and tissue inhibitor of matrix metalloproteinases 2 in esophageal cancer patients. <i>Folia Histochemica Et Cytobiologica</i> , 2012 , 50, 590-598	1.4	6
13	EP α receptor-mediated contraction of human pulmonary arteries and inhibition of neurogenic tachycardia in pithed rats. <i>Pharmacological Reports</i> , 2012 , 64, 1526-36	3.9	5
12	Serum concentrations of receptor for interleukin 8 in patients with esophageal cancer. <i>Polish Archives of Internal Medicine</i> , 2016 , 126, 854-861	1.9	5
11	Lymphatics-associated genes are downregulated at transcription level in non-small cell lung cancer. <i>Oncology Letters</i> , 2018 , 15, 6752-6762	2.6	4
10	Stem cell factor in the serum of patients with esophageal cancer in relation to its histological types. <i>Archives of Medical Science</i> , 2017 , 13, 1357-1364	2.9	3
9	The alcohol dehydrogenase isoenzyme (ADH IV) as a candidate tumour marker of esophageal cancer.. <i>Acta Biochimica Polonica</i> , 2013 , 60,	2	3
8	Reduction of the serotonin 5-HT and 5-HT receptor-mediated contraction of human pulmonary artery by the combined 5-HT receptor antagonist and serotonin transporter inhibitor LY393558. <i>Pharmacological Reports</i> , 2020 , 72, 756-762	3.9	2
7	Activity of Alcohol Dehydrogenase and Aldehyde Dehydrogenase in Lung Cancer Cells. <i>Anticancer Research</i> , 2020 , 40, 3857-3863	2.3	2
6	Significance of EGFR signaling pathway genetic alterations in radically resected non-small cell lung cancers from a Polish cohort. One institutional study. <i>Advances in Medical Sciences</i> , 2015 , 60, 277-86	2.8	1
5	GPR18-Mediated Relaxation of Human Isolated Pulmonary Arteries.. <i>International Journal of Molecular Sciences</i> , 2022 , 23,	6.3	1
4	Histone deacetylases affect transcriptional regulation of CCL2 and CXCL8 expression by pulmonary fibroblasts in vitro. <i>Advances in Respiratory Medicine</i> , 2017 , 85, 307-312	0.8	1
3	The Ability of Metabolomics to Discriminate Non-Small-Cell Lung Cancer Subtypes Depends on the Stage of the Disease and the Type of Material Studied. <i>Cancers</i> , 2021 , 13,	6.6	1

- | | | |
|---|---|-----|
| 2 | P-172POLO-LIKE KINASE 1 AND 3 mRNA EXPRESSION IN EARLY-STAGE NON-SMALL CELL LUNG CANCER. <i>Interactive Cardiovascular and Thoracic Surgery</i> , 2015 , 21, S47-S48 | 1.8 |
| 1 | THORACIC SURGERY Intratumoral lymphatic vessel density, and intratumoral and peritumoral lymphatic vessel invasion as predictive factors of lymph node metastasis and prognostic factors in esophageal cancer. <i>Kardiochirurgia I Torakochirurgia Polska</i> , 2013 , 2, 120-129 | 0.3 |