Elizabeth I Hamelin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9366468/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Use of Diagnostic Ions for the Detection of Fentanyl Analogs in Human Matrices by LC–QTOF. Journal of the American Society for Mass Spectrometry, 2021, 32, 2852-2859.	1.2	6
2	Application of the fentanyl analog screening kit toward the identification of emerging synthetic opioids in human plasma and urine by LC-QTOF. Toxicology Letters, 2020, 320, 87-94.	0.4	16
3	Rapid, Sensitive, and Accurate Point-of-Care Detection of Lethal Amatoxins in Urine. Toxins, 2020, 12, 123.	1.5	19
4	Surveillance for Harmful Algal Bloom Events and Associated Human and Animal Illnesses — One Health Harmful Algal Bloom System, United States, 2016–2018. Morbidity and Mortality Weekly Report, 2020, 69, 1889-1894.	9.0	33
5	Designing traceable opioid material§ kits to improve laboratory testing during the U.S. opioid overdose crisis. Toxicology Letters, 2019, 317, 53-58.	0.4	13
6	Measurement of Microcystin and Nodularin Activity in Human Urine by Immunocapture-Protein Phosphatase 2A Assay. Toxins, 2019, 11, 729.	1.5	13
7	Determination of Fentanyl Analog Exposure Using Dried Blood Spots with LC–MS-MS. Journal of Analytical Toxicology, 2019, 43, 266-276.	1.7	26
8	Investigation of dried blood sampling with liquid chromatography tandem mass spectrometry to confirm human exposure to nerve agents. Analytica Chimica Acta, 2018, 1033, 100-107.	2.6	11
9	Detection of α-, β-, and γ-amanitin in urine by LC-MS/MS using 15N10-α-amanitin as the internal standard. Toxicon, 2018, 152, 71-77.	0.8	26
10	Quantification of Microcystin-LR in Human Urine by Immunocapture Liquid Chromatography Tandem Mass Spectrometry. Chemical Research in Toxicology, 2018, 31, 898-903.	1.7	15
11	Quantification of saxitoxin in human blood by ELISA. Toxicon, 2017, 133, 110-115.	0.8	26
12	Quantitation of fentanyl analogs in dried blood spots by flow-through desorption coupled to online solid phase extraction tandem mass spectrometry. Analytical Methods, 2017, 9, 3876-3883.	1.3	16
13	Evaluation of Multiple Blood Matrices for Assessment of Human Exposure to Nerve Agents. Journal of Analytical Toxicology, 2016, 40, 229-235.	1.7	10
14	Bridging the gap between sample collection and laboratory analysis: using dried blood spots to identify human exposure to chemical agents. , 2016, 98630, 98630P-98630P9.		7
15	Quantitation of five organophosphorus nerve agent metabolites in serum using hydrophilic interaction liquid chromatography and tandem mass spectrometry. Analytical and Bioanalytical Chemistry, 2014, 406, 5195-5202.	1.9	43
16	Comparison of two automated solid phase extractions for the detection of ten fentanyl analogs and metabolites in human urine using liquid chromatography tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2014, 962, 52-58.	1.2	22
17	Comparison of highâ€resolution and tandem mass spectrometry for the analysis of nerve agent metabolites in urine. Rapid Communications in Mass Spectrometry, 2013, 27, 1697-1704.	0.7	22
18	Quantification of monofluoroacetate and monochloroacetate in human urine by isotope dilution liquid chromatography tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2010, 878, 1045-1050.	1.2	8

#	Article	IF	CITATIONS
19	The determination of organophosphonate nerve agent metabolites in human urine by hydrophilic interaction liquid chromatography tandem mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2007, 852, 235-243.	1.2	86
20	Enhancing the response of alkyl methylphosphonic acids in negative electrospray ionization liquid chromatography tandem mass spectrometry by post-column addition of organic solvents. Journal of the American Society for Mass Spectrometry, 2007, 18, 1821-1826.	1.2	31