Jonathan Aldridge

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9374553/publications.pdf

Version: 2024-02-01

1478505 1588992 10 118 6 8 citations h-index g-index papers 11 11 11 189 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	T helper cells in synovial fluid of patients with rheumatoid arthritis primarily have a Th1 and a CXCR3+Th2 phenotype. Arthritis Research and Therapy, 2020, 22, 245.	3.5	29
2	Extracellular vesicles from human mesenchymal stem cells expedite chondrogenesis in 3D human degenerative disc cell cultures. Stem Cell Research and Therapy, 2020, 11, 323.	5.5	29
3	Sex-based differences in association between circulating T cell subsets and disease activity in untreated early rheumatoid arthritis patients. Arthritis Research and Therapy, 2018, 20, 150.	3.5	14
4	Blood PD-1+TFh and CTLA-4+CD4+ T cells predict remission after CTLA-4lg treatment in early rheumatoid arthritis. Rheumatology, 2022, 61, 1233-1242.	1.9	14
5	Investigation of the Effect of Secreted Factors from Mesenchymal Stem Cells on Disc Cells from Degenerated Discs. Cells Tissues Organs, 2019, 208, 76-88.	2.3	10
6	Recombinant Adiponectin Induces the Production of Pro-Inflammatory Chemokines and Cytokines in Circulating Mononuclear Cells and Fibroblast-Like Synoviocytes From Non-Inflamed Subjects. Frontiers in Immunology, 2020, 11, 569883.	4.8	10
7	Plasma interferon-alpha is associated with double-positivity for autoantibodies but is not a predictor of remission in early rheumatoid arthritis—a spin-off study of the NORD-STAR randomized clinical trial. Arthritis Research and Therapy, 2021, 23, 189.	3.5	5
8	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. Clinical and Experimental Rheumatology, 0, , .	0.8	4
9	AB0083â€ADIPONECTIN STIMULATES PRO-INFLAMMATORY CYTOKINE PRODUCTION BY PERIPHERAL BLOOD MONONUCLEAR CELLS FROM PATIENTS WITH EARLY UNTREATED RHEUMATOID ARTHRITIS. , 2019, , .		0
10	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. Clinical and Experimental Rheumatology, 2021, , .	0.8	0