## Jonathan Aldridge

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9374553/publications.pdf

Version: 2024-02-01

1478505 1588992 10 118 6 8 citations h-index g-index papers 11 11 11 189 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Blood PD-1+TFh and CTLA-4+CD4+ T cells predict remission after CTLA-4lg treatment in early rheumatoid arthritis. Rheumatology, 2022, 61, 1233-1242.	1.9	14
2	Plasma interferon-alpha is associated with double-positivity for autoantibodies but is not a predictor of remission in early rheumatoid arthritis—a spin-off study of the NORD-STAR randomized clinical trial. Arthritis Research and Therapy, 2021, 23, 189.	3.5	5
3	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. Clinical and Experimental Rheumatology, 2021, , .	0.8	0
4	T helper cells in synovial fluid of patients with rheumatoid arthritis primarily have a Th1 and a CXCR3+Th2 phenotype. Arthritis Research and Therapy, 2020, 22, 245.	3.5	29
5	Extracellular vesicles from human mesenchymal stem cells expedite chondrogenesis in 3D human degenerative disc cell cultures. Stem Cell Research and Therapy, 2020, 11, 323.	5.5	29
6	Recombinant Adiponectin Induces the Production of Pro-Inflammatory Chemokines and Cytokines in Circulating Mononuclear Cells and Fibroblast-Like Synoviocytes From Non-Inflamed Subjects. Frontiers in Immunology, 2020, 11, 569883.	4.8	10
7	AB0083â€ADIPONECTIN STIMULATES PRO-INFLAMMATORY CYTOKINE PRODUCTION BY PERIPHERAL BLOOD MONONUCLEAR CELLS FROM PATIENTS WITH EARLY UNTREATED RHEUMATOID ARTHRITIS. , 2019, , .		0
8	Investigation of the Effect of Secreted Factors from Mesenchymal Stem Cells on Disc Cells from Degenerated Discs. Cells Tissues Organs, 2019, 208, 76-88.	2.3	10
9	Sex-based differences in association between circulating T cell subsets and disease activity in untreated early rheumatoid arthritis patients. Arthritis Research and Therapy, 2018, 20, 150.	3.5	14
10	Blood chemokine levels are markers of disease activity but not predictors of remission in early rheumatoid arthritis. Clinical and Experimental Rheumatology, 0, , .	0.8	4