

# Sam D Hutchings

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9376294/publications.pdf>

Version: 2024-02-01

7  
papers

102  
citations

1684188

5  
h-index

1872680

6  
g-index

7  
all docs

7  
docs citations

7  
times ranked

165  
citing authors

#	ARTICLE	IF	CITATIONS
1	Provision of acute renal replacement therapy, using three separate modalities, in critically ill patients during the COVID-19 pandemic. An after action review from a UK tertiary critical care centre. <i>Journal of Critical Care</i> , 2021, 62, 190-196.	2.2	9
2	Critically Ill COVID-19 Patients With Acute Kidney Injury Have Reduced Renal Blood Flow and Perfusion Despite Preserved Cardiac Function: A Case-Control Study Using Contrast-Enhanced Ultrasound. <i>Shock</i> , 2021, 55, 479-487.	2.1	24
3	Microcirculatory, Endothelial, and Inflammatory Responses in Critically Ill Patients With COVID-19 Are Distinct From Those Seen in Septic Shock: A Case Control Study. <i>Shock</i> , 2021, 55, 752-758.	2.1	25
4	Quantification of stroke volume in a simulated healthy volunteer model of traumatic haemorrhage; a comparison of two non-invasive monitoring devices using error grid analysis alongside traditional measures of agreement. <i>PLoS ONE</i> , 2021, 16, e0261546.	2.5	2
5	Prospective longitudinal observational study of the macro and micro haemodynamic responses to septic shock in the renal and systemic circulations: a protocol for the MICROSHOCK "RENAL" study. <i>BMJ Open</i> , 2019, 9, e028364.	1.9	5
6	Observational study of the effects of traumatic injury, haemorrhagic shock and resuscitation on the microcirculation: a protocol for the MICROSHOCK study. <i>BMJ Open</i> , 2016, 6, e010893.	1.9	20
7	Microcirculatory perfusion shows wide inter-individual variation and is important in determining shock reversal during resuscitation in a porcine experimental model of complex traumatic hemorrhagic shock. <i>Intensive Care Medicine Experimental</i> , 2016, 4, 17.	1.9	17