
Zohreh Lasjerdi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/938098/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Inhibition of anti-inflammatory cytokines, IL-10 and TGF-β, in Leishmania major infected macrophage by miRNAs: A new therapeutic modality against leishmaniasis. Microbial Pathogenesis, 2021, 153, 104777.	1.3	9
2	Molecular identification of Acanthamoeba genotypes isolated from oral cavity of heart transplant patients in Iran. Transplant Infectious Disease, 2021, , 13744.	0.7	0
3	Isolation and Phylogenetic Analysis of Free-Living Amoebae (Acanthamoeba, Naegleria, and) Tj ETQq1 1 0.7843	14 rgBT /C	overlock 10 Tf
4	Recent researches in effective antileishmanial herbal compounds: narrative review. Parasitology Research, 2020, 119, 3929-3946.	0.6	11
5	Isolation of N. philippinensis and N. americana strains from irrigation waters of farmland soils in Iran. Environmental Science and Pollution Research, 2020, 27, 24568-24573.	2.7	6
6	Comparative Expression Profile Analysis of Apoptosis-Related miRNA and Its Target Gene in Leishmania major Infected Macrophages. Iranian Journal of Parasitology, 2020, 15, 332-340.	0.6	7
7	MicroRNAs Expression Induces Apoptosis of Macrophages in Response to Leishmania major (MRHO/IR/75/ER): An In-Vitro and In-Vivo Study. Iranian Journal of Parasitology, 2020, 15, 475-487.	0.6	9
8	Isolation, identification, and phylogenetic analysis of potentially pathogenic free-living amoebae isolated from nasal and oral mucosa of HIV/AIDS patients in Iran. Parasitology Research, 2019, 118, 3061-3066.	0.6	7
9	Comparison of the antibacterial effects of a short cationic peptide and 1% silver bioactive glass against extensively drug-resistant bacteria, Pseudomonas aeruginosa and Acinetobacter baumannii, isolated from burn patients. Amino Acids, 2018, 50, 1617-1628.	1.2	21
10	Occurrence of Free-living Amoebae in Nasal Swaps of Patients of Intensive Care Unit (ICU) and Critical Care Unit (CCU) and Their Surrounding Environments. Iranian Journal of Public Health, 2018, 47, 908-913.	0.3	1
11	An Experimental Model of Primary Amoebic Meningoence phalitis Due to in Iran. Iranian Journal of Parasitology, 2018, 13, 369-372.	0.6	1
12	Occurrence of Naegleria species in therapeutic geothermal water sources, Northern Iran. Acta Parasitologica, 2017, 62, 104-109.	0.4	26
13	Molecular identification of waterborne free living amoebae (Acanthamoeba, Naegleria and) Tj ETQq1 1 0.7843 north half of Iran. Experimental Parasitology, 2017, 183, 240-244.	14 rgBT /Ov 0.5	verlock 10 Tf 41
14	Molecular characterization of Acanthamoeba strains isolated from the oral cavity of hemodialysis patients in Iran. Parasitology Research, 2017, 116, 2965-2969.	0.6	12
15	SEROPREVALENCE AND RISK FACTORS ASSOCIATED WITH Toxoplasma gondii INFECTION AMONG RURAL COMMUNITIES IN NORTHERN IRAN. Revista Do Instituto De Medicina Tropical De Sao Paulo, 2016, 58, 70.	0.5	39
16	Presence of <i>Balamuthia mandrillaris</i> in hot springs from Mazandaran province, northern Iran. Epidemiology and Infection, 2016, 144, 2456-2461.	1.0	24
17	Isolation and molecular characterization of Acanthamoeba genotypes isolated from soil sources of public and recreational areas in Iran. Acta Parasitologica, 2016, 61, 784-789.	0.4	15
18	Distribution of Acanthamoeba Genotypes Isolated from Recreational and Therapeutic Geothermal Water Sources in Southwestern Iran, Environmental Health Insights, 2016, 10, EHI S38349	0.6	23

Zohreh Lasjerdi

#	Article	IF	CITATIONS
19	Isolation of Balamuthia mandrillaris from soil samples in North-Western Iran. Parasitology Research, 2016, 115, 541-545.	0.6	20
20	Molecular Characterization of Pathogenic <i>Acanthamoeba</i> Isolated from Drinking and Recreational water in East Azerbaijan, Northwest Iran. Environmental Health Insights, 2015, 9, EHI.S27811.	0.6	21
21	Occurrence of pathogenic Acanthamoeba genotypes in nasal swabs of cancer patients in Iran. Parasitology Research, 2015, 114, 1907-1912.	0.6	31
22	Optimization of fluoride-containing bioactive glasses as a novel scolicidal agent adjunct to hydatid surgery. Acta Tropica, 2015, 148, 105-114.	0.9	26
23	Ophthalmology hospital wards contamination to pathogenic free living Amoebae in Iran. Acta Parasitologica, 2015, 60, 417-22.	0.4	23
24	High Occurrence of Potentially Pathogenic Free Living Amoebae in Water Bodies of Kaleybar and Khodaafarin, East Azerbaijan Province. Current World Environment Journal, 2015, 10, 727-731.	0.2	0
25	Morphological and Molecular Survey of Naegleria spp. in Water Bodies Used for Recreational Purposes in Rasht city, Northern Iran. Iranian Journal of Parasitology, 2015, 10, 523-9.	0.6	9
26	Comparison of the RE and B1 gene for detection of Toxoplasma gondii infection in children with cancer. Parasitology International, 2014, 63, 37-41.	0.6	59
27	Application of Multiplex PCR for Detection and Differentiation of Entamoeba histolytica, Entamoeba dispar and Entamoeba moshkovskii. Iranian Journal of Parasitology, 2014, 9, 466-73.	0.6	8
28	Screening of recreational areas of rivers for potentially pathogenic free-living amoebae in the suburbs of Tehran, Iran. Journal of Water and Health, 2012, 10, 140-146.	1.1	37
29	Potentially pathogenic free-living amoebae isolated from hospital wards with immunodeficient patients in Tehran, Iran. Parasitology Research, 2011, 109, 575-580.	0.6	68