

# Marjan Hosseini

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9388875/publications.pdf>

Version: 2024-02-01

10  
papers

170  
citations

1478505

6  
h-index

1474206

9  
g-index

10  
all docs

10  
docs citations

10  
times ranked

296  
citing authors

#	ARTICLE	IF	CITATIONS
1	Rutin, a quercetin glycoside, alleviates acute endotoxemic kidney injury in C57BL/6 mice via suppression of inflammation and up-regulation of antioxidants and SIRT1. <i>European Journal of Pharmacology</i> , 2018, 833, 307-313.	3.5	67
2	Spinal 5-HT3 receptor mediates nociceptive effect on central neuropathic pain; possible therapeutic role for tropisetron. <i>Journal of Spinal Cord Medicine</i> , 2016, 39, 212-219.	1.4	27
3	Huperzine A ameliorates cognitive dysfunction and neuroinflammation in kainic acid-induced epileptic rats by antioxidant activity and <i>NLRP3/caspase-1</i> pathway inhibition. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2019, 46, 360-372.	1.9	20
4	The Effect of Intrathecal Administration of Muscimol on Modulation of Neuropathic Pain Symptoms Resulting from Spinal Cord Injury; an Experimental Study. <i>Emergency</i> , 2014, 2, 151-7.	0.6	18
5	Simultaneous intrathecal injection of muscimol and endomorphin-1 alleviates neuropathic pain in rat model of spinal cord injury. <i>Brain and Behavior</i> , 2020, 10, e01576.	2.2	16
6	Netrin-1 protects the SH-SY5Y cells against amyloid beta neurotoxicity through NF- $\kappa$ B/Nrf2 dependent mechanism. <i>Molecular Biology Reports</i> , 2020, 47, 9271-9277.	2.3	8
7	The inhibiting role of periaqueductal gray metabotropic glutamate receptor subtype 8 in a rat model of central neuropathic pain. <i>Neurological Research</i> , 2020, 42, 515-521.	1.3	6
8	Targeting Caveolin-1 and Claudin-5 with AY9944, Improve Blood-Brain Barrier Permeability; Computational Simulation and Experimental Study. <i>Cellular and Molecular Neurobiology</i> , 2022, 42, 1125-1139.	3.3	5
9	Evaluation of the Effect of (S)-3,4-Dicarboxyphenylglycine as a Metabotropic Glutamate Receptors Subtype 8 Agonist on Thermal Nociception Following Central Neuropathic Pain. <i>Asian Spine Journal</i> , 2021, 15, 200-206.	2.0	3
10	The effect of periaqueductal gray's metabotropic glutamate receptor subtype 8 activation on locomotor function following spinal cord injury. <i>Scandinavian Journal of Pain</i> , 2020, 20, 785-793.	1.3	0