

Jari Jokelainen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9397469/jari-jokelainen-publications-by-year.pdf>

Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

209 papers	5,654 citations	42 h-index	65 g-index
215 ext. papers	6,576 ext. citations	4.3 avg, IF	5.41 L-index

#	Paper	IF	Citations
209	Association of personalised care plans with monitoring and control of clinical outcomes, prescription of medication and utilisation of primary care services in patients with type 2 diabetes: an observational real-world study.. <i>Scandinavian Journal of Primary Health Care</i> , 2022 , 1-9	2.7	0
208	Ocular Symptoms and Rosacea: A Population-Based Study.. <i>Dermatology</i> , 2022 , 1-5	4.4	
207	Skin cancers and their risk factors in older persons: a population-based study.. <i>BMC Geriatrics</i> , 2022 , 22, 269	4.1	1
206	Characterizing Vocal Tract Dimensions in the Vocal Modes Using Magnetic Resonance Imaging. <i>Journal of Voice</i> , 2021 , 35, 804.e27-804.e42	1.9	2
205	Dose-escalation of adalimumab, golimumab or ustekinumab in inflammatory bowel diseases: characterization and implications in real-life clinical practice.. <i>Scandinavian Journal of Gastroenterology</i> , 2021 , 1-9	2.4	0
204	The association between chronic venous disease and measures of physical performance in older people: a population-based study. <i>BMC Geriatrics</i> , 2021 , 21, 556	4.1	1
203	GLP-1 Analogs and SGLT2 Inhibitors Do Not Increase Risk of Bullous Pemphigoid. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 2969-2972.e1	4.3	0
202	Long-term outcomes of lifestyle intervention to prevent type 2 diabetes in people at high risk in primary health care. <i>Primary Care Diabetes</i> , 2021 , 15, 444-450	2.4	4
201	Inflammation, hippocampal volume, and cognition in schizophrenia: results from the Northern Finland Birth Cohort 1966. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2021 , 271, 609-622	5.1	9
200	Atopic Dermatitis Is Associated with Dermatitis Herpetiformis and Celiac Disease in Children. <i>Journal of Investigative Dermatology</i> , 2021 , 141, 191-193.e2	4.3	4
199	Association of hyperglycaemia with periodontal status: Results of the Northern Finland Birth Cohort 1966 study. <i>Journal of Clinical Periodontology</i> , 2021 , 48, 24-36	7.7	2
198	Characterisation of healthcare utilisation and cost of haemophilia care in real-life: A 4-year follow-up study in Finland. <i>Haemophilia</i> , 2021 , 27, e30-e39	3.3	
197	Incidence and remission of aeroallergen sensitization in adults in Northern Finland: 15 years longitudinal study. <i>Scientific Reports</i> , 2021 , 11, 4249	4.9	3
196	Measures of empathy and the capacity for self-reflection in dental and medical students. <i>BMC Medical Education</i> , 2021 , 21, 114	3.3	1
195	A Childhood Farm Environment Protects from Allergic Sensitization until Middle Age but Not from New-Onset Sensitization in Adulthood: A 15 Year Longitudinal Study. <i>International Journal of Environmental Research and Public Health</i> , 2021 , 18,	4.6	2
194	Association between accelerometer-measured physical activity, glucose metabolism, and waist circumference in older adults. <i>Diabetes Research and Clinical Practice</i> , 2021 , 178, 108937	7.4	0
193	Association of long-term obesity and weight gain with periodontal pocketing: Results of the Northern Finland Birth Cohort 1966 study. <i>Journal of Clinical Periodontology</i> , 2021 , 48, 1344-1355	7.7	0

192	Male patients with rosacea have increased risk for migraine: a population-based study. <i>British Journal of Dermatology</i> , 2021 , 185, 1058-1061	4	1
191	Long-term dysglycemia as a risk factor for faster cognitive decline during aging: A 12-year follow-up study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 180, 109045	7.4	2
190	Factors predicting 31-year survival among a population cohort in Northern Finland. <i>International Journal of Circumpolar Health</i> , 2021 , 80, 1909334	1.7	
189	Association of mitochondrial DNA haplogroups J and K with low response in exercise training among Finnish military conscripts. <i>BMC Genomics</i> , 2021 , 22, 75	4.5	0
188	Association between psychosocial distress, sexual disorders, self-esteem and quality of life with male androgenetic alopecia: a population-based study with men at age 46. <i>BMJ Open</i> , 2021 , 11, e049853		
187	Implementation of a Personalized Care Plan for Patients With Type 2 Diabetes Is Associated With Improvements in Clinical Outcomes: An Observational Real-World Study. <i>Journal of Primary Care and Community Health</i> , 2020 , 11, 2150132720921700	2.1	4
186	The prevalence of chronic kidney disease and screening of renal function in type 2 diabetic patients in Finnish primary healthcare. <i>Primary Care Diabetes</i> , 2020 , 14, 639-644	2.4	2
185	Population-based Data at Ages 31 and 46 Show Decreased HRQoL and Life Satisfaction in Women with PCOS Symptoms. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	11
184	Could vitamin D reduce obesity-associated inflammation? Observational and Mendelian randomization study. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1036-1047	7	12
183	Overweight, obesity and hyperandrogenemia are associated with gestational diabetes mellitus: A follow-up cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020 , 99, 1311-1319	3.8	8
182	Effect of Hidradenitis Suppurativa Disease Duration on Psychiatric Comorbidity. <i>Acta Dermato-Venereologica</i> , 2020 , 100, adv00216	2.2	1
181	Comorbidities of Alopecia Areata in Finland between 1987 and 2016. <i>Acta Dermato-Venereologica</i> , 2020 , 100, adv00063	2.2	0
180	Current use of combined hormonal contraception is associated with glucose metabolism disorders in perimenopausal women. <i>European Journal of Endocrinology</i> , 2020 , 183, 619-626	6.5	1
179	Type 1 and type 2 diabetes after gestational diabetes: a 23-year cohort study. <i>Diabetologia</i> , 2020 , 63, 2123-2128	10.3	11
178	The High Prevalence of Skin Diseases in Adults Aged 70 and Older. <i>Journal of the American Geriatrics Society</i> , 2020 , 68, 2565-2571	5.6	9
177	Cold weather-related cardiorespiratory symptoms predict higher morbidity and mortality. <i>Environmental Research</i> , 2020 , 191, 110108	7.9	8
176	Nurses' Self-assessments of adherence to guidelines on safe medication preparation and administration in long-term elderly care. <i>Scandinavian Journal of Caring Sciences</i> , 2020 , 34, 108-117	2.3	10
175	Circulating metabolites and the risk of type 2 diabetes: a prospective study of 11,896 young adults from four Finnish cohorts. <i>Diabetologia</i> , 2019 , 62, 2298-2309	10.3	54

174	Drugs used for neurologic and psychiatric conditions increase the risk for bullous pemphigoid: A case-control study. <i>Journal of the American Academy of Dermatology</i> , 2019 , 81, 250-253	4.5	8
173	Genome-wide association study identifies seven novel loci associating with circulating cytokines and cell adhesion molecules in Finns. <i>Journal of Medical Genetics</i> , 2019 , 56, 607-616	5.8	21
172	Climacteric Status at the Age of 46: Impact on Metabolic Outcomes in Population-Based Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 2701-2711	5.6	5
171	Long-term metabolic syndrome is associated with periodontal pockets and alveolar bone loss. <i>Journal of Clinical Periodontology</i> , 2019 , 46, 799-808	7.7	9
170	Effect of Physical Activity on Plasma PCSK9 in Subjects With High Risk for Type 2 Diabetes. <i>Frontiers in Physiology</i> , 2019 , 10, 456	4.6	2
169	Vitamin D Status and Components of Metabolic Syndrome in Older Subjects from Northern Finland (Latitude 65°North). <i>Nutrients</i> , 2019 , 11,	6.7	18
168	REPLY to MS#JAAD-D-19-0054 "Bullous pemphigoid and neuropsychiatric medications: An influence of drugs or of underlying conditions?". <i>Journal of the American Academy of Dermatology</i> , 2019 ,	4.5	
167	Glucose status and depressive symptoms: a cohort study of elderly people in northwest Finland. <i>Scandinavian Journal of Primary Health Care</i> , 2019 , 37, 242-248	2.7	2
166	Association of Physical Activity With Telomere Length Among Elderly Adults - The Oulu Cohort 1945. <i>Frontiers in Physiology</i> , 2019 , 10, 444	4.6	6
165	Adult Patients with Atopic Eczema have a High Burden of Psychiatric Disease: A Finnish Nationwide Registry Study. <i>Acta Dermato-Venereologica</i> , 2019 , 99, 647-651	2.2	7
164	Increased Risk of Cardiovascular Diseases in Female Rosacea Patients: A Nested Case-control Study. <i>Acta Dermato-Venereologica</i> , 2019 , 99, 705-706	2.2	9
163	Dermatitis Herpetiformis and Celiac Disease Increase the Risk of Bullous Pemphigoid. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 600-604	4.3	14
162	Validation of the Zung self-rating depression scale (SDS) in older adults. <i>Scandinavian Journal of Primary Health Care</i> , 2019 , 37, 353-357	2.7	44
161	High Home Blood Pressure Variability Associates With Exaggerated Blood Pressure Response to Cold Stress. <i>American Journal of Hypertension</i> , 2019 , 32, 538-546	2.3	3
160	Pemphigus Foliaceus and Pemphigus Erythematosus are the Most Common Subtypes of Pemphigus in Northern Finland. <i>Acta Dermato-Venereologica</i> , 2019 , 99, 1127-1130	2.2	1
159	Characterization of inflammatory bowel disease management by vedolizumab and concomitant treatments in real-life clinical practice. <i>Biologicals</i> , 2019 , 58, 50-56	1.8	2
158	Insulin resistance and systemic metabolic changes in oral glucose tolerance test in 5340 individuals: an interventional study. <i>BMC Medicine</i> , 2019 , 17, 217	11.4	19
157	Substantially reduced life expectancy in patients with hidradenitis suppurativa: a Finnish nationwide registry study. <i>British Journal of Dermatology</i> , 2019 , 180, 1543-1544	4	11

156	Pressure pain sensitivity is associated with dental fear in adults in middle age: Findings from the Northern Finland 1966 birth cohort study. <i>Community Dentistry and Oral Epidemiology</i> , 2019 , 47, 193-200 ^{2.8}	6
155	Vildagliptin Significantly Increases the Risk of Bullous Pemphigoid: A Finnish Nationwide Registry Study. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 1659-1661	4.3 77
154	High treatment persistence rate and significant endoscopic healing among real-life patients treated with vedolizumab - a Finnish Nationwide Inflammatory Bowel Disease Cohort Study (FINVEDO). <i>Scandinavian Journal of Gastroenterology</i> , 2018 , 53, 158-167	2.4 23
153	Somatic and psychiatric comorbidities of hidradenitis suppurativa in children and adolescents. <i>Journal of the American Academy of Dermatology</i> , 2018 , 79, 514-519	4.5 33
152	The Association Between Low Grade Systemic Inflammation and Skin Diseases: A Cross-sectional Survey in the Northern Finland Birth Cohort 1966. <i>Acta Dermato-Venereologica</i> , 2018 , 98, 65-69	2.2 17
151	Adipokines and inflammatory markers in elderly subjects with high risk of type 2 diabetes and cardiovascular disease. <i>Scientific Reports</i> , 2018 , 8, 12816	4.9 26
150	Patients with Hidradenitis Suppurativa Have a High Psychiatric Disease Burden: A Finnish Nationwide Registry Study. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 46-51	4.3 36
149	Relationship between BMI and emotion-handling capacity in an adult Finnish population: The Northern Finland Birth Cohort 1966. <i>PLoS ONE</i> , 2018 , 13, e0203660	3.7 3
148	Women with Hidradenitis Suppurativa Have an Elevated Risk of Suicide. <i>Journal of Investigative Dermatology</i> , 2018 , 138, 2672-2674	4.3 16
147	Oral diabetes medications other than dipeptidyl peptidase 4 inhibitors are not associated with bullous pemphigoid: A Finnish nationwide case-control study. <i>Journal of the American Academy of Dermatology</i> , 2018 , 79, 1034-1038.e5	4.5 32
146	Early determinants of metabolically healthy obesity in young adults: study of the Northern Finland Birth Cohort 1966. <i>International Journal of Obesity</i> , 2018 , 42, 1704-1714	5.5 5
145	Effect of physical activity on pulse wave velocity in elderly subjects with normal glucose, prediabetes or Type 2 Diabetes. <i>Scientific Reports</i> , 2018 , 8, 8045	4.9 11
144	Primary health care utilisation and its costs among middle-aged smokers. <i>European Journal of Health Economics</i> , 2017 , 18, 351-360	3.6 3
143	The Modified Frontal Behavioral Inventory (FBI-mod) for Patients with Frontotemporal Lobar Degeneration, Alzheimer's Disease, and Mild Cognitive Impairment. <i>Journal of Alzheimer's Disease</i> , 2017 , 56, 1241-1251	4.3 9
142	Depressive symptom profiles and glucose tolerance status. <i>World Psychiatry</i> , 2017 , 16, 112	14.4 3
141	Potential determinants of vitamin D in Finnish adults: a cross-sectional study from the Northern Finland birth cohort 1966. <i>BMJ Open</i> , 2017 , 7, e013161	3 22
140	Psychological Distress Is More Prevalent in Fertile Age and Premenopausal Women With PCOS Symptoms: 15-Year Follow-Up. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 1861-1869	5.6 49
139	Diabetes and impaired glucose metabolism is associated with more cold-related cardiorespiratory symptoms. <i>Diabetes Research and Clinical Practice</i> , 2017 , 129, 116-125	7.4 5

138	Neuropsychological Profile in the C9ORF72 Associated Behavioral Variant Frontotemporal Dementia. <i>Journal of Alzheimer's Disease</i> , 2017 , 58, 479-489	4.3	8
137	Alcohol Consumption and Binge Drinking in Young Men as Predictors of Body Composition Changes During Military Service. <i>Alcohol and Alcoholism</i> , 2017 , 52, 365-371	3.5	
136	Long-term adolescent multi-site musculoskeletal pain is associated with psychological distress and anxiety. <i>Journal of Psychosomatic Research</i> , 2017 , 93, 28-32	4.1	15
135	Overweight and obese but not normal weight women with PCOS are at increased risk of Type 2 diabetes mellitus-a prospective, population-based cohort study. <i>Human Reproduction</i> , 2017 , 32, 423-431	5.7	75
134	Association of Multiple Melanocytic Naevi with Education, Sex and Skin Type. A Northern Finland Birth Cohort 1966 Study with 46 Years Follow-up. <i>Acta Dermato-Venereologica</i> , 2017 , 97, 219-224	2.2	3
133	The association of generalized anxiety disorder and Somatic Symptoms with frequent attendance to health care services: A cross-sectional study from the Northern Finland Birth Cohort 1966. <i>International Journal of Psychiatry in Medicine</i> , 2017 , 52, 147-159	1	8
132	Low level activity thresholds for changes in NMR biomarkers and genes in high risk subjects for Type 2 Diabetes. <i>Scientific Reports</i> , 2017 , 7, 11267	4.9	2
131	Abnormal skin in toe webs is a marker for abnormal glucose metabolism. A cross-sectional survey among 1,849 adults in Finland. <i>Scientific Reports</i> , 2017 , 7, 9125	4.9	2
130	Finnish Dentists Find Smoking Cessation Important but Seldom Offer Practical Support for Their Patients. <i>Community Dental Health</i> , 2017 , 34, 241-247	0.9	
129	Uterine fibroids and cardiovascular risk. <i>Human Reproduction</i> , 2016 , 31, 2689-2703	5.7	18
128	Psychiatric and neurological disorders are associated with bullous pemphigoid - a nationwide Finnish Care Register study. <i>Scientific Reports</i> , 2016 , 6, 37125	4.9	40
127	Work disabilities and unmet needs for health care and rehabilitation among jobseekers: a community-level investigation using multidimensional work ability assessments. <i>Scandinavian Journal of Primary Health Care</i> , 2016 , 34, 343-351	2.7	7
126	Metabolic profiling of pregnancy: cross-sectional and longitudinal evidence. <i>BMC Medicine</i> , 2016 , 14, 205	11.4	85
125	Association of maternal smoking during pregnancy with aerobic fitness of offspring in young adulthood: a prospective cohort study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1789-95	3.7	7
124	The association between impaired glucose tolerance and soluble CD40 ligand: a 15-year prospective cohort study. <i>Aging Clinical and Experimental Research</i> , 2016 , 28, 1243-1249	4.8	2
123	Risk of Death in Bullous Pemphigoid: A Retrospective Database Study in Finland. <i>Acta Dermato-Venereologica</i> , 2016 , 96, 758-61	2.2	9
122	A Decrease in Temperature and Humidity Precedes Human Rhinovirus Infections in a Cold Climate. <i>Viruses</i> , 2016 , 8,	6.2	33
121	Cardiovascular disease risk factors in relation to smoking behaviour and history: a population-based cohort study. <i>Open Heart</i> , 2016 , 3, e000358	3	31

120	The Association Between Hospital-Treated Injuries and ADHD Symptoms in Childhood and Adolescence: A Follow-Up Study in the Northern Finland Birth Cohort 1986. <i>Journal of Attention Disorders</i> , 2016 , 20, 3-10	3.7	20
119	Generalised anxiety disorder symptoms and utilisation of health care services. A cross-sectional study from the "Northern Finland 1966 Birth Cohort". <i>Scandinavian Journal of Primary Health Care</i> , 2016 , 34, 151-8	2.7	11
118	Changes in lifestyle modestly reduce the estimated cardiovascular disease risk in one-year follow-up of the Finnish diabetes prevention program (FIN-D2D). <i>European Journal of Cardiovascular Nursing</i> , 2015 , 14, 145-52	3.3	13
117	Depression and Insulin Resistance: Additional Support for the Novel Heuristic Model in Perimenopausal Depression. <i>American Journal of Psychiatry</i> , 2015 , 172, 796-7	11.9	2
116	Transcranial bright light and symptoms of jet lag: a randomized, placebo-controlled trial. <i>Aerospace Medicine and Human Performance</i> , 2015 , 86, 344-50	1.1	7
115	Physicians discuss the risks of smoking with their patients, but seldom offer practical cessation support. <i>Substance Abuse Treatment, Prevention, and Policy</i> , 2015 , 10, 43	3.4	22
114	Changes in glucose metabolism in people with different glucose metabolism disorders at baseline: follow-up results of a Finnish national diabetes prevention programme. <i>Diabetic Medicine</i> , 2015 , 32, 1611-8	3.5	2
113	Do depressive symptoms have an impact on the effectiveness of lifestyle counseling in prevention of type 2 diabetes? One-year follow-up of FIN-D2D. <i>Primary Care Diabetes</i> , 2014 , 8, 43-7	2.4	2
112	Comparative cost analysis of generalized anxiety disorder and major depressive disorder patients in secondary care from a national hospital registry in Finland. <i>Nordic Journal of Psychiatry</i> , 2014 , 68, 306-10	2.3	1
111	Decline in temperature and humidity increases the occurrence of influenza in cold climate. <i>Environmental Health</i> , 2014 , 13, 22	6	83
110	Increasing incidence of bullous pemphigoid in Northern Finland: a retrospective database study in Oulu University Hospital. <i>British Journal of Dermatology</i> , 2014 , 171, 1223-6	4	43
109	Light physical activity determined by a motion sensor decreases insulin resistance, improves lipid homeostasis and reduces visceral fat in high-risk subjects: PreDiabEx study RCT. <i>International Journal of Obesity</i> , 2014 , 38, 1089-96	5.5	46
108	Prevalence of anxiety disorders among Finnish primary care high utilizers and validation of Finnish translation of GAD-7 and GAD-2 screening tools. <i>Scandinavian Journal of Primary Health Care</i> , 2014 , 32, 78-83	2.7	26
107	Cold-related cardiorespiratory symptoms among subjects with and without hypertension: the National FINRISK Study 2002. <i>European Journal of Public Health</i> , 2014 , 24, 237-43	2.1	17
106	Lifestyle intervention in prevention of type 2 diabetes in women with a history of gestational diabetes mellitus: one-year results of the FIN-D2D project. <i>Journal of Women's Health</i> , 2014 , 23, 506-12	3	20
105	Transcranial bright light treatment via the ear canals in seasonal affective disorder: a randomized, double-blind dose-response study. <i>BMC Psychiatry</i> , 2014 , 14, 288	4.2	10
104	The association between erectile dysfunction, depressive symptoms and testosterone levels among middle-aged men. <i>Scandinavian Journal of Public Health</i> , 2014 , 42, 677-82	3	11
103	Altered resting-state activity in seasonal affective disorder. <i>Human Brain Mapping</i> , 2014 , 35, 161-72	5.9	26

102	High prevalence of skin diseases and need for treatment in a middle-aged population. A Northern Finland Birth Cohort 1966 study. <i>PLoS ONE</i> , 2014 , 9, e99533	3.7	19
101	The association between physical fitness and depressive symptoms among young adults: results of the Northern Finland 1966 birth cohort study. <i>BMC Public Health</i> , 2013 , 13, 535	4.1	36
100	Clinical outcome after crush versus culotte stenting of coronary artery bifurcation lesions: the Nordic Stent Technique Study 36-month follow-up results. <i>JACC: Cardiovascular Interventions</i> , 2013 , 6, 1160-5	5	42
99	Predictors of success of a lifestyle intervention in relation to weight loss and improvement in glucose tolerance among individuals at high risk for type 2 diabetes: the FIN-D2D project. <i>Journal of Primary Care and Community Health</i> , 2013 , 4, 59-66	2.1	11
98	Reduction in metabolic syndrome among obese young men is associated with exercise-induced body composition changes during military service. <i>Diabetes Research and Clinical Practice</i> , 2012 , 98, 312-9	3.4	9
97	Aerobic performance and body composition changes during military service. <i>Scandinavian Journal of Primary Health Care</i> , 2012 , 30, 95-100	2.7	18
96	Can transcranial brain-targeted bright light treatment via ear canals be effective in relieving symptoms in seasonal affective disorder? A pilot study. <i>Medical Hypotheses</i> , 2012 , 78, 511-5	3.8	14
95	Participation, socioeconomic status and group or individual counselling intervention in individuals at high risk for type 2 diabetes: one-year follow-up study of the FIN-D2D-project. <i>Primary Care Diabetes</i> , 2012 , 6, 277-83	2.4	10
94	Family history of diabetes and effectiveness of lifestyle counselling on the cardio-metabolic risk profile in individuals at high risk of Type 2 diabetes: 1-year follow-up of the FIN-D2D project. <i>Diabetic Medicine</i> , 2012 , 29, 207-11	3.5	12
93	Unacylated ghrelin is associated with changes in insulin sensitivity and lipid profile during an exercise intervention. <i>Clinical Endocrinology</i> , 2012 , 76, 39-45	3.4	19
92	Prevalence of metabolic syndrome components among the elderly using three different definitions: a cohort study in Finland. <i>Scandinavian Journal of Primary Health Care</i> , 2012 , 30, 29-34	2.7	18
91	Validation of the Whooley questions and the Beck Depression Inventory in older adults. <i>Scandinavian Journal of Primary Health Care</i> , 2012 , 30, 259-64	2.7	18
90	Precariousness and discontinuous work history in association with health. <i>Scandinavian Journal of Public Health</i> , 2012 , 40, 360-7	3	32
89	Do statins interfere with lifestyle intervention in the prevention of diabetes in primary healthcare? One-year follow-up of the FIN-D2D project. <i>BMJ Open</i> , 2012 , 2,	3	8
88	Unacylated ghrelin is associated with changes in body composition and body fat distribution during long-term exercise intervention. <i>European Journal of Endocrinology</i> , 2011 , 165, 243-8	6.5	23
87	The association between anxiety and C-reactive protein (CRP) levels: results from the Northern Finland 1966 birth cohort study. <i>European Psychiatry</i> , 2011 , 26, 363-9	6	79
86	Exercise during military training improves cardiovascular risk factors in young men. <i>Atherosclerosis</i> , 2011 , 216, 489-95	3.1	15
85	Increase in physical activity and cardiometabolic risk profile change during lifestyle intervention in primary healthcare: 1-year follow-up study among individuals at high risk for type 2 diabetes. <i>BMJ Open</i> , 2011 , 1, e000292	3	29

84	Socioeconomic position and effectiveness of lifestyle intervention in prevention of type 2 diabetes: one-year follow-up of the FIN-D2D project. <i>Scandinavian Journal of Public Health</i> , 2011 , 39, 561-70	3	22
83	Limited overlap between intermediate hyperglycemia as defined by A1C 5.7-6.4%, impaired fasting glucose, and impaired glucose tolerance. <i>Diabetes Care</i> , 2011 , 34, 2314-6	14.6	36
82	Determinants of functional recovery after myocardial infarction of patients treated with bone marrow-derived stem cells after thrombolytic therapy. <i>Heart</i> , 2010 , 96, 362-7	5.1	36
81	Postchallenge glucose, A1C, and fasting glucose as predictors of type 2 diabetes and cardiovascular disease: a 10-year prospective cohort study. <i>Diabetes Care</i> , 2010 , 33, 2077-83	14.6	80
80	Depressive symptoms and 10-year risk for cardiovascular morbidity and mortality. <i>World Journal of Biological Psychiatry</i> , 2010 , 11, 834-9	3.8	38
79	Despite evidence-based guidelines, systolic blood pressure remains inadequately controlled in older hypertensive adults. <i>Journal of Human Hypertension</i> , 2010 , 24, 439-46	2.6	15
78	Prevalence of dementia a rising challenge among ageing populations. <i>European Geriatric Medicine</i> , 2010 , 1, 330-333	3	11
77	Cardiometabolic profile of people screened for high risk of type 2 diabetes in a national diabetes prevention programme (FIN-D2D). <i>Primary Care Diabetes</i> , 2010 , 4, 231-9	2.4	22
76	Deviant intrauterine growth and risk of schizophrenia: a 34-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2010 , 124, 223-30	3.6	36
75	Serum ghrelin and the prediction of the development of impaired glucose regulation and Type 2 diabetes in middle-aged subjects. <i>Journal of Endocrinological Investigation</i> , 2010 , 33, 496-500	5.2	15
74	Lifestyle intervention for prevention of type 2 diabetes in primary health care: one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D). <i>Diabetes Care</i> , 2010 , 33, 2146-51	14.6	228
73	The relationships between high-sensitivity C-reactive protein and incident depressed mood among older adults. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2010 , 70, 75-9	2	15
72	Effect of menopause and use of contraceptives/hormone therapy on association of C-reactive protein and depression: a population-based study. <i>Journal of Psychosomatic Research</i> , 2010 , 68, 573-9	4.1	4
71	Differences in COPD-related readmissions to primary and secondary care hospitals. <i>Scandinavian Journal of Primary Health Care</i> , 2009 , 27, 80-4	2.7	2
70	Falls risk among a very old home-dwelling population. <i>Scandinavian Journal of Primary Health Care</i> , 2009 , 27, 25-30	2.7	56
69	Depressive symptoms predispose females to metabolic syndrome: a 7-year follow-up study. <i>Acta Psychiatrica Scandinavica</i> , 2009 , 119, 137-42	6.5	78
68	Depressive symptomatology is associated with decreased interleukin-1 beta and increased interleukin-1 receptor antagonist levels in males. <i>Psychiatry Research</i> , 2009 , 167, 73-9	9.9	41
67	Cold temperature and low humidity are associated with increased occurrence of respiratory tract infections. <i>Respiratory Medicine</i> , 2009 , 103, 456-62	4.6	180

66	Occurrence of frostbite in the general population--work-related and individual factors. <i>Scandinavian Journal of Work, Environment and Health</i> , 2009 , 35, 384-93	4.3	37
65	Low serum adiponectin level as a predictor of impaired glucose regulation and type 2 diabetes mellitus in a middle-aged Finnish population. <i>Metabolism: Clinical and Experimental</i> , 2008 , 57, 1130-4	12.7	42
64	Treatment of hepatitis C with interferon-alpha and ribavirine in a patient with long-term clozapine treatment. <i>Psychosomatics</i> , 2008 , 49, 86-8	2.6	3
63	Does place of treatment affect prognosis for chronic obstructive pulmonary disease (COPD)?. <i>European Journal of General Practice</i> , 2008 , 14, 123-8	2.8	1
62	Recurrent traumatic brain injury is predicted by the index injury occurring under the influence of alcohol. <i>Brain Injury</i> , 2008 , 22, 780-5	2.1	27
61	Exercise and risk of injurious fall in home-dwelling elderly. <i>International Journal of Circumpolar Health</i> , 2008 , 67, 235-44	1.7	14
60	Autonomic nervous function during whole-body cold exposure before and after cold acclimation. <i>Aviation, Space, and Environmental Medicine</i> , 2008 , 79, 875-82		63
59	Levels of adiponectin, C-reactive protein and interleukin-1 receptor antagonist are associated with insulin sensitivity: a population-based study. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 378-83	3.5	28
58	Risk of dementia associated with the ApoE epsilon4 allele and falls causing head injury without explicit traumatic brain injury. <i>Acta Neurologica Scandinavica</i> , 2008 , 118, 153-8	3.8	9
57	Metabolic syndrome predisposes to depressive symptoms: a population-based 7-year follow-up study. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 178-82	4.6	124
56	Traumatic brain injuries in children and young adults: a birth cohort study from northern Finland. <i>Neuroepidemiology</i> , 2007 , 29, 136-42	5.4	28
55	Parental alcohol misuse is a powerful predictor for the risk of traumatic brain injury in childhood. <i>Brain Injury</i> , 2007 , 21, 1079-85	2.1	30
54	Somatization symptoms in young adult Finnish population--associations with sex, educational level and mental health. <i>Nordic Journal of Psychiatry</i> , 2007 , 61, 219-24	2.3	11
53	Obesity and low-grade inflammation among young Finnish men with early-onset alopecia. <i>Dermatology</i> , 2007 , 214, 125-9	4.4	50
52	Insulin resistance and depressive symptoms in young adult males: findings from Finnish military conscripts. <i>Psychosomatic Medicine</i> , 2007 , 69, 723-8	3.7	32
51	C-reactive protein levels and sleep disturbances: observations based on the Northern Finland 1966 Birth Cohort study. <i>Psychosomatic Medicine</i> , 2007 , 69, 756-61	3.7	51
50	Pragmatic exercise-oriented prevention of falls among the elderly: a population-based, randomized, controlled trial. <i>Preventive Medicine</i> , 2007 , 44, 265-71	4.3	44
49	Metabolic syndrome in connection with BMI in young Finnish male adults. <i>Diabetes Research and Clinical Practice</i> , 2007 , 76, 404-9	7.4	20

48	Co-occurrence of personality disorders with mood, anxiety, and substance use disorders in a young adult population. <i>Journal of Personality Disorders</i> , 2006 , 20, 102-12	2.6	14
47	Prevention of disability by exercise among the elderly: a population-based, randomized, controlled trial. <i>Scandinavian Journal of Primary Health Care</i> , 2006 , 24, 199-205	2.7	42
46	The association between C-reactive protein levels and depression: Results from the northern Finland 1966 birth cohort study. <i>Biological Psychiatry</i> , 2006 , 60, 825-30	7.9	121
45	Late pregnancy blood pressure in nulliparas and subsequent abnormal glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 2006 , 71, 220-4	7.4	2
44	Adolescents' drinking habits predict later occurrence of traumatic brain injury: 35-year follow-up of the northern Finland 1966 birth cohort. <i>Journal of Adolescent Health</i> , 2006 , 39, 275.e1-7	5.8	17
43	Temporal changes in the frequency of falling accidents among the elderly during the 1990s: a population-based study. <i>Public Health</i> , 2006 , 120, 418-20	4	3
42	Common polymorphisms (single-nucleotide polymorphisms SNP+45 and SNP+276) of the adiponectin gene regulate serum adiponectin concentrations and blood pressure in young Finnish men. <i>Molecular Genetics and Metabolism</i> , 2006 , 87, 147-51	3.7	48
41	Depressive symptoms and insulin resistance in young adult males: results from the Northern Finland 1966 birth cohort. <i>Molecular Psychiatry</i> , 2006 , 11, 929-33	15.1	47
40	The Pro12Ala polymorphism of the PPAR gamma 2 gene influences sex hormone-binding globulin level and its relationship to the development of the metabolic syndrome in young Finnish men. <i>Endocrine</i> , 2006 , 30, 185-90		16
39	Factors affecting outdoor exposure in winter: population-based study. <i>International Journal of Biometeorology</i> , 2006 , 51, 27-36	3.7	44
38	Association of insulin resistance linked diseases and hair loss in elderly men. Finnish population-based study. <i>Central European Journal of Public Health</i> , 2006 , 14, 78-81	1.2	47
37	Insulin resistance and depression: cross sectional study. <i>BMJ, The</i> , 2005 , 330, 17-8	5.9	100
36	Common polymorphisms in the PPARGgamma2 and IRS-1 genes and their interaction influence serum adiponectin concentration in young Finnish men. <i>Molecular Genetics and Metabolism</i> , 2005 , 84, 344-8	3.7	22
35	Poor premorbid school performance is associated with later cigarette smoking among schizophrenia patients. <i>Psychiatry Research</i> , 2005 , 137, 137-41	9.9	7
34	Lack of increase of serum adiponectin concentrations with a moderate weight loss during six months on a high-caloric diet in military service among a young male Finnish population. <i>Endocrine</i> , 2005 , 26, 65-9		17
33	Effect of the Pro12Ala polymorphism of the PPARG2 gene on serum adiponectin changes. <i>Endocrine</i> , 2005 , 27, 307-9		13
32	Fall-related brain injuries and the risk of dementia in elderly people: a population-based study. <i>European Journal of Neurology</i> , 2005 , 12, 86-92	6	43
31	Infant developmental milestones: a 31-year follow-up. <i>Developmental Medicine and Child Neurology</i> , 2005 , 47, 581-586	3.3	51

30 2005, 330, 965.3-966

2

29	Infant developmental milestones: a 31-year follow-up. <i>Developmental Medicine and Child Neurology</i> , 2005 , 47, 581	3.3	31
28	Serum lipids in schizophrenia and other functional psychoses: a general population northern Finland 1966 birth cohort survey. <i>Acta Psychiatrica Scandinavica</i> , 2004 , 110, 279-85	6.5	26
27	Seasonal variation in postnatal depression. <i>Journal of Affective Disorders</i> , 2004 , 78, 111-8	6.6	27
26	Fish consumption and depression: the Northern Finland 1966 birth cohort study. <i>Journal of Affective Disorders</i> , 2004 , 82, 447-52	6.6	175
25	Somatization disorder in young adult population. <i>General Hospital Psychiatry</i> , 2004 , 26, 9-12	5.6	15
24	Childhood central nervous system infections and risk for schizophrenia. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2004 , 254, 9-13	5.1	79
23	Comparison of hospital-treated personality disorders and personality disorders in a general population sample. <i>Nordic Journal of Psychiatry</i> , 2004 , 58, 357-62	2.3	12
22	Schizophrenia in the offspring of antenatally depressed mothers: a 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2004 , 66, 79-81	3.6	16
21	Vitamin D supplementation during the first year of life and risk of schizophrenia: a Finnish birth cohort study. <i>Schizophrenia Research</i> , 2004 , 67, 237-45	3.6	224
20	The persistence of developmental markers in childhood and adolescence and risk for schizophrenic psychoses in adult life. A 34-year follow-up of the Northern Finland 1966 birth cohort. <i>Schizophrenia Research</i> , 2004 , 71, 213-25	3.6	49
19	Age at menarche and depression at the age of 31 years: findings from the Northern Finland 1966 Birth Cohort Study. <i>Journal of Psychosomatic Research</i> , 2004 , 57, 359-62	4.1	12
18	Higher serum triglyceride levels in early-onset schizophrenia. <i>American Journal of Psychiatry</i> , 2004 , 161, 176	11.9	12
17	Hyperlipidemia in persons using antipsychotic medication: a general population-based birth cohort study. <i>Journal of Clinical Psychiatry</i> , 2004 , 65, 547-50	4.6	34
16	The Relationship between Childhood Family Background and Educational Performance, with Special Reference to Single-parent Families: A Longitudinal Study. <i>Social Psychology of Education</i> , 2003 , 6, 349-365	2	9
15	Criminality in the offspring of antenatally depressed mothers: a 33-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Journal of Affective Disorders</i> , 2003 , 74, 273-8	6.6	36
14	Atopy and depression: results from the Northern Finland 1966 Birth Cohort Study. <i>Molecular Psychiatry</i> , 2003 , 8, 738-44	15.1	90
13	Presence of atopy in first-degree relatives as a predictor of a female proband's depression: results from the Northern Finland 1966 Birth Cohort. <i>Journal of Allergy and Clinical Immunology</i> , 2003 , 111, 1249-54	11.5	30

12	Ability to speak at the age of 1 year and alexithymia 30 years later. <i>Journal of Psychosomatic Research</i> , 2003 , 54, 491-5	4.1	19
11	The Hopkins Symptom Checklist-25 in screening DSM-III-R axis-I disorders. <i>Nordic Journal of Psychiatry</i> , 2003 , 57, 119-23	2.3	111
10	Low educational performance is associated with drunk driving: a 31-year follow-up of the northern Finland 1966 birth cohort. <i>Alcohol and Alcoholism</i> , 2003 , 38, 219-23	3.5	16
9	Association between skin test diagnosed atopy and professionally diagnosed depression: a Northern Finland 1966 Birth Cohort study. <i>Biological Psychiatry</i> , 2002 , 52, 349-55	7.9	43
8	Juvenile and early adulthood smoking and adult educational achievements--a 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Scandinavian Journal of Public Health</i> , 2001 , 29, 87-95	3	10
7	Psychometric properties of the Finnish 20-item Toronto Alexithymia Scale. <i>Nordic Journal of Psychiatry</i> , 2001 , 55, 123-7	2.3	101
6	Early developmental milestones in adult schizophrenia and other psychoses. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Schizophrenia Research</i> , 2001 , 52, 1-19	3.6	133
5	Prevalence and sociodemographic correlates of alexithymia in a population sample of young adults. <i>Comprehensive Psychiatry</i> , 2001 , 42, 471-6	7.3	140
4	Educational consequences of mental disorders treated in hospital. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. <i>Psychological Medicine</i> , 2001 , 31, 339-49	6.9	40
3	Is grand multiparity associated with offspring's hospital-treated mental disorders? A 28-year follow-up of the North Finland 1966 birth cohort. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2000 , 35, 104-8	4.5	29
2	School predictors of schizophrenia. <i>Archives of General Psychiatry</i> , 2000 , 57, 813		5
1	School performance as a predictor of psychiatric hospitalization in adult life. A 28-year follow-up in the Northern Finland 1966 Birth Cohort. <i>Psychological Medicine</i> , 1998 , 28, 967-74	6.9	88