

Carmen Fenoll

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9409183/carmen-fenoll-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

64
papers

1,742
citations

24
h-index

40
g-index

69
ext. papers

2,161
ext. citations

6.9
avg, IF

4.43
L-index

| # | Paper | IF | Citations |
|----|--|------|-----------|
| 64 | Laser Microdissection of Cells and Isolation of High-Quality RNA After Cryosectioning. <i>Methods in Molecular Biology</i> , 2021 , 2170, 35-43 | 1.4 | 0 |
| 63 | Compatible interactions between plants and endoparasitic nematodes – a follow-up of ABR volume 73: Plant nematode interactions – a view on compatible interrelationships. <i>Advances in Botanical Research</i> , 2021 , 237-248 | 2.2 | 0 |
| 62 | Root-knot nematodes induce gall formation by recruiting developmental pathways of post-embryonic organogenesis and regeneration to promote transient pluripotency. <i>New Phytologist</i> , 2020 , 227, 200-215 | 9.8 | 15 |
| 61 | The Tomato Genome Encodes SPCH, MUTE, and FAMA Candidates That Can Replace the Endogenous Functions of Their Orthologs. <i>Frontiers in Plant Science</i> , 2019 , 10, 1300 | 6.2 | 11 |
| 60 | A Genetic Dissection of Natural Variation for Stomatal Abundance Traits in. <i>Frontiers in Plant Science</i> , 2019 , 10, 1392 | 6.2 | 4 |
| 59 | A role for ALF4 during gall and giant cell development in the biotic interaction between Arabidopsis and Meloidogyne spp. <i>Physiologia Plantarum</i> , 2019 , 165, 17-28 | 4.6 | 2 |
| 58 | sRNAs involved in the regulation of plant developmental processes are altered during the root-knot nematode interaction for feeding site formation. <i>European Journal of Plant Pathology</i> , 2018 , 152, 945-955 | 2.1 | 1 |
| 57 | Silenced retrotransposons are major rasiRNAs targets in Arabidopsis galls induced by Meloidogyne javanica. <i>Molecular Plant Pathology</i> , 2018 , 19, 2431-2445 | 5.7 | 13 |
| 56 | A Phenotyping Method of Giant Cells from Root-Knot Nematode Feeding Sites by Confocal Microscopy Highlights a Role for CHITINASE-LIKE 1 in Arabidopsis. <i>International Journal of Molecular Sciences</i> , 2018 , 19, | 6.3 | 14 |
| 55 | A role for the gene regulatory module microRNA172/TARGET OF EARLY ACTIVATION TAGGED 1/FLOWERING LOCUS T (miRNA172/TOE1/FT) in the feeding sites induced by Meloidogyne javanica in Arabidopsis thaliana. <i>New Phytologist</i> , 2018 , 217, 813-827 | 9.8 | 24 |
| 54 | POLAR-guided signalling complex assembly and localization drive asymmetric cell division. <i>Nature</i> , 2018 , 563, 574-578 | 50.4 | 82 |
| 53 | Overexpression of a Gene From Wild Tomato Decreases Stomatal Density and Enhances Dehydration Avoidance in Arabidopsis and Cultivated Tomato. <i>Frontiers in Plant Science</i> , 2018 , 9, 940 | 6.2 | 25 |
| 52 | A Standardized Method to Assess Infection Rates of Root-Knot and Cyst Nematodes in Arabidopsis thaliana Mutants with Alterations in Root Development Related to Auxin and Cytokinin Signaling. <i>Methods in Molecular Biology</i> , 2017 , 1569, 73-81 | 1.4 | 7 |
| 51 | A Mutation in the bHLH Domain of the SPCH Transcription Factor Uncovers a BR-Dependent Mechanism for Stomatal Development. <i>Plant Physiology</i> , 2017 , 174, 823-842 | 6.6 | 16 |
| 50 | Molecular Transducers from Roots Are Triggered in Arabidopsis Leaves by Root-Knot Nematodes for Successful Feeding Site Formation: A Conserved Post-Embryogenic Organogenesis Program?. <i>Frontiers in Plant Science</i> , 2017 , 8, 875 | 6.2 | 13 |
| 49 | Belowground Defence Strategies Against Sedentary Nematodes. <i>Signaling and Communication in Plants</i> , 2016 , 221-251 | 1 | 1 |
| 48 | Differentially expressed small RNAs in Arabidopsis galls formed by Meloidogyne javanica: a functional role for miR390 and its TAS3-derived tasiRNAs. <i>New Phytologist</i> , 2016 , 209, 1625-40 | 9.8 | 63 |

| | | | |
|----|---|-----|----|
| 47 | Too many faces for TOO MANY MOUTHS?. <i>New Phytologist</i> , 2016 , 210, 779-85 | 9.8 | 9 |
| 46 | Long-Term In Vitro System for Maintenance and Amplification of Root-Knot Nematodes in <i>Cucumis sativus</i> Roots. <i>Frontiers in Plant Science</i> , 2016 , 7, 124 | 6.2 | 13 |
| 45 | A Reliable Protocol for In situ microRNAs Detection in Feeding Sites Induced by Root-Knot Nematodes. <i>Frontiers in Plant Science</i> , 2016 , 7, 966 | 6.2 | 7 |
| 44 | The Power of Omics to Identify Plant Susceptibility Factors and to Study Resistance to Root-knot Nematodes. <i>Current Issues in Molecular Biology</i> , 2016 , 19, 53-72 | 2.9 | 6 |
| 43 | Developmental Pathways Mediated by Hormones in Nematode Feeding Sites. <i>Advances in Botanical Research</i> , 2015 , 73, 167-188 | 2.2 | 13 |
| 42 | Overview of Root-Knot Nematodes and Giant Cells. <i>Advances in Botanical Research</i> , 2015 , 73, 1-32 | 2.2 | 35 |
| 41 | Transcriptional profiles of Arabidopsis stomataless mutants reveal developmental and physiological features of life in the absence of stomata. <i>Frontiers in Plant Science</i> , 2015 , 6, 456 | 6.2 | 7 |
| 40 | Genes co-regulated with LBD16 in nematode feeding sites inferred from in silico analysis show similarities to regulatory circuits mediated by the auxin/cytokinin balance in Arabidopsis. <i>Plant Signaling and Behavior</i> , 2015 , 10, e990825 | 2.5 | 12 |
| 39 | Phenotyping nematode feeding sites: three-dimensional reconstruction and volumetric measurements of giant cells induced by root-knot nematodes in Arabidopsis. <i>New Phytologist</i> , 2015 , 206, 868-80 | 9.8 | 25 |
| 38 | NEMATIC: a simple and versatile tool for the in silico analysis of plant-nematode interactions. <i>Molecular Plant Pathology</i> , 2014 , 15, 627-36 | 5.7 | 24 |
| 37 | Transcriptomic signatures of transfer cells in early developing nematode feeding cells of Arabidopsis focused on auxin and ethylene signaling. <i>Frontiers in Plant Science</i> , 2014 , 5, 107 | 6.2 | 24 |
| 36 | A role for LATERAL ORGAN BOUNDARIES-DOMAIN 16 during the interaction Arabidopsis-Meloidogyne spp. provides a molecular link between lateral root and root-knot nematode feeding site development. <i>New Phytologist</i> , 2014 , 203, 632-645 | 9.8 | 40 |
| 35 | The TRANSPLANTA collection of Arabidopsis lines: a resource for functional analysis of transcription factors based on their conditional overexpression. <i>Plant Journal</i> , 2014 , 77, 944-53 | 6.9 | 61 |
| 34 | Distinct and conserved transcriptomic changes during nematode-induced giant cell development in tomato compared with Arabidopsis: a functional role for gene repression. <i>New Phytologist</i> , 2013 , 197, 1276-1290 | 9.8 | 76 |
| 33 | Timely expression of the Arabidopsis stoma-fate master regulator MUTE is required for specification of other epidermal cell types. <i>Plant Journal</i> , 2013 , 75, 808-22 | 6.9 | 16 |
| 32 | Dynamic analysis of epidermal cell divisions identifies specific roles for COP10 in Arabidopsis stomatal lineage development. <i>Planta</i> , 2012 , 236, 447-61 | 4.7 | 21 |
| 31 | Laser microdissection of cells and isolation of high-quality RNA after cryosectioning. <i>Methods in Molecular Biology</i> , 2012 , 883, 87-95 | 1.4 | 13 |
| 30 | Roles of constitutive photomorphogenic 10 in Arabidopsis stomata development. <i>Plant Signaling and Behavior</i> , 2012 , 7, 990-3 | 2.5 | 3 |

| | | | |
|----|--|------|-----|
| 29 | Arabidopsis as a Tool for the Study of Plant-Nematode Interactions 2011 , 139-156 | | 3 |
| 28 | Natural variation in stomatal abundance of <i>Arabidopsis thaliana</i> includes cryptic diversity for different developmental processes. <i>Annals of Botany</i> , 2011 , 107, 1247-58 | 4.1 | 32 |
| 27 | Activation of geminivirus V-sense promoters in roots is restricted to nematode feeding sites. <i>Molecular Plant Pathology</i> , 2010 , 11, 409-17 | 5.7 | 4 |
| 26 | Early transcriptomic events in microdissected <i>Arabidopsis</i> nematode-induced giant cells. <i>Plant Journal</i> , 2010 , 61, 698-712 | 6.9 | 173 |
| 25 | Isolation of RNA from laser-capture-microdissected giant cells at early differentiation stages suitable for differential transcriptome analysis. <i>Molecular Plant Pathology</i> , 2009 , 10, 523-35 | 5.7 | 33 |
| 24 | Distinct heat-shock element arrangements that mediate the heat shock, but not the late-embryogenesis induction of small heat-shock proteins, correlate with promoter activation in root-knot nematode feeding cells. <i>Plant Molecular Biology</i> , 2008 , 66, 151-64 | 4.6 | 28 |
| 23 | Evaluation of different RNA extraction methods for small quantities of plant tissue: Combined effects of reagent type and homogenization procedure on RNA quality-integrity and yield. <i>Physiologia Plantarum</i> , 2006 , 128, 1-7 | 4.6 | 28 |
| 22 | Differential activation of ABI3 and LEA genes upon plant parasitic nematode infection. <i>Molecular Plant Pathology</i> , 2005 , 6, 321-5 | 5.7 | 16 |
| 21 | Regulation of MSV and WDV virion-sense promoters by WDV nonstructural proteins: a role for their retinoblastoma protein-binding motifs. <i>Virology</i> , 2003 , 306, 313-23 | 3.6 | 16 |
| 20 | Stomatal precursors in <i>Arabidopsis</i> : prohibiting the fulfilment of a general rule. <i>New Phytologist</i> , 2003 , 158, 427-430 | 9.8 | 3 |
| 19 | Induction of the Hahsp17.7G4 promoter by root-knot nematodes: involvement of heat-shock elements in promoter activity in giant cells. <i>Molecular Plant-Microbe Interactions</i> , 2003 , 16, 1062-8 | 3.6 | 29 |
| 18 | Specification of stomatal fate in <i>Arabidopsis</i> : evidences for cellular interactions. <i>New Phytologist</i> , 2002 , 153, 399-404 | 9.8 | 9 |
| 17 | Clonal analysis of stomatal development and patterning in <i>Arabidopsis</i> leaves. <i>Developmental Biology</i> , 2002 , 241, 24-33 | 3.1 | 31 |
| 16 | Gene expression in nematode feeding sites. <i>Annual Review of Phytopathology</i> , 2002 , 40, 191-219 | 10.8 | 246 |
| 15 | Concerted Efforts To Develop Handles For Plant Parasitic Nematode Control. <i>Developments in Plant Genetics and Breeding</i> , 2000 , 6, 159-167 | | |
| 14 | Stomatal development and patterning in <i>Arabidopsis</i> leaves. <i>Physiologia Plantarum</i> , 2000 , 109, 351-358 | 4.6 | 24 |
| 13 | Cis-Elements In Nematode-Responsive Promoters. <i>Developments in Plant Genetics and Breeding</i> , 2000 , 6, 177-182 | | 1 |
| 12 | Stomatal development in <i>Arabidopsis</i> : how to make a functional pattern. <i>Trends in Plant Science</i> , 2000 , 5, 458-60 | 13.1 | 31 |

| | | | |
|----|--|-----|----|
| 11 | Isolation of the LEMMI9 gene and promoter analysis during a compatible plant-nematode interaction. <i>Molecular Plant-Microbe Interactions</i> , 1999 , 12, 440-9 | 3.6 | 30 |
| 10 | Tracing the ontogeny of stomatal clusters in Arabidopsis with molecular markers. <i>Plant Journal</i> , 1997 , 12, 747-55 | 6.9 | 67 |
| 9 | Regulation of Gene Expression in Feeding Sites. <i>Developments in Plant Pathology</i> , 1997 , 133-149 | | 8 |
| 8 | The two nonstructural proteins from wheat dwarf virus involved in viral gene expression and replication are retinoblastoma-binding proteins. <i>Virology</i> , 1996 , 219, 324-9 | 3.6 | 65 |
| 7 | The intergenic region of maize streak virus contains a GC-rich element that activates rightward transcription and binds maize nuclear factors. <i>Plant Molecular Biology</i> , 1990 , 15, 865-77 | 4.6 | 42 |
| 6 | The intergenic region of maize streak virus contains promoter elements involved in rightward transcription of the viral genome. <i>EMBO Journal</i> , 1988 , 7, 1589-1596 | 13 | 35 |
| 5 | Construction and Homologous Expression of a Maize Adh1 Based NcoI Cassette Vector. <i>Plant Physiology</i> , 1987 , 85, 327-30 | 6.6 | 16 |
| 4 | On the mechanism of respiratory and photosynthetic electron transfer in <i>Rhodospirillum rubrum</i> . <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1987 , 892, 172-184 | 4.6 | 17 |
| 3 | Purification of a light-harvesting B880 complex from wild-type <i>Rhodospirillum rubrum</i> . <i>Analytical Biochemistry</i> , 1986 , 152, 29-34 | 3.1 | 4 |
| 2 | The membrane potential of intact <i>Rhodospirillum rubrum</i> cells in the absence of light-dependent and oxygen-linked electron transfer. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1985 , 806, 168-174 | 4.6 | 6 |
| 1 | Simultaneous presence of two terminal oxidases in the respiratory system of dark aerobically grown <i>Rhodospirillum rubrum</i> . <i>Archives of Microbiology</i> , 1984 , 137, 42-46 | 3 | 11 |