

Jason Jungsik Song

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/941485/jason-jungsik-song-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

136
papers

671
citations

13
h-index

18
g-index

167
ext. papers

887
ext. citations

3
avg, IF

4.42
L-index

#	Paper	IF	Citations
136	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis.. <i>Journal of Clinical Laboratory Analysis</i> , 2022 , e24237	3	1
135	Association Between Idiopathic Cutaneous Leukocytoclastic Angiitis and ANCA-negative Microscopic Polyangiitis. <i>Journal of Rheumatic Diseases</i> , 2022 , 29, 40-45	1.2	
134	Incidence and Patterns of Interstitial Lung Disease and Their Clinical Impact on Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Korean Single-Centre Observational Study. <i>Journal of Immunology Research</i> , 2022 , 2022, 1-7	4.5	0
133	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e13929	2.9	1
132	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. <i>Clinical Rheumatology</i> , 2021 , 1	3.9	2
131	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021 , 69, 91-95	2.9	
130	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 1016-1022	3	1
129	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e24048	3	1
128	B-cell metabolism regulator IM156 contributes to the mitigation of systemic lupus erythematosus. <i>Korean Journal of Transplantation</i> , 2021 , 35, S31-S31	0.2	
127	Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Monocentric Study. <i>Journal of Clinical Rheumatology</i> , 2021 , 27, e330-e335	1.1	2
126	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021 , 40, 3703-3710	3.9	0
125	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 441-446	2.5	0
124	Association between Follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021 , 134, 1168-1174	2.9	
123	Total Haemolytic Complement Activity at Diagnosis as an Indicator of the Baseline Activity of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2021 , 28, 85-93	1.2	1
122	Detection of intracellular monosodium urate crystals in gout synovial fluid using optical diffraction tomography. <i>Scientific Reports</i> , 2021 , 11, 10019	4.9	2
121	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 731-741	2.5	1
120	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23885	3	1

119	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021 , 75, e14512	2.9	0
118	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021 , 40, 1423-1430	3.9	0
117	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021 , 53, 1631-1638	2.3	1
116	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021 , 35, 1761-1768	2.3	0
115	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021 , 21, 399-406	4.9	1
114	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021 , 35, e23731	3	2
113	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021 , 2021, 6668884	4.3	0
112	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021 , 16, e0254226	3.7	0
111	Efficacy of tacrolimus as maintenance therapy after cyclophosphamide for treating antineutrophil cytoplasmic antibody-associated vasculitis. <i>Medicine (United States)</i> , 2021 , 100, e26956	1.8	
110	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021 , 36, 1221-1232	2.5	1
109	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , 2021 , 76, e2501	2.3	
108	Antineutrophil Cytoplasmic Antibody Positivity Is Associated with Vascular Involvement in Behçet's Disease. <i>Yonsei Medical Journal</i> , 2021 , 62, 149-158	3	1
107	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021 , 36, e120	4.7	1
106	The significance of cytoplasmic antinuclear antibody patterns in autoimmune liver disease. <i>PLoS ONE</i> , 2021 , 16, e0244950	3.7	2
105	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021 , 62, 494-502	3	1
104	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39 Suppl 129, 88-93	2.2	1
103	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39 Suppl 129, 94-100	2.2	
102	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39 Suppl 131, 111-118	2.2	

101	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 94-100	2.2	
100	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 88-93	2.2	4
99	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021 , 39, 111-118	2.2	
98	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis.. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
97	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace. <i>Annals of the Rheumatic Diseases</i> , 2020 ,	2.4	7
96	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020 , 39, 3009-3015	3.9	2
95	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020 , 76, 407-412	3	0
94	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020 , 20, 147	3.5	2
93	Clinical characteristics and long-term outcomes of Libman-Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020 , 29, 1115-1120	2.6	1
92	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 138	5.7	2
91	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020 , 21, 258	2.7	2
90	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 251, 117-123	2.4	3
89	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020 , 61, 951-957	3	2
88	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020 , 61, 712-719	3	3
87	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020 , 61, 720-725	3	2
86	Lipid Profiles in Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis: A Cross-sectional Analysis. <i>Journal of Rheumatic Diseases</i> , 2020 , 27, 261-269	1.2	3
85	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020 , 61, 970-975	3	1
84	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients: Results From the Korean College of Rheumatology Biologics Registry. <i>Journal of Clinical Rheumatology</i> , 2020 , Publish Ahead of Print,	1.1	2

83	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020 , 40, 1063-1070	3.6	3
82	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020 , 74, e13431	2.9	2
81	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020 , 39, 831-840	3.9	1
80	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020 , 19, 184	4.4	4
79	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020 , 2020, 8847306	3.2	0
78	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020 , 34, e23022	3	5
77	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020 , 40, 1089-1099	3.6	4
76	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020 , 50, 1414-1420	5.3	2
75	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020 , 22, 73	5.7	4
74	MiR-451 suppresses inflammatory responses in ankylosing spondylitis by targeting macrophage migration inhibitory factor. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 275-281	2.2	4
73	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 236-237	2.2	2
72	Application of the 2019 classification criteria for systemic lupus erythematosus to patients with established ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 243-244	2.2	
71	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019 , 39, 1859-1866	3.6	3
70	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019 , 39, 1937-1944	3.6	7
69	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019 , 19, 90	3	8
68	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019 , 39, 1359-1367	3.6	2
67	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019 , 60, 454-460	3	6
66	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019 , 39, 885-892	3.6	1

65	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019 , 20, e299-e308	4.9	0
64	Uterine Artery Embolization in Patients With Autoimmune Disease: A Matched Case-Control Study. <i>American Journal of Roentgenology</i> , 2019 , 1-6	5.4	0
63	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019 , 38, 1685-1690	3.9	4
62	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019 , 22, 797-804	2.3	8
61	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019 , 39, 879-884	3.6	3
60	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019 , 38, 3485-3491	3.9	3
59	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019 , 14, e0219970	3.7	0
58	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019 , 39, 1899-1905	3.6	2
57	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019 , 2019, 7508240	3	5
56	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019 , 11,	6.7	5
55	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010-2018. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
54	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019 , 60, 1164-1173 ³		4
53	Metabolic Reprogramming by the Excessive AMPK Activation Exacerbates Antigen-Specific Memory CD8 T Cell Differentiation after Acute Lymphocytic Choriomeningitis Virus Infection. <i>Immune Network</i> , 2019 , 19, e11	6.1	3
52	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019 , 39, 395-401	1.5	1
51	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 487-494	3.9	9
50	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019 , 23, 223-230	2.5	12
49	Systemic immune-inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019 , 24, 711-717	2.2	27
48	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019 , 38, 1147-1154	3.9	19

47	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37 Suppl 117, 116-121	2.2	2
46	Risk of Primary Spontaneous Pneumothorax According to Chest Configuration. <i>Thoracic and Cardiovascular Surgeon</i> , 2018 , 66, 583-588	1.6	5
45	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018 , 37, 2133-2141	3.9	10
44	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018 , 38, 1095-1101	3.6	7
43	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 294-302	3	4
42	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018 , 19, 187	2.7	21
41	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018 , 59, 452-456	3	16
40	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018 , 59, 397-405	3	12
39	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018 , 38, 1865-1871	3.6	21
38	Serum leucine-rich β -glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018 , 486, 253-258	6.2	6
37	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018 , 37, 1333-1340	3.9	5
36	Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018 , 25, 131	1.2	1
35	Treat-to-Target Strategy for Asian Patients with Early Rheumatoid Arthritis: Result of a Multicenter Trial in Korea. <i>Journal of Korean Medical Science</i> , 2018 , 33, e346	4.7	3
34	A novel antimicrobial peptide acting via formyl peptide receptor 2 shows therapeutic effects against rheumatoid arthritis. <i>Scientific Reports</i> , 2018 , 8, 14664	4.9	12
33	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology Biologics (KOBIO) registry. <i>PLoS ONE</i> , 2018 , 13, e0205651	3.7	8
32	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 165-166	2.2	2
31	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 85-87	2.2	12
30	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 115-120	2.2	2

29	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 533-539	2.2	2
28	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 73-77	2.2	3
27	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36 Suppl 111, 65-72	2.2	8
26	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 261-268	2.3	7
25	Five factor score of more than 1 is associated with relapse during the first 2 year-follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1261-1268	2.3	9
24	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017 , 47, 216-221	5.3	21
23	Evaluation of Spleen Glucose Metabolism Using F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017 , 58, 507-513	8.9	24
22	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017 , 44, 996-1003	4.1	31
21	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017 , 56, 795-801	3.9	6
20	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1028-1035	2.3	24
19	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017 , 37, 685-694	3.6	11
18	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017 , 36, 2751-2759	3.9	8
17	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017 , 12, e0184333	3.7	3
16	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017 , 19, 193	5.7	9
15	Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017 , 24, 143	1.2	
14	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017 , 58, 1128-1134	3	5
13	The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017 , 96, e7901	1.8	4
12	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017 , 20, 1593-1605	2.3	12

11	Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017 , 96, e7431	1.8	6
10	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017 , 84, 583-587	2.9	8
9	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017 , 12, e0181590	3.7	7
8	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 47-54	2.2	10
7	Differential expressions of NOD-like receptors and their associations with inflammatory responses in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35, 630-637	2.2	17
6	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017 , 35 Suppl 103, 111-118	2.2	13
5	Brain meningioma in a patient with systemic lupus erythematosus. <i>Yeungnam University Journal of Medicine</i> , 2016 , 33, 159	1.3	1
4	Anti-Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016 , 19, 897-902	2.3	16
3	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , 2015 , 17, 77	5.7	15
2	Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, S68-74	2.2	4
1	Active tuberculosis risk with tumor necrosis factor inhibitors after treating latent tuberculosis. <i>Journal of Clinical Rheumatology</i> , 2014 , 20, 68-73	1.1	9