

Carmen Diniz

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9416477/carmen-diniz-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

47
papers

1,030
citations

14
h-index

31
g-index

53
ext. papers

1,207
ext. citations

4.9
avg, IF

4.22
L-index

#	Paper	IF	Citations
47	Impact of the PdSpm (Spermine) Complex on the Metabolism of Triple-Negative Breast Cancer Tumors of a Xenograft Mouse Model. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
46	Insights into Nuclear G-Protein-Coupled Receptors as Therapeutic Targets in Non-Communicable Diseases. <i>Pharmaceuticals</i> , 2021 , 14,	5.2	3
45	Implication of RAS in Postnatal Cardiac Remodeling, Fibrosis and Dysfunction Induced by Fetal Undernutrition.. <i>Pathophysiology</i> , 2021 , 28, 273-290	1.8	1
44	Novel Insights into Mice Multi-Organ Metabolism upon Exposure to a Potential Anticancer Pd(II)-Agent. <i>Metabolites</i> , 2021 , 11,	5.6	3
43	Preclinical Pharmacokinetics and Biodistribution of Anticancer Dinuclear Palladium(II)-Spermine Complex (PdSpm) in Mice. <i>Pharmaceuticals</i> , 2021 , 14,	5.2	6
42	The health-promoting potential of peptides from brewing by-products: An up-to-date review. <i>Trends in Food Science and Technology</i> , 2021 , 118, 143-143	15.3	1
41	Insights into sympathetic nervous system and GPCR interplay in fetal programming of hypertension: a bridge for new pharmacological strategies. <i>Drug Discovery Today</i> , 2020 , 25, 739-747	8.8	2
40	Experimental and Clinical Evidence of Endothelial Dysfunction in Inflammatory Bowel Disease. <i>Current Pharmaceutical Design</i> , 2020 , 26, 3733-3747	3.3	0
39	Adenosine A and A Receptors as Targets for the Treatment of Hypertensive-Diabetic Nephropathy. <i>Biomedicines</i> , 2020 , 8,	4.8	3
38	Fast and reliable ICP-MS quantification of palladium and platinum-based drugs in animal pharmacokinetic and biodistribution studies. <i>Analytical Methods</i> , 2020 , 12, 4806-4812	3.2	5
37	Nuclear G-protein-coupled receptors as putative novel pharmacological targets. <i>Drug Discovery Today</i> , 2019 , 24, 2192-2201	8.8	13
36	Diabetes downregulates renal adenosine A2A receptors in an experimental model of hypertension. <i>PLoS ONE</i> , 2019 , 14, e0217552	3.7	6
35	Anticancer activity of palladium-based complexes against triple-negative breast cancer. <i>Drug Discovery Today</i> , 2019 , 24, 1044-1058	8.8	54
34	Delphinidin-3-O-glucoside inhibits angiogenesis via VEGFR2 downregulation and migration through actin disruption. <i>Journal of Functional Foods</i> , 2019 , 54, 393-402	5.1	7
33	Multi-Organ NMR Metabolomics to Assess In Vivo Overall Metabolic Impact of Cisplatin in Mice. <i>Metabolites</i> , 2019 , 9,	5.6	7
32	Vascular angiotensin AT1 receptor neuromodulation in fetal programming of hypertension. <i>Vascular Pharmacology</i> , 2019 , 117, 27-34	5.9	9
31	Vascular impairment of adenosinergic system in hypertension: increased adenosine bioavailability and differential distribution of adenosine receptors and nucleoside transporters. <i>Histochemistry and Cell Biology</i> , 2019 , 151, 407-418	2.4	2

30	Adenosine Receptor Ligands on Cancer Therapy: A Review of Patent Literature. <i>Recent Patents on Anti-Cancer Drug Discovery</i> , 2018 , 13, 40-69	2.6	13
29	New insights into the antiangiogenic and proangiogenic properties of dietary polyphenols. <i>Molecular Nutrition and Food Research</i> , 2017 , 61, 1600912	5.9	18
28	The Adenosinergic System as a Therapeutic Target in the Vasculature: New Ligands and Challenges. <i>Molecules</i> , 2017 , 22,	4.8	18
27	Anti-Invasive and Anti-Proliferative Synergism between Docetaxel and a Polynuclear Pd-Spermine Agent. <i>PLoS ONE</i> , 2016 , 11, e0167218	3.7	13
26	Endothelial dysfunction impairs vascular neurotransmission in tail arteries. <i>Neurochemistry International</i> , 2015 , 80, 7-13	4.4	7
25	Inosine strongly enhances proliferation of human C32 melanoma cells through PLC-PKC-MEK1/2-ERK1/2 and PI3K pathways. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2015 , 116, 25-36	3.1	13
24	Endothelial and Neuronal Nitric Oxide Activate Distinct Pathways on Sympathetic Neurotransmission in Rat Tail and Mesenteric Arteries. <i>PLoS ONE</i> , 2015 , 10, e0129224	3.7	8
23	Combination of Cl-IB-MECA with paclitaxel is a highly effective cytotoxic therapy causing mTOR-dependent autophagy and mitotic catastrophe on human melanoma cells. <i>Journal of Cancer Research and Clinical Oncology</i> , 2014 , 140, 921-35	4.9	11
22	The combination of Cl-IB-MECA with paclitaxel: a new anti-metastatic therapeutic strategy for melanoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2014 , 74, 847-60	3.5	8
21	Lack of endogenous adenosine tonus on sympathetic neurotransmission in spontaneously hypertensive rat mesenteric artery. <i>PLoS ONE</i> , 2014 , 9, e105540	3.7	14
20	Potential of cytotoxicity of paclitaxel in combination with Cl-IB-MECA in human C32 metastatic melanoma cells: A new possible therapeutic strategy for melanoma. <i>Biomedicine and Pharmacotherapy</i> , 2013 , 67, 777-89	7.5	12
19	Differential inhibition of noradrenaline release mediated by inhibitory A ₁ adenosine receptors in the mesenteric vein and artery from normotensive and hypertensive rats. <i>Neurochemistry International</i> , 2013 , 62, 399-405	4.4	6
18	Impaired inhibitory function of presynaptic A ₁ -adenosine receptors in SHR mesenteric arteries. <i>Journal of Pharmacological Sciences</i> , 2013 , 122, 59-70	3.7	11
17	The anticancer properties of dietary polyphenols and its relation with apoptosis. <i>Current Pharmaceutical Design</i> , 2010 , 16, 114-34	3.3	12.2
16	PRESYNAPTIC ADENOSINE RECEPTORS IN SUPERIOR MESENTERIC ARTERY FROM SPONTANEOUSLY HYPERTENSIVE RATS: A FUNCTIONAL AND MORPHOLOGICAL STUDY: PP.29.169. <i>Journal of Hypertension</i> , 2010 , 28, e499-e500	1.9	1
15	Biologically relevant O,S-donor compounds. Synthesis, molybdenum complexation and xanthine oxidase inhibition. <i>Dalton Transactions</i> , 2008 , 1773-82	4.3	15
14	Ligands and therapeutic perspectives of adenosine A _{2A} receptors. <i>Current Pharmaceutical Design</i> , 2008 , 14, 1698-722	3.3	18
13	Immunohistochemical characterization of adenosine receptors in rat aorta and tail arteries. <i>Microscopy Research and Technique</i> , 2008 , 71, 703-9	2.8	16

12	Immunohistochemical localization of angiotensin II receptor types 1 and 2 in the mesenteric artery from spontaneously hypertensive rats. <i>Microscopy Research and Technique</i> , 2007 , 70, 677-81	2.8	6
11	A2A adenosine-receptor-mediated facilitation of noradrenaline release in rat tail artery involves protein kinase C activation and betagamma subunits formed after alpha2-adrenoceptor activation. <i>Neurochemistry International</i> , 2007 , 51, 47-56	4.4	14
10	New insights on the anticancer properties of dietary polyphenols. <i>Medicinal Research Reviews</i> , 2006 , 26, 747-66	14.4	414
9	Cytotoxic and COX-2 Inhibition Properties of Hydroxycinnamic Derivatives. <i>Letters in Drug Design and Discovery</i> , 2006 , 3, 316-320	0.8	10
8	Semiautomated computer-assisted image analysis to quantify 3,3'-diaminobenzidine tetrahydrochloride-immunostained small tissues. <i>Analytical Biochemistry</i> , 2006 , 357, 137-43	3.1	25
7	Regional differences in extracellular purine degradation in the prostatic and epididymal portions of the rat vas deferens. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2005 , 32, 721-7	3	9
6	Adenosine receptors involved in modulation of noradrenaline release in isolated rat tail artery. <i>European Journal of Pharmacology</i> , 2004 , 504, 17-25	5.3	18
5	Facilitation of noradrenaline release by activation of adenosine A(2A) receptors triggers both phospholipase C and adenylate cyclase pathways in rat tail artery. <i>Cardiovascular Research</i> , 2004 , 63, 739-46	9.9	33
4	Regional differences in the adenosine A(2) receptor-mediated modulation of contractions in rat vas deferens. <i>European Journal of Pharmacology</i> , 2003 , 460, 191-9	5.3	13
3	Facilitation of noradrenaline release by adenosine A(2A) receptors in the epididymal portion and adenosine A(2B) receptors in the prostatic portion of the rat vas deferens. <i>European Journal of Pharmacology</i> , 2002 , 448, 45-50	5.3	13
2	Release inhibitory receptors activation favours the A2A-adenosine receptor-mediated facilitation of noradrenaline release in isolated rat tail artery. <i>British Journal of Pharmacology</i> , 2002 , 136, 230-6	8.6	21
1	Taurine release in the rat vas deferens is modulated by Ca ²⁺ and is independent of contractions. <i>European Journal of Pharmacology</i> , 1999 , 376, 273-8	5.3	1