

Kousuke Watanabe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9417158/publications.pdf>

Version: 2024-02-01

26
papers

523
citations

623188

14
h-index

642321

23
g-index

26
all docs

26
docs citations

26
times ranked

991
citing authors

#	ARTICLE	IF	CITATIONS
1	Immune checkpoint inhibitor combination therapies very frequently induce secondary adrenal insufficiency. <i>Scientific Reports</i> , 2021, 11, 11617.	1.6	9
2	Dual EGFR and ABL Tyrosine Kinase Inhibitor Treatment in a Patient with Concomitant EGFR-Mutated Lung Adenocarcinoma and BCR-ABL1-Positive CML. <i>Case Reports in Oncological Medicine</i> , 2020, 2020, 1-6.	0.2	2
3	Integrin $\alpha 11$ in non-small cell lung cancer is associated with tumor progression and postoperative recurrence. <i>Cancer Science</i> , 2020, 111, 200-208.	1.7	24
4	NRXN1 as a novel potential target of antibody-drug conjugates for small cell lung cancer. <i>Oncotarget</i> , 2020, 11, 3590-3600.	0.8	8
5	Rapid temporal improvement of pembrolizumab-induced pneumonitis using the anti-TNF- α antibody infliximab. <i>Drug Discoveries and Therapeutics</i> , 2019, 13, 164-167.	0.6	14
6	Spontaneous Transdifferentiation from Small Cell Lung Carcinoma to Squamous Cell Carcinoma. <i>Journal of Thoracic Oncology</i> , 2019, 14, e31-e34.	0.5	3
7	Development of mild drug-induced sclerosing cholangitis after discontinuation of nivolumab. <i>European Journal of Cancer</i> , 2019, 107, 93-96.	1.3	17
8	Identification of a metastatic lung adenocarcinoma of the palate mucosa through genetic and histopathological analysis: a rare case report and literature review. <i>BMC Cancer</i> , 2019, 19, 52.	1.1	5
9	Glutamate-cysteine ligase catalytic subunit is associated with cisplatin resistance in lung adenocarcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 303-307.	0.6	21
10	Altered editing level of microRNAs is a potential biomarker in lung adenocarcinoma. <i>Cancer Science</i> , 2018, 109, 3326-3335.	1.7	32
11	Mean platelet volume and lymphocyte-to-monocyte ratio are associated with shorter progression-free survival in EGFR-mutant lung adenocarcinoma treated by EGFR tyrosine kinase inhibitor. <i>PLoS ONE</i> , 2018, 13, e0203625.	1.1	30
12	High expression of IRE1 in lung adenocarcinoma is associated with a lower rate of recurrence. <i>Japanese Journal of Clinical Oncology</i> , 2017, 47, 543-550.	0.6	7
13	Histone methylation-mediated silencing of miR-139 enhances invasion of non-small cell lung cancer. <i>Cancer Medicine</i> , 2015, 4, 1573-1582.	1.3	41
14	The chimeric transcript RUNX1-GLRX5: a biomarker for good postoperative prognosis in Stage IA non-small-cell lung cancer. <i>Japanese Journal of Clinical Oncology</i> , 2015, 46, 187.	0.6	1
15	Oncogenic TPM3-ALK activation requires dimerization through the coiled-coil structure of TPM3. <i>Biochemical and Biophysical Research Communications</i> , 2015, 457, 457-460.	1.0	20
16	Alternative polyadenylation is associated with lower expression of PABPN1 and poor prognosis in non-small cell lung cancer. <i>Cancer Science</i> , 2014, 105, 1135-1141.	1.7	36
17	Detection of novel paralogous ring finger tyrosine kinase mRNA chimeras is associated with poor postoperative prognosis in non-small cell lung cancer. <i>Cancer Science</i> , 2013, 104, 1447-1454.	1.7	12
18	Disruption of the expression and function of microRNAs in lung cancer as a result of epigenetic changes. <i>Frontiers in Genetics</i> , 2013, 4, 275.	1.1	17

#	ARTICLE	IF	CITATIONS
19	FER overexpression is associated with poor postoperative prognosis and cancer-cell survival in non-small cell lung cancer. <i>International Journal of Clinical and Experimental Pathology</i> , 2013, 6, 598-612.	0.5	16
20	Pulmonary Venous Invasion, Determined by Chest Computed Tomographic Scan, as a Potential Early Indicator of Zygomycosis Infection. <i>Journal of Thoracic Imaging</i> , 2012, 27, W97-W99.	0.8	0
21	Genome structure-based screening identified epigenetically silenced microRNA associated with invasiveness in non-small cell lung cancer. <i>International Journal of Cancer</i> , 2012, 130, 2580-2590.	2.3	83
22	CpG island methylation of microRNAs is associated with tumor size and recurrence of non-small cell lung cancer. <i>Cancer Science</i> , 2011, 102, 2126-2131.	1.7	33
23	Polymerase reaction without primers throughout for the reconstruction of full-length cDNA from products of rapid amplification of cDNA ends (RACE). <i>Biotechnology Letters</i> , 2011, 33, 1301-1307.	1.1	4
24	Identification of <i>GOS2</i> as a gene frequently methylated in squamous lung cancer by combination of <i>in silico</i> and experimental approaches. <i>International Journal of Cancer</i> , 2010, 126, 1895-1902.	2.3	48
25	Treatment of PCR products with exonuclease I and heat-labile alkaline phosphatase improves the visibility of combined bisulfite restriction analysis. <i>Biochemical and Biophysical Research Communications</i> , 2010, 399, 422-424.	1.0	9
26	Impact of DNA demethylation of the <i>GOS2</i> gene on the transcription of <i>GOS2</i> in squamous lung cancer cell lines with or without nuclear receptor agonists. <i>Biochemical and Biophysical Research Communications</i> , 2009, 390, 1283-1287.	1.0	31