

Michael Hultström

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9419468/michael-hultstrom-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

106
papers

1,070
citations

19
h-index

28
g-index

126
ext. papers

1,607
ext. citations

4.5
avg, IF

4.97
L-index

#	Paper	IF	Citations
106	The Contribution of Plasma Urea to Total Osmolality During Iatrogenic Fluid Reduction in Critically Ill Patients.. <i>Function</i> , 2022 , 3, zqab055	6.1	2
105	Soluble TNF receptors predict acute kidney injury and mortality in critically ill COVID-19 patients: A prospective observational study. <i>Cytokine</i> , 2022 , 149, 155727	4	5
104	Infectious SARS-CoV-2 is rarely present in the nasopharynx samples collected from Swedish hospitalized critically ill COVID-19 patients.. <i>Irish Journal of Medical Science</i> , 2022 , 1	1.9	
103	Neutrophil extracellular traps promote cancer-associated inflammation and myocardial stress.. <i>Oncotmmunology</i> , 2022 , 11, 2049487	7.2	0
102	Limitations of the ARDS criteria during high-flow oxygen or non-invasive ventilation: evidence from critically ill COVID-19 patients.. <i>Critical Care</i> , 2022 , 26, 55	10.8	0
101	How the Innate Immune System of the Blood Contributes to Systemic Pathology in COVID-19-Induced ARDS and Provides Potential Targets for Treatment.. <i>Frontiers in Immunology</i> , 2022 , 13, 840137	8.4	1
100	The Evolution of Blood Cell Phenotypes, Intracellular and Plasma Cytokines and Morphological Changes in Critically Ill COVID-19 Patients. <i>Biomedicines</i> , 2022 , 10, 934	4.8	0
99	Surgical trauma is associated with renal immune cell activation in rats: A microarray study. <i>Physiological Reports</i> , 2021 , 9, e15142	2.6	1
98	Common, low-frequency, rare, and ultra-rare coding variants contribute to COVID-19 severity. <i>Human Genetics</i> , 2021 , 141, 147	6.3	3
97	Plasma endostatin correlates with hypoxia and mortality in COVID-19-associated acute respiratory failure. <i>Biomarkers in Medicine</i> , 2021 , 15, 1509-1517	2.3	
96	Plasma hyaluronan, hyaluronidase activity and endogenous hyaluronidase inhibition in sepsis: an experimental and clinical cohort study. <i>Intensive Care Medicine Experimental</i> , 2021 , 9, 53	3.7	0
95	Intensive care-treated COVID-19 patients' perception of their illness and remaining symptoms. <i>Acta Anaesthesiologica Scandinavica</i> , 2021 ,	1.9	2
94	The impact of viremia on organ failure, biomarkers and mortality in a Swedish cohort of critically ill COVID-19 patients. <i>Scientific Reports</i> , 2021 , 11, 7163	4.9	10
93	Case report: An unusual presentation of renal hypertension after damage control surgery. <i>International Journal of Surgery Case Reports</i> , 2021 , 82, 105872	0.8	
92	Point of care ultrasound screening for deep vein thrombosis in critically ill COVID-19 patients, an observational study. <i>Thrombosis Journal</i> , 2021 , 19, 38	5.6	1
91	Impaired diffusing capacity for carbon monoxide is common in critically ill Covid-19 patients at four months post-discharge. <i>Respiratory Medicine</i> , 2021 , 182, 106394	4.6	9
90	COVID-19 patients in intensive care develop predominantly oliguric acute kidney injury. <i>Acta Anaesthesiologica Scandinavica</i> , 2021 , 65, 364-372	1.9	16

89	Increased levels of plasma cytokines and correlations to organ failure and 30-day mortality in critically ill Covid-19 patients. <i>Cytokine</i> , 2021 , 138, 155389	4	24
88	High expression of neutrophil and monocyte CD64 with simultaneous lack of upregulation of adhesion receptors CD11b, CD162, CD15, CD65 on neutrophils in severe COVID-19. <i>Therapeutic Advances in Infectious Disease</i> , 2021 , 8, 20499361211034065	2.8	2
87	Severe acute kidney injury associated with progression of chronic kidney disease after critical COVID-19. <i>Critical Care</i> , 2021 , 25, 37	10.8	13
86	The Outcome of Critically Ill COVID-19 Patients Is Linked to Thromboinflammation Dominated by the Kallikrein/Kinin System. <i>Frontiers in Immunology</i> , 2021 , 12, 627579	8.4	16
85	The swedish covid-19 intensive care cohort: Risk factors of ICU admission and ICU mortality. <i>Acta Anaesthesiologica Scandinavica</i> , 2021 , 65, 525	1.9	24
84	A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity. <i>Nature Medicine</i> , 2021 , 27, 659-667	50.5	52
83	Critical illness polyneuropathy, myopathy and neuronal biomarkers in COVID-19 patients: A prospective study. <i>Clinical Neurophysiology</i> , 2021 , 132, 1733-1740	4.3	40
82	Evolution of NETosis markers and DAMPs have prognostic value in critically ill COVID-19 patients. <i>Scientific Reports</i> , 2021 , 11, 15701	4.9	20
81	A quantitative analysis of extension and distribution of lung injury in COVID-19: a prospective study based on chest computed tomography. <i>Critical Care</i> , 2021 , 25, 276	10.8	2
80	Histone H3 Cleavage in Severe COVID-19 ICU Patients. <i>Frontiers in Cellular and Infection Microbiology</i> , 2021 , 11, 694186	5.9	4
79	Urinary cytokines correlate with acute kidney injury in critically ill COVID-19 patients. <i>Cytokine</i> , 2021 , 146, 155589	4	4
78	Plasma Leptin Is Increased in Intensive Care Patients with COVID-19-An Investigation Performed in the PronMed-Cohort.. <i>Biomedicines</i> , 2021 , 10,	4.8	2
77	ECG pathology and its association with death in critically ill COVID-19 patients, a cohort study.. <i>PLoS ONE</i> , 2021 , 16, e0261315	3.7	3
76	Optimal cutting temperature medium embedding and cryostat sectioning are valid for cardiac myofilament function assessment. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2020 , 319, H235-H241	5.2	
75	Patient satisfaction with continuous epidural analgesia after major surgical procedures at a Swedish University hospital. <i>PLoS ONE</i> , 2020 , 15, e0235636	3.7	0
74	Analgesic effects of dexmedetomidine and remifentanyl on periprocedural pain during percutaneous ablation of renal carcinoma. <i>Uppsala Journal of Medical Sciences</i> , 2020 , 125, 52-57	2.8	3
73	Presence of SARS-CoV-2 in urine is rare and not associated with acute kidney injury in critically ill COVID-19 patients. <i>Critical Care</i> , 2020 , 24, 587	10.8	17
72	Optimal Cutting Temperature Medium-Embedding Is a Valid Method for Storing and Preparing Myocardial Biopsies Preceding Myofilament Function-Assessment. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	

71	Inadequate prophylactic effect of low-molecular weight heparin in critically ill COVID-19 patients. <i>Journal of Critical Care</i> , 2020 , 60, 249-252	4	15
70	Weak anti-SARS-CoV-2 antibody response is associated with mortality in a Swedish cohort of COVID-19 patients in critical care. <i>Critical Care</i> , 2020 , 24, 639	10.8	5
69	Hyperreninemia and low total body water may contribute to acute kidney injury in COVID-19 patients in intensive care. <i>Journal of Hypertension</i> , 2020 , 38, 1613-1614	1.9	8
68	Blood type A associates with critical COVID-19 and death in a Swedish cohort. <i>Critical Care</i> , 2020 , 24, 496	10.8	13
67	Mannose-Binding Lectin is Associated with Thrombosis and Coagulopathy in Critically Ill COVID-19 Patients. <i>Thrombosis and Haemostasis</i> , 2020 , 120, 1720-1724	7	39
66	Quantitative trait loci associated with angiotensin II and high-salt diet induced acute decompensated heart failure in Balb/CJ mice. <i>Physiological Genomics</i> , 2019 , 51, 279-289	3.6	1
65	Time course of decompensation after angiotensin II and high-salt diet in Balb/CJ mice suggests pulmonary hypertension-induced cardiorenal syndrome. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2019 , 316, R563-R570	3.2	5
64	Angiotensin II and salt-induced decompensation in Balb/CJ mice is aggravated by fluid retention related to low oxidative stress. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, F914-F933	4.3	3
63	Release of a contractile factor and reduced nitric oxide from isolated pulmonary resistance vessels from Balb/CJ mice cause higher reactivity to angiotensin II compared to C57BL/6J mice. <i>FASEB Journal</i> , 2019 , 33, 550.10	0.9	
62	ADAMTS13 protects mice against renal ischemia-reperfusion injury by reducing inflammation and improving endothelial function. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, F134-F145	4.3	13
61	NFAT5 regulates renal gene expression in response to angiotensin II through Annexin-A2-mediated posttranscriptional regulation in hypertensive rats. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, F101-F112	4.3	5
60	Comparison of acute kidney injury of different etiology reveals in-common mechanisms of tissue damage. <i>Physiological Genomics</i> , 2018 , 50, 127-141	3.6	28
59	Losartan does not decrease renal oxygenation and norepinephrine effects in rats after resuscitated hemorrhage. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 315, F241-F246	4.3	1
58	In-Common And Unique Gene Expression Patterns In Acute Kidney Injury Of Different Aetiology Implicates MYC-Pathway In Damage Progression. <i>FASEB Journal</i> , 2018 , 32, 849.7	0.9	
57	BALB/CJBom Treated with Angiotensin II and High Salt Diet Develop Pulmonary Hypertension and Right Sided Heart Failure while C57BL/6J Mice do not. <i>FASEB Journal</i> , 2018 , 32, 892.10	0.9	
56	ICU mortality and variables associated with ICU survival in Poland: A nationwide database study. <i>European Journal of Anaesthesiology</i> , 2018 , 35, 949-954	2.3	14
55	Osthole Ameliorates Renal Fibrosis in Mice by Suppressing Fibroblast Activation and Epithelial-Mesenchymal Transition. <i>Frontiers in Physiology</i> , 2018 , 9, 1650	4.6	18
54	Mortality rate is higher in Polish intensive care units than in other European countries. <i>Intensive Care Medicine</i> , 2017 , 43, 1430-1432	14.5	15

53	Mortality rate in Polish intensive care units is lower than predicted according to the APACHE II scoring system. <i>Intensive Care Medicine</i> , 2017 , 43, 1745-1746	14.5	16
52	c-Jun N-terminal Kinase mediates prostaglandin-induced sympathoexcitation in rats with chronic heart failure by reducing GAD1 and GABRA1 expression. <i>Acta Physiologica</i> , 2017 , 219, 494-509	5.6	2
51	Sympathoexcitation in Rats With Chronic Heart Failure Depends on Homeobox D10 and MicroRNA-7b Inhibiting GABBR1 Translation in Paraventricular Nucleus. <i>Circulation: Heart Failure</i> , 2016 , 9, e002261	7.6	5
50	Unilateral renal ischaemia in rats induces a rapid secretion of inflammatory markers to renal lymph and increased capillary permeability. <i>Journal of Physiology</i> , 2016 , 594, 1709-26	3.9	11
49	Increased hydrogen peroxide impairs angiotensin II contractions of afferent arterioles in mice after renal ischaemia-reperfusion injury. <i>Acta Physiologica</i> , 2016 , 218, 136-45	5.6	23
48	Distinct protein signature of hypertension-induced damage in the renal proteome of the two-kidney, one-clip rat model. <i>Journal of Hypertension</i> , 2015 , 33, 126-35	1.9	12
47	Noradrenaline enhances angiotensin II responses via p38 MAPK activation after hypoxia/re-oxygenation in renal interlobar arteries. <i>Acta Physiologica</i> , 2015 , 213, 920-32	5.6	11
46	Commentaries on Viewpoint: Can elite athletes benefit from dietary nitrate supplementation?. <i>Journal of Applied Physiology</i> , 2015 , 119, 762-9	3.7	13
45	Identification of a common molecular pathway in hypertensive renal damage: comparison of rat and human gene expression profiles. <i>Journal of Hypertension</i> , 2015 , 33, 584-96; discussion 596	1.9	7
44	Matrix Metalloproteinase-2 Knockout and Heterozygote Mice Are Protected from Hydronephrosis and Kidney Fibrosis after Unilateral Ureteral Obstruction. <i>PLoS ONE</i> , 2015 , 10, e0143390	3.7	21
43	Validation of uromodulin as a candidate gene for human essential hypertension. <i>Hypertension</i> , 2014 , 63, 551-8	8.5	83
42	Intradermal insulin delivery: a promising future for diabetes management. <i>Journal of Diabetes Science and Technology</i> , 2014 , 8, 453-7	4.1	25
41	Renal neurohormonal regulation in heart failure decompensation. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2014 , 307, R493-7	3.2	28
40	Ingrid Toft (June 2, 1959-April 26, 2014). <i>Blood Pressure</i> , 2014 , 23, 255	1.7	
39	Sex-specific prevalence of fatty liver disease and associated metabolic factors in Wuhan, south central China. <i>European Journal of Gastroenterology and Hepatology</i> , 2014 , 26, 1015-21	2.2	25
38	Lower oxidative stress is associated with angiotensin II and salt-induced acute cardiorenal failure in BalbC mice but not C57Black6 (860.10). <i>FASEB Journal</i> , 2014 , 28, 860.10	0.9	
37	Nucleic acid binding of annexin A2 is regulated through angiotensin II/AT1 signaling in kidneys of hypertensive rats (1088.2). <i>FASEB Journal</i> , 2014 , 28, 1088.2	0.9	
36	Genomic differences in glutathione metabolism determines susceptibility to cardiorenal failure in mice (860.11). <i>FASEB Journal</i> , 2014 , 28, 860.11	0.9	

35	Prevalence and associated metabolic factors of fatty liver disease in the elderly. <i>Experimental Gerontology</i> , 2013 , 48, 705-9	4.5	45
34	Arterial damage precedes the development of interstitial damage in the nonclipped kidney of two-kidney, one-clip hypertensive rats. <i>Journal of Hypertension</i> , 2013 , 31, 152-9	1.9	15
33	Nitric oxide in afferent arterioles after uninephrectomy depends on extracellular L-arginine. <i>American Journal of Physiology - Renal Physiology</i> , 2013 , 304, F1088-98	4.3	2
32	Neurohormonal interactions on the renal oxygen delivery and consumption in haemorrhagic shock-induced acute kidney injury. <i>Acta Physiologica</i> , 2013 , 209, 11-25	5.6	21
31	Renal ischemia-reperfusion (I/R) injury induces a rapid activation of local inflammatory markers and causes increased peritubular permeability.. <i>FASEB Journal</i> , 2013 , 27, 682.10	0.9	
30	Attenuated contractility in afferent arterioles during development of proteinuria in two-kidney, one-clip hypertensive rats. <i>FASEB Journal</i> , 2013 , 27, 1110.15	0.9	
29	Proteomic analysis of outer and juxtamedullary cortex of non-clipped kidneys in 2K1C hypertensive rats. <i>FASEB Journal</i> , 2013 , 27, 909.15	0.9	
28	Collagen-binding proteins in age-dependent changes in renal collagen turnover: microarray analysis of mRNA expression. <i>Physiological Genomics</i> , 2012 , 44, 576-86	3.6	5
27	Development of structural kidney damage in spontaneously hypertensive rats. <i>Journal of Hypertension</i> , 2012 , 30, 1087-91	1.9	57
26	MMP2 deficient mice are protected from hydronephrosis after unilateral urethral obstruction. <i>FASEB Journal</i> , 2012 , 26, 868.12	0.9	1
25	Renal extracellular matrix in three rat-models of hypertensive kidney damage: A microarray study of SHR, SHRSP and 2K1C. <i>FASEB Journal</i> , 2012 , 26, 872.32	0.9	
24	Adenosine sensitization after angiotensin II stimulation in afferent arterioles from normal rats does not occur during two-kidney, one-clip hypertension. <i>Acta Physiologica</i> , 2011 , 201, 289-94	5.6	7
23	Afferent arteriopathy and glomerular collapse but not segmental sclerosis induce tubular atrophy in old spontaneously hypertensive rats. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2011 , 459, 99-108	5.1	23
22	Osteopontin is upregulated in damaged non-clipped kidney cortex from rats with renal hypertension. <i>FASEB Journal</i> , 2010 , 24, 791.4	0.9	
21	Trefoil factor-3 is down regulated while CYP24a1 is increased in the ageing rat kidney. <i>FASEB Journal</i> , 2010 , 24, 791.5	0.9	
20	Norepinephrine increases calcium sensitivity of mouse afferent arteriole, thereby enhancing angiotensin II-mediated vasoconstriction. <i>Kidney International</i> , 2009 , 76, 953-9	9.9	14
19	AT(1) receptor activation regulates the mRNA expression of CAT1, CAT2, arginase-1, and DDAH2 in preglomerular vessels from angiotensin II hypertensive rats. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F163-8	4.3	14
18	Angiotensin II-induced contraction is attenuated by nitric oxide in afferent arterioles from the nonclipped kidney in 2K1C. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 296, F78-86	4.3	22

17	Sympathectomy suppresses tumor growth and alters gene-expression profiles in rat tongue cancer. <i>European Journal of Oral Sciences</i> , 2009 , 117, 351-61	2.3	19
16	Renal vascular L-arginine metabolism, NO release and contraction in Angiotensin II hypertensive rats. <i>FASEB Journal</i> , 2009 , 23, 606.6	0.9	
15	Norepinephrine Treatment Enhances the Constriction of the Afferent Arterioles to Angiotensin II by Increasing the Calcium Sensitivity. <i>FASEB Journal</i> , 2009 , 23, 804.2	0.9	
14	Renal damage in the non-clipped kidney in two kidney one clip rat is most pronounced in the juxtamedullary cortex.. <i>FASEB Journal</i> , 2009 , 23, 1017.12	0.9	
13	Tetradecylthioacetic acid downregulates cyclooxygenase 2 in the renal cortex of two-kidney, one-clip hypertensive rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008 , 295, R1866-73	3.2	5
12	Upregulation of tissue inhibitor of metalloproteases-1 (TIMP-1) and procollagen-N-peptidase in hypertension-induced renal damage. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 896-903	4.3	26
11	Collagen metabolism and renal damage in 2k1c rats. <i>FASEB Journal</i> , 2008 , 22, 968.5	0.9	
10	Compensatory hyperfiltration and NO in 2k1c and uninephrectomized rats. <i>FASEB Journal</i> , 2008 , 22, 761.4	0.9	
9	Moderate hypothermia induces a preferential increase in pancreatic islet blood flow in anesthetized rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2007 , 293, R1438-43	3.2	1
8	Adenosine triphosphate increases the reactivity of the afferent arteriole to low concentrations of norepinephrine. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2007 , 293, R2225-31	3.2	10
7	Protein expression of factors involved in the development of renal interstitial fibrosis in old SHR. <i>FASEB Journal</i> , 2007 , 21, A899	0.9	
6	The mRNA expression of eNOS, iNOS and L-arginine transporters in the afferent arterioles (AA) of 2K1C hypertensive rats. <i>FASEB Journal</i> , 2007 , 21, A899	0.9	
5	Prevention of hypertension and organ damage in 2-kidney, 1-clip rats by tetradecylthioacetic acid. <i>Hypertension</i> , 2006 , 48, 460-6	8.5	17
4	Markers of NETosis and DAMPs are altered in critically ill COVID-19 patients		3
3	Critical Illness Polyneuropathy and Myopathy in COVID-19 Patients: A Prospective Observational Intensive Care Unit Cross-Sectional Cohort Study		4
2	A Neanderthal OAS1 isoform Protects Against COVID-19 Susceptibility and Severity: Results from Mendelian Randomization and Case-Control Studies		4
1	Elevated Angiotensin-2 inhibits thrombomodulin-mediated anticoagulation in critically ill COVID-19 patients		2