

# Keiko Arai

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9427435/publications.pdf>

Version: 2024-02-01

14  
papers

203  
citations

1307594

7  
h-index

1125743

13  
g-index

14  
all docs

14  
docs citations

14  
times ranked

228  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Cross sectional study on proportion of sulfonylureas among various oral antidiabetic drugs using for Japanese patients with type 2 diabetes, analyzed from NSAID Study-2. Diabetology International, 2022, 13, 169-176.   | 1.4 | 0         |
| 2  | Differences in Dental Care Referral for Diabetic Patients Between General Practitioners and Diabetes Specialists in Japan, Analyzed from NSAID-Study 3. Diabetes Therapy, 2022, 13, 379-385.  | 2.5 | 1         |
| 3  | Current status of oral antidiabetic drug prescribing patterns based on the body mass index for Japanese type 2 diabetes mellitus patients and yearly changes in diabetologists' prescribing patterns from 2002 to 2019 (JDDM61). Journal of Diabetes Investigation, 2021, , .                 | 2.4 | 14        |
| 4  | Nationwide Survey on Actual Interventions for Type 2 Diabetes by Japanese Practitioners (NSAID) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50  | 2.5 | 3         |
| 5  | Effect of canagliflozin on the overall clinical state including insulin resistance in Japanese patients with type 2 diabetes mellitus. Diabetes Research and Clinical Practice, 2019, 149, 140-146.   | 2.8 | 32        |
| 6  | Relationship between the efficacy of oral antidiabetic drugs and clinical features in type 2 diabetic patients (JDDM38). Journal of Diabetes Investigation, 2016, 7, 386-395.   | 2.4 | 15        |
| 7  | Both Glimpiride and High-Dose Metformin Are Important for Sustained Glucose Lowering in Japanese Type 2 Diabetes Patients on Glimpiride+Sitagliptin+Metformin Therapy: Subanalysis of a Single-Center, Open-Label, Randomized Study. Diabetes Technology and Therapeutics, 2014, 16, 442-446. | 4.4 | 1         |
| 8  | Glimpiride Strongly Enhances the Glucose-Lowering Effect in Triple Oral Antidiabetes Therapy with Sitagliptin and Metformin for Japanese Patients with Type 2 Diabetes Mellitus. Diabetes Technology and Therapeutics, 2013, 15, 335-341.   | 4.4 | 7         |
| 9  | Short duration of diabetes and disuse of sulfonylurea have any association with insulin cessation of the patients with type 2 diabetes in a clinical setting in Japan (JDDM 30). Endocrine Journal, 2013, 60, 305-310.  | 1.6 | 3         |
| 10 | Combination Therapy with a Dipeptidyl peptidase-4 Inhibitor, Sulfonylurea, and Metformin Markedly Improves HbA <sub>1c</sub> Levels in Japanese Patients with Type 2 Diabetes Mellitus. Japanese Clinical Medicine, 2012, 3, JCM.S8571.   | 1.9 | 6         |
| 11 | Present status of insulin therapy for type 2 diabetes treated by general practitioners and diabetes specialists in Japan: Third report of a cross-sectional survey of 15,652 patients. Journal of Diabetes Investigation, 2012, 3, 396-401.   | 2.4 | 10        |
| 12 | Present status of sulfonylurea treatment for type 2 diabetes in Japan: Second report of a cross-sectional survey of 15,652 patients. Endocrine Journal, 2010, 57, 499-507.  | 1.6 | 31        |
| 13 | The status of glycemic control by general practitioners and specialists for diabetes in Japan: A cross-sectional survey of 15,652 patients with diabetes mellitus. Diabetes Research and Clinical Practice, 2009, 83, 397-401.  | 2.8 | 37        |
| 14 | Six-month multicentric, open-label, randomized trial of twice-daily injections of biphasic insulin aspart 30 versus multiple daily injections of insulin aspart in Japanese type 2 diabetic patients (JDDM 11). Diabetes Research and Clinical Practice, 2008, 79, 171-176.                   | 2.8 | 43        |