

# Yong-Beom Park

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/9429330/yong-beom-park-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

206  
papers

2,152  
citations

21  
h-index

40  
g-index

242  
ext. papers

2,711  
ext. citations

3.7  
avg, IF

5.07  
L-index

#	Paper	IF	Citations
206	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis.. <i>Journal of Clinical Laboratory Analysis</i> , <b>2022</b> , e24237	3	1
205	Association Between Idiopathic Cutaneous Leukocytoclastic Angiitis and ANCA-negative Microscopic Polyangiitis. <i>Journal of Rheumatic Diseases</i> , <b>2022</b> , 29, 40-45	1.2	
204	Aortic valve surgery in patients with Takayasu's arteritis: a nationwide analysis of 1,197 patients during a 9-year period. <i>Clinical and Experimental Rheumatology</i> , <b>2022</b> , 40, 568-574	2.2	
203	New body mass index for predicting prognosis in patients with antineutrophil cytoplasmic antibody-associated vasculitis.. <i>Journal of Clinical Laboratory Analysis</i> , <b>2022</b> , e24357	3	0
202	Incidence and Patterns of Interstitial Lung Disease and Their Clinical Impact on Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Korean Single-Centre Observational Study. <i>Journal of Immunology Research</i> , <b>2022</b> , 2022, 1-7	4.5	0
201	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , <b>2021</b> , 75, e13929	2.9	1
200	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. <i>Clinical Rheumatology</i> , <b>2021</b> , 1	3.9	2
199	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , <b>2021</b> , 69, 91-95	2.9	
198	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2021</b> , 62, 1016-1022	3	1
197	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , <b>2021</b> , 35, e24048	3	1
196	Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Monocentric Study. <i>Journal of Clinical Rheumatology</i> , <b>2021</b> , 27, e330-e335	4.1	2
195	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , <b>2021</b> , 23, 77	5.7	0
194	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , <b>2021</b> , 40, 3703-3710	3.9	0
193	Risk of Stroke in Systemic Necrotizing Vasculitis: A Nationwide Study Using the National Claims Database. <i>Frontiers in Immunology</i> , <b>2021</b> , 12, 629902	8.4	0
192	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , <b>2021</b> , 36, 111-116	2.5	0
191	Unclassifiable repeated antineutrophil cytoplasmic antibody (ANCA) positivity in diseases other than ANCA-associated vasculitis. <i>Zeitschrift Fur Rheumatologie</i> , <b>2021</b> , 1	1.9	2
190	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , <b>2021</b> , 134, 1168-1174	2.9	

189	Total Haemolytic Complement Activity at Diagnosis as an Indicator of the Baseline Activity of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , <b>2021</b> , 28, 85-93 <sup>1,2</sup>		1
188	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , <b>2021</b> , 36, 731-741	2.5	1
187	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , <b>2021</b> , 35, e23885	3	1
186	Comparison of Retention Rates Between Tumor Necrosis Factor- $\alpha$ Inhibitors in Patients With Ankylosing Spondylitis: Data From the Korean College of Rheumatology Biologics Registry. <i>Frontiers in Medicine</i> , <b>2021</b> , 8, 689609	4.9	2
185	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , <b>2021</b> , 75, e14512	2.9	0
184	Increased prevalence rate of metabolic syndrome is an independent predictor of cardiovascular disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Rheumatology International</i> , <b>2021</b> , 1	3.6	2
183	Effect of statins on the prevention of recurrent thrombosis in thrombotic antiphospholipid syndrome. <i>Rheumatology</i> , <b>2021</b> ,	3.9	1
182	Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus. <i>Annals of the Rheumatic Diseases</i> , <b>2021</b> , 80, 632-640	2.4	3 <sup>1</sup>
181	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , <b>2021</b> , 40, 1423-1430	3.9	0
180	LBEC0101, an etanercept biosimilar for the treatment of rheumatoid arthritis. <i>Expert Opinion on Biological Therapy</i> , <b>2021</b> , 21, 1-8	5.4	
179	Serological Biomarkers and Indices for the Current Activity and Prognosis of ANCA-Associated Vasculitis: Experience in a Single Centre in Korea. <i>Yonsei Medical Journal</i> , <b>2021</b> , 62, 279-287	3	3
178	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , <b>2021</b> , 53, 1631-1638	2.3	1
177	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , <b>2021</b> , 35, 1761-1768	2.3	0
176	A clinical comparison of an endothelin receptor antagonist and phosphodiesterase type 5 inhibitors for treating digital ulcers of systemic sclerosis. <i>Rheumatology</i> , <b>2021</b> , 60, 5814-5819	3.9	2
175	Fucoxanthin Suppresses Osteoclastogenesis via Modulation of MAP Kinase and Nrf2 Signaling. <i>Marine Drugs</i> , <b>2021</b> , 19,	6	3
174	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , <b>2021</b> , 21, 399-406	4.9	1
173	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , <b>2021</b> , 35, e23731	3	2
172	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , <b>2021</b> , 2021, 6668884	4.3	0

171	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , <b>2021</b> , 16, e0254226	3.7	0
170	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , <b>2021</b> , 36, 1221-1232	2.5	1
169	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , <b>2021</b> , 76, e2501	2.3	
168	Antineutrophil Cytoplasmic Antibody Positivity Is Associated with Vascular Involvement in Behçet's Disease. <i>Yonsei Medical Journal</i> , <b>2021</b> , 62, 149-158	3	1
167	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , <b>2021</b> , 36, e120	4.7	1
166	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , <b>2021</b> , 62, 494-502	3	1
165	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39 Suppl 129, 88-93	2.2	1
164	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39 Suppl 129, 94-100	2.2	
163	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39 Suppl 131, 111-118	2.2	
162	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39, 94-100	2.2	
161	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39, 88-93	2.2	4
160	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , <b>2021</b> , 39, 111-118	2.2	
159	Transcriptional Interactomic Inhibition of ROR $\gamma$ Suppresses Th17-Related Inflammation.. <i>Journal of Inflammation Research</i> , <b>2021</b> , 14, 7091-7105	4.8	0
158	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis.. <i>Journal of Clinical Medicine</i> , <b>2021</b> , 10,	5.1	2
157	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace. <i>Annals of the Rheumatic Diseases</i> , <b>2020</b> ,	2.4	7
156	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , <b>2020</b> , 39, 3009-3015	3.9	2
155	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , <b>2020</b> , 76, 407-412	3	0
154	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , <b>2020</b> , 20, 147	3.5	2

153	Clinical characteristics and long-term outcomes of Libman-Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , <b>2020</b> , 29, 1115-1120	2.6	1
152	Methotrexate-loaded multifunctional nanoparticles with near-infrared irradiation for the treatment of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , <b>2020</b> , 22, 146	5.7	14
151	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , <b>2020</b> , 22, 138	5.7	2
150	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , <b>2020</b> , 21, 258	2.7	2
149	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , <b>2020</b> , 251, 117-123	2.4	3
148	Efficacy and Safety of Rituximab in Korean Patients with Refractory Inflammatory Myopathies. <i>Journal of Korean Medical Science</i> , <b>2020</b> , 35, e335	4.7	1
147	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , <b>2020</b> , 61, 951-957	3	2
146	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , <b>2020</b> , 61, 712-719	3	3
145	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2020</b> , 61, 720-725	3	2
144	Lipid Profiles in Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis: A Cross-sectional Analysis. <i>Journal of Rheumatic Diseases</i> , <b>2020</b> , 27, 261-269	1.2	3
143	Novel PAR2 antagonist ameliorates progression of lupus nephritis in NZB/Z F1 mice. <i>FASEB Journal</i> , <b>2020</b> , 34, 1-1	0.9	
142	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , <b>2020</b> , 61, 970-975	3	1
141	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients: Results From the Korean College of Rheumatology Biologics Registry. <i>Journal of Clinical Rheumatology</i> , <b>2020</b> , Publish Ahead of Print,	1.1	2
140	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , <b>2020</b> , 40, 1063-1070	3.6	3
139	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , <b>2020</b> , 74, e13431	2.9	2
138	Retention Rate and Safety of Biosimilar CT-P13 in Rheumatoid Arthritis: Data from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , <b>2020</b> , 34, 89-98	7.9	9
137	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , <b>2020</b> , 39, 831-840	3.9	1
136	Retention Rate and Efficacy of the Biosimilar CT-P13 Versus Reference Infliximab in Patients with Ankylosing Spondylitis: A Propensity Score-Matched Analysis from the Korean College of Rheumatology Biologics Registry. <i>BioDrugs</i> , <b>2020</b> , 34, 529-539	7.9	4

135	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	6
134	Incidence of Tuberculosis in Systemic Necrotizing Vasculitides: A Population-Based Study From an Intermediate-Burden Country. <i>Frontiers in Medicine</i> , <b>2020</b> , 7, 550004	4.9	
133	Disease-specific factors associated with cardiovascular events in patients with Takayasu arteritis. <i>Arthritis Research and Therapy</i> , <b>2020</b> , 22, 180	5.7	3
132	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , <b>2020</b> , 19, 184	4.4	4
131	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , <b>2020</b> , 2020, 8847306	3.2	0
130	Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , <b>2020</b> , 30, 70-77	3.3	11
129	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , <b>2020</b> , 34, e23022	3	5
128	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , <b>2020</b> , 40, 1089-1099	3.6	4
127	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , <b>2020</b> , 50, 1414-1420	5.3	2
126	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , <b>2020</b> , 22, 73	5.7	4
125	MiR-451 suppresses inflammatory responses in ankylosing spondylitis by targeting macrophage migration inhibitory factor. <i>Clinical and Experimental Rheumatology</i> , <b>2020</b> , 38, 275-281	2.2	4
124	Retention rate and long-term safety of biosimilar CT-P13 in patients with ankylosing spondylitis: data from the Korean College of Rheumatology Biologics registry. <i>Clinical and Experimental Rheumatology</i> , <b>2020</b> , 38, 267-274	2.2	6
123	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , <b>2020</b> , 38 Suppl 124, 236-237	2.2	2
122	Application of the 2019 classification criteria for systemic lupus erythematosus to patients with established ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2020</b> , 38 Suppl 124, 243-244	2.2	
121	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , <b>2019</b> , 39, 1859-1866	3.6	3
120	Classification of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , <b>2019</b> , 26, 156	1.2	6
119	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , <b>2019</b> , 39, 1937-1944	3.6	7
118	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , <b>2019</b> , 19, 90	3	8

117	Long-term efficacy, safety and immunogenicity in patients with rheumatoid arthritis continuing on an etanercept biosimilar (LBEC0101) or switching from reference etanercept to LBEC0101: an open-label extension of a phase III multicentre, randomised, double-blind, parallel-group study. <i>Arthritis Research and Therapy</i> , <b>2019</b> , 21, 122	5.7	5
116	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , <b>2019</b> , 39, 1359-1367	3.6	2
115	Thyroid Dysfunction in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis: A Monocentric Retrospective Study. <i>Journal of Rheumatology</i> , <b>2019</b> , 46, 1248-1250	4.1	0
114	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2019</b> , 60, 454-460	3	6
113	Amino acid signatures of HLA Class-I and II molecules are strongly associated with SLE susceptibility and autoantibody production in Eastern Asians. <i>PLoS Genetics</i> , <b>2019</b> , 15, e1008092	6	21
112	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , <b>2019</b> , 39, 885-892	3.6	1
111	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , <b>2019</b> , 20, e299-e308	4.9	0
110	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , <b>2019</b> , 38, 1685-1690	3.9	4
109	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , <b>2019</b> , 22, 797-804	2.3	8
108	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , <b>2019</b> , 39, 879-884	3.6	3
107	Clinical significance of ANCA positivity in patients with IgA vasculitis: a retrospective monocentric study. <i>Rheumatology International</i> , <b>2019</b> , 39, 1927-1936	3.6	2
106	Atalizumab, a humanized anti-aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , <b>2019</b> , 220, 119408	15.6	7
105	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , <b>2019</b> , 38, 3485-3491	3.9	3
104	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , <b>2019</b> , 14, e0219970	3.7	0
103	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , <b>2019</b> , 39, 1899-1905	3.6	2
102	Causal effect of alcohol consumption on hyperuricemia using a Mendelian randomization design. <i>International Journal of Rheumatic Diseases</i> , <b>2019</b> , 22, 1912-1919	2.3	10
101	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , <b>2019</b> , 2019, 7508240	3	5
100	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , <b>2019</b> , 11,	6.7	5

99	Effects of tapering tumor necrosis factor inhibitor on the achievement of inactive disease in patients with axial spondyloarthritis: a nationwide cohort study. <i>Arthritis Research and Therapy</i> , <b>2019</b> , 21, 163	5.7	7
98	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010-2018. <i>Journal of Clinical Medicine</i> , <b>2019</b> , 8,	5.1	3
97	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2019</b> , 60, 1164-1173 <sup>3</sup>		4
96	Eosinophilic Granulomatosis with Polyangiitis: Experiences in Korean Patients. <i>Yonsei Medical Journal</i> , <b>2019</b> , 60, 705-712	3	7
95	Risk of malignancy in patients with rheumatoid arthritis after anti-tumor necrosis factor therapy: results from Korean National Health Insurance claims data. <i>Korean Journal of Internal Medicine</i> , <b>2019</b> , 34, 669-677	2.5	8
94	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , <b>2019</b> , 39, 395-401	1.5	1
93	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , <b>2019</b> , 38, 487-494	3.9	9
92	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , <b>2019</b> , 23, 223-230	2.5	12
91	Systemic immune-inflammation index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , <b>2019</b> , 24, 711-717	2.2	27
90	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , <b>2019</b> , 38, 1147-1154	3.9	19
89	Antineutrophil Cytoplasmic Antibody-Associated Vasculitis in Korea: A Narrative Review. <i>Yonsei Medical Journal</i> , <b>2019</b> , 60, 10-21	3	8
88	Impact of EUSTAR standardized training on accuracy of modified Rodnan skin score in patients with systemic sclerosis. <i>International Journal of Rheumatic Diseases</i> , <b>2019</b> , 22, 96-102	2.3	4
87	Subclinical but significant liver fibrosis in patients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2019</b> , 37 Suppl 117, 26-31	2.2	1
86	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , <b>2019</b> , 37 Suppl 117, 116-121	2.2	2
85	Intranuclear delivery of the transcription modulation domain of Tbet-improved lupus nephritis in (NZB/NZW) F1 lupus-prone mice. <i>Kidney International</i> , <b>2018</b> , 93, 1118-1130	9.9	6
84	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , <b>2018</b> , 37, 2133-2141	3.9	10
83	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , <b>2018</b> , 38, 1095-1101	3.6	7
82	Increased serum interleukin-32 levels in patients with Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , <b>2018</b> , 21, 2167-2174	2.3	17



81	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , <b>2018</b> , 59, 294-302	3	4
80	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , <b>2018</b> , 19, 187	2.7	21
79	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , <b>2018</b> , 59, 452-456	3	16
78	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2018</b> , 59, 397-405	3	12
77	Extravascular manifestations of Takayasu arteritis: focusing on the features shared with spondyloarthritis. <i>Arthritis Research and Therapy</i> , <b>2018</b> , 20, 142	5.7	11
76	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , <b>2018</b> , 38, 1865-1871	3.6	21
75	C-Reactive Protein to Serum Albumin Ratio Is an Independent Predictor of All-Cause Mortality in Patients with ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , <b>2018</b> , 59, 865-871	3	25
74	Serum leucine-rich $\alpha$ -glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , <b>2018</b> , 486, 253-258	6.2	6
73	Intraperitoneal infusion of mesenchymal stem cell attenuates severity of collagen antibody induced arthritis. <i>PLoS ONE</i> , <b>2018</b> , 13, e0198740	3.7	14
72	Red blood cell distribution width is useful in discriminating adult onset Still's disease and sepsis within 24 hours after hospitalization. <i>Korean Journal of Internal Medicine</i> , <b>2018</b> , 33, 1234-1240	2.5	2
71	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , <b>2018</b> , 37, 1333-1340	3.9	5
70	Rheumatoid factor false positivity in patients with ANCA-associated vasculitis not having medical conditions producing rheumatoid factor. <i>Clinical Rheumatology</i> , <b>2018</b> , 37, 2771-2779	3.9	6
69	The Expression of Immunomodulation-Related Cytokines and Genes of Adipose- and Bone Marrow-Derived Human Mesenchymal Stromal Cells from Early to Late Passages. <i>Tissue Engineering and Regenerative Medicine</i> , <b>2018</b> , 15, 771-779	4.5	12
68	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , <b>2018</b> , 13, e0205651	3.7	8
67	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , <b>2018</b> , 21, 1838-1843	2.3	1
66	HBsAg-negative and anti-HBc-positive in eosinophilic granulomatosis with polyangiitis: a retrospective pilot study. <i>Rheumatology International</i> , <b>2018</b> , 38, 1531-1538	3.6	4
65	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36 Suppl 111, 165-166	2.2	2
64	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36 Suppl 111, 85-87	2.2	12

63	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36 Suppl 111, 115-120	2.2	2
62	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36, 533-539	2.2	2
61	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36 Suppl 111, 73-77	2.2	3
60	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , <b>2018</b> , 36 Suppl 111, 65-72	2.2	8
59	A phase III, multicentre, randomised, double-blind, active-controlled, parallel-group trial comparing safety and efficacy of HD203, with innovator etanercept, in combination with methotrexate, in patients with rheumatoid arthritis: the HERA study. <i>Annals of the Rheumatic Diseases</i> , <b>2017</b> , 76, 65-71	2.4	32
58	Glycated albumin increases with disease activity in rheumatoid factor positive rheumatoid arthritis patients with normal fasting glucose and HbA. <i>Joint Bone Spine</i> , <b>2017</b> , 84, 115-118	2.9	4
57	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , <b>2017</b> , 20, 261-268	2.3	7
56	Five factor score of more than 1 is associated with relapse during the first 2 year-follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , <b>2017</b> , 20, 1261-1268	2.3	9
55	Switching profiles in a population-based cohort of rheumatoid arthritis receiving biologic therapy: results from the KOBIO registry. <i>Clinical Rheumatology</i> , <b>2017</b> , 36, 1013-1022	3.9	16
54	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , <b>2017</b> , 47, 216-221	5.3	21
53	Evaluation of Spleen Glucose Metabolism Using F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , <b>2017</b> , 58, 507-513	8.9	24
52	Monthly Oral Ibandronate Reduces Bone Loss in Korean Women With Rheumatoid Arthritis and Osteopenia Receiving Long-term Glucocorticoids: A 48-week Double-blinded Randomized Placebo-controlled Investigator-initiated Trial. <i>Clinical Therapeutics</i> , <b>2017</b> , 39, 268-278.e2	3.5	21
51	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , <b>2017</b> , 44, 996-1003	4.1	31
50	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , <b>2017</b> , 56, 795-801	3.9	6
49	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , <b>2017</b> , 20, 1028-1035	2.3	24
48	Control of dental-derived induced pluripotent stem cells through modified surfaces for dental application. <i>Acta Odontologica Scandinavica</i> , <b>2017</b> , 75, 309-318	2.2	2
47	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , <b>2017</b> , 37, 685-694	3.6	11
46	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , <b>2017</b> , 36, 2751-2759	3.9	8

45	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , <b>2017</b> , 12, e0184337	3
44	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , <b>2017</b> , 19, 193	5-7 9
43	Autologous Haematopoietic Stem Cell Transplantation for Refractory Rheumatic Diseases. <i>Journal of Rheumatic Diseases</i> , <b>2017</b> , 24, 171	1.2
42	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , <b>2017</b> , 58, 1128-1134	3 5
41	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , <b>2017</b> , 20, 1593-1605	2.3 12
40	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , <b>2017</b> , 84, 583-587	2.9 8
39	Survival and prognostic factors in patients with connective tissue disease-associated pulmonary hypertension diagnosed by echocardiography: results from a Korean nationwide registry. <i>International Journal of Rheumatic Diseases</i> , <b>2017</b> , 20, 1227-1236	2.3 15
38	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , <b>2017</b> , 12, e0181590	3-7 7
37	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , <b>2017</b> , 35 Suppl 103, 47-54	2.2 10
36	Differential expressions of NOD-like receptors and their associations with inflammatory responses in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , <b>2017</b> , 35, 630-637	2.2 17
35	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , <b>2017</b> , 35 Suppl 103, 111-118	2.2 13
34	Real-time monitoring of 3D cell culture using a 3D capacitance biosensor. <i>Biosensors and Bioelectronics</i> , <b>2016</b> , 77, 56-61	11.8 39
33	High-density genotyping of immune-related loci identifies new SLE risk variants in individuals with Asian ancestry. <i>Nature Genetics</i> , <b>2016</b> , 48, 323-30	36.3 156
32	Clinical features of psoriatic arthritis in Korean patients with psoriasis: a cross-sectional observational study of 196 patients with psoriasis using psoriatic arthritis screening questionnaires. <i>Rheumatology International</i> , <b>2016</b> , 36, 207-12	3.6 19
31	Brain meningioma in a patient with systemic lupus erythematosus. <i>Yeungnam University Journal of Medicine</i> , <b>2016</b> , 33, 159	1.3 1
30	Gastrointestinal Risk Factors and Non-steroidal Anti-inflammatory Drugs Use in Rheumatoid Arthritis and Osteoarthritis Patients in Korea. <i>Journal of Rheumatic Diseases</i> , <b>2016</b> , 23, 47	1.2 1
29	Anti-Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , <b>2016</b> , 19, 897-902	2.3 16
28	Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , <b>2016</b> , 26, 576-82	3.3 2

27	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , <b>2015</b> , 17, 77	5.7	15
26	Drug-loaded gold/iron/gold plasmonic nanoparticles for magnetic targeted chemo-photothermal treatment of rheumatoid arthritis. <i>Biomaterials</i> , <b>2015</b> , 61, 95-102	15.6	97
25	A Case of Aortitis with Systemic Lupus Erythematosus. <i>Journal of Rheumatic Diseases</i> , <b>2015</b> , 22, 205	1.2	1
24	Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , <b>2015</b> , 33, S68-74	2.2	4
23	Plasma chemerin levels in rheumatoid arthritis are correlated with disease activity rather than obesity. <i>Joint Bone Spine</i> , <b>2014</b> , 81, 189-90	2.9	25
22	Secreted Frizzled-related protein 5 suppresses inflammatory response in rheumatoid arthritis fibroblast-like synoviocytes through down-regulation of c-Jun N-terminal kinase. <i>Rheumatology</i> , <b>2014</b> , 53, 1704-11	3.9	11
21	Delta neutrophil index as an early marker for differential diagnosis of adult-onset Still's disease and sepsis. <i>Yonsei Medical Journal</i> , <b>2014</b> , 55, 753-9	3	15
20	Differences in clinical manifestations and outcomes between adult and child patients with Henoch-Schönlein purpura. <i>Journal of Korean Medical Science</i> , <b>2014</b> , 29, 198-203	4.7	76
19	Usefulness of serum leucine-rich alpha-2 glycoprotein as a disease activity biomarker in patients with rheumatoid arthritis. <i>Journal of Korean Medical Science</i> , <b>2014</b> , 29, 1199-204	4.7	28
18	Inhibition of glycogen synthase kinase-3 $\beta$ suppresses inflammatory responses in rheumatoid arthritis fibroblast-like synoviocytes and collagen-induced arthritis. <i>Joint Bone Spine</i> , <b>2014</b> , 81, 240-6	2.9	21
17	Soluble CD93 levels in patients with acute myocardial infarction and its implication on clinical outcome. <i>PLoS ONE</i> , <b>2014</b> , 9, e96538	3.7	12
16	Treating rheumatoid arthritis to target: recommendations assessment questionnaire in Korea. <i>Clinical Rheumatology</i> , <b>2013</b> , 32, 1791-7	3.9	3
15	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , <b>2013</b> , 23, 509-515	3.3	3
14	Targeted chemo-photothermal treatments of rheumatoid arthritis using gold half-shell multifunctional nanoparticles. <i>ACS Nano</i> , <b>2013</b> , 7, 50-7	16.7	181
13	Typical 18-FDG-PET/CT Findings of Polymyalgia Rheumatica: A Case Report. <i>Journal of Rheumatic Diseases</i> , <b>2013</b> , 20, 113	1.2	4
12	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , <b>2013</b> , 23, 509-15	3.3	1
11	The role of (18) F-fluorodeoxyglucose-positron emission tomography in the assessment of disease activity in patients with takayasu arteritis. <i>Arthritis and Rheumatism</i> , <b>2012</b> , 64, 866-75		74
10	Tuberculin reaction is not attenuated in patients with rheumatoid arthritis living in a region with intermediate burden of tuberculosis. <i>Rheumatology International</i> , <b>2012</b> , 32, 1421-4	3.6	5

9	Multicenter retrospective analysis of the effectiveness and safety of rituximab in Korean patients with refractory systemic lupus erythematosus. <i>Autoimmune Diseases</i> , <b>2012</b> , 2012, 565039	2.9	15
8	The Effect of Hypouricemic Treatment on the Renal Function in Patients with Gout. <i>Journal of Rheumatic Diseases</i> , <b>2011</b> , 18, 26	1.2	1
7	A Case of Ankylosing Spondylitis with Follicular Lymphoma. <i>Journal of Rheumatic Diseases</i> , <b>2011</b> , 18, 315	1.2	
6	ANCA-associated Vasculitis after Scrub Typhus. <i>Infection and Chemotherapy</i> , <b>2011</b> , 43, 82	3.9	1
5	Clinical Manifestations of Korean Adult Patients with Henoch-Schönlein Purpura. <i>The Journal of the Korean Rheumatism Association</i> , <b>2010</b> , 17, 133		3
4	The mid-range of the adjusted level of ferritin can predict the chronic course in patients with adult onset Still's disease. <i>Journal of Rheumatology</i> , <b>2009</b> , 36, 156-62	4.1	23
3	Atherosclerosis in rheumatoid arthritis: morphologic evidence obtained by carotid ultrasound. <i>Arthritis and Rheumatism</i> , <b>2002</b> , 46, 1714-9		191
2	Effects of antirheumatic therapy on serum lipid levels in patients with rheumatoid arthritis: a prospective study. <i>American Journal of Medicine</i> , <b>2002</b> , 113, 188-93	2.4	141
1	GeTe Nanosheets as Theranostic Agents for Multimodal Imaging and Therapy of Inflammatory Bowel Disease. <i>Advanced Functional Materials</i> , 2107433	15.6	1