

# Masahito Yoshinari

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9433952/publications.pdf>

Version: 2024-02-01

21  
papers

223  
citations

1040056

9  
h-index

1058476

14  
g-index

21  
all docs

21  
docs citations

21  
times ranked

388  
citing authors

#	ARTICLE	IF	CITATIONS
1	Incidence of diabetic foot ulcer in Japanese patients with type 2 diabetes mellitus: The Fukuoka diabetes registry. <i>Diabetes Research and Clinical Practice</i> , 2018, 137, 183-189.	2.8	39
2	The serum creatinine to cystatin C ratio predicts bone fracture in patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2018, 146, 202-210.	2.8	23
3	Impact of age at menarche on obesity and glycemic control in Japanese patients with type 2 diabetes: Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1216-1223.	2.4	20
4	Impact of Body Weight Loss From Maximum Weight on Fragility Bone Fractures in Japanese Patients With Type 2 Diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Care</i> , 2018, 41, 1061-1067.	8.6	19
5	Additive effects of green tea and coffee on all-cause mortality in patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001252.	2.8	19
6	Serum adiponectin predicts fracture risk in individuals with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Diabetologia</i> , 2017, 60, 1922-1930.	6.3	13
7	Association Between Genetic Risk and Development of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 3213-3222.	3.6	12
8	Comparison of the contributions of impaired beta cell function and insulin resistance to the development of type 2 diabetes in a Japanese community: the Hisayama Study. <i>Diabetologia</i> , 2021, 64, 1775-1784.	6.3	10
9	Polypharmacy and bone fracture risk in patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2021, 181, 109097.	2.8	10
10	Serum Ethylamine Levels as an Indicator of L-Theanine Consumption and the Risk of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. <i>Diabetes Care</i> , 2019, 42, 1234-1240.	8.6	9
11	Incidence of stroke and its association with glycemic control and lifestyle in Japanese patients with type 2 diabetes mellitus: The Fukuoka diabetes registry. <i>Diabetes Research and Clinical Practice</i> , 2021, 172, 108518.	2.8	7
12	Incidence of end-stage renal disease and risk factors for progression of renal dysfunction in Japanese patients with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Clinical and Experimental Nephrology</i> , 2021, , 1.	1.6	7
13	The gene-treatment interaction of paraoxonase-1 gene polymorphism and statin therapy on insulin secretion in Japanese patients with type 2 diabetes: Fukuoka diabetes registry. <i>BMC Medical Genetics</i> , 2017, 18, 146.	2.1	6
14	30-minute postload plasma glucose levels during an oral glucose tolerance test predict the risk of future type 2 diabetes: the Hisayama Study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001156.	2.8	5
15	Impact of hip fracture on all-cause mortality in Japanese patients with type 2 diabetes mellitus: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2020, 11, 62-69.	2.4	4
16	Incidence of severe hypoglycemia and its association with serum adiponectin in Japanese patients with type 1 and insulin-treated type 2 diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1258-1264.	2.4	4
17	Constipation and diabetic kidney disease: The Fukuoka Diabetes Registry. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 1247-1254.	1.6	4
18	Relationship of coffee consumption with a decline in kidney function among patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2022, 13, 1030-1038.	2.4	4

#	ARTICLE	IF	CITATIONS
19	Defecation frequency and glycemic control in patients with diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes and Its Complications</i> , 2021, 35, 107751.	2.3	3
20	Usefulness of urinary tubule injury markers for predicting progression of renal dysfunction in patients with type 2 diabetes and albuminuria: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109840.	2.8	3
21	Prospective study of cancer in Japanese patients with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Diabetology International</i> , 2019, 10, 260-267.	1.4	2