

Masahito Yoshinari

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9433952/publications.pdf>

Version: 2024-02-01

21
papers

223
citations

1040056

9
h-index

1058476

14
g-index

21
all docs

21
docs citations

21
times ranked

388
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Relationship of coffee consumption with a decline in kidney function among patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2022, 13, 1030-1038. | 2.4 | 4 |
| 2 | Usefulness of urinary tubule injury markers for predicting progression of renal dysfunction in patients with type 2 diabetes and albuminuria: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109840. | 2.8 | 3 |
| 3 | Defecation frequency and glycemic control in patients with diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes and Its Complications</i> , 2021, 35, 107751. | 2.3 | 3 |
| 4 | Incidence of stroke and its association with glycemic control and lifestyle in Japanese patients with type 2 diabetes mellitus: The Fukuoka diabetes registry. <i>Diabetes Research and Clinical Practice</i> , 2021, 172, 108518. | 2.8 | 7 |
| 5 | Comparison of the contributions of impaired beta cell function and insulin resistance to the development of type 2 diabetes in a Japanese community: the Hisayama Study. <i>Diabetologia</i> , 2021, 64, 1775-1784. | 6.3 | 10 |
| 6 | Constipation and diabetic kidney disease: The Fukuoka Diabetes Registry. <i>Clinical and Experimental Nephrology</i> , 2021, 25, 1247-1254. | 1.6 | 4 |
| 7 | Incidence of end-stage renal disease and risk factors for progression of renal dysfunction in Japanese patients with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Clinical and Experimental Nephrology</i> , 2021, , 1. | 1.6 | 7 |
| 8 | Polypharmacy and bone fracture risk in patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2021, 181, 109097. | 2.8 | 10 |
| 9 | Impact of hip fracture on all-cause mortality in Japanese patients with type 2 diabetes mellitus: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2020, 11, 62-69. | 2.4 | 4 |
| 10 | Additive effects of green tea and coffee on all-cause mortality in patients with type 2 diabetes mellitus: the Fukuoka Diabetes Registry. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001252. | 2.8 | 19 |
| 11 | 30-minute postload plasma glucose levels during an oral glucose tolerance test predict the risk of future type 2 diabetes: the Hisayama Study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001156. | 2.8 | 5 |
| 12 | Incidence of severe hypoglycemia and its association with serum adiponectin in Japanese patients with type 1 and insulin-treated type 2 diabetes: The Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1258-1264. | 2.4 | 4 |
| 13 | Association Between Genetic Risk and Development of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 3213-3222. | 3.6 | 12 |
| 14 | Serum Ethylamine Levels as an Indicator of Theanine Consumption and the Risk of Type 2 Diabetes in a General Japanese Population: The Hisayama Study. <i>Diabetes Care</i> , 2019, 42, 1234-1240. | 8.6 | 9 |
| 15 | Prospective study of cancer in Japanese patients with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Diabetology International</i> , 2019, 10, 260-267. | 1.4 | 2 |
| 16 | Incidence of diabetic foot ulcer in Japanese patients with type 2 diabetes mellitus: The Fukuoka diabetes registry. <i>Diabetes Research and Clinical Practice</i> , 2018, 137, 183-189. | 2.8 | 39 |
| 17 | Impact of age at menarche on obesity and glycemic control in Japanese patients with type 2 diabetes: Fukuoka Diabetes Registry. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1216-1223. | 2.4 | 20 |
| 18 | Impact of Body Weight Loss From Maximum Weight on Fragility Bone Fractures in Japanese Patients With Type 2 Diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Care</i> , 2018, 41, 1061-1067. | 8.6 | 19 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The serum creatinine to cystatin C ratio predicts bone fracture in patients with type 2 diabetes: The Fukuoka Diabetes Registry. <i>Diabetes Research and Clinical Practice</i> , 2018, 146, 202-210. | 2.8 | 23 |
| 20 | Serum adiponectin predicts fracture risk in individuals with type 2 diabetes: the Fukuoka Diabetes Registry. <i>Diabetologia</i> , 2017, 60, 1922-1930. | 6.3 | 13 |
| 21 | The gene-treatment interaction of paraoxonase-1 gene polymorphism and statin therapy on insulin secretion in Japanese patients with type 2 diabetes: Fukuoka diabetes registry. <i>BMC Medical Genetics</i> , 2017, 18, 146. | 2.1 | 6 |