Thomas F Parkerton

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/9440743/thomas-f-parkerton-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

62 89 4,097 31 h-index g-index citations papers 4,595 5.22 91 5.3 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
89	Assessing the Toxicity of Individual Aromatic Compounds and Mixtures to American Lobster (Homarus americanus) Larvae Using a Passive Dosing System. <i>Environmental Toxicology and Chemistry</i> , 2021 , 40, 1379-1388	3.8	O
88	Miniaturised marine tests as indicators of aromatic hydrocarbon toxicity: Potential applicability to oil spill assessment. <i>Marine Pollution Bulletin</i> , 2021 , 165, 112151	6.7	3
87	Assessing toxicity of hydrophobic aliphatic and monoaromatic hydrocarbons at the solubility limit using novel dosing methods. <i>Chemosphere</i> , 2021 , 265, 129174	8.4	1
86	Toxicity of two representative petroleum hydrocarbons, toluene and phenanthrene, to five Atlantic coral species. <i>Marine Pollution Bulletin</i> , 2021 , 169, 112560	6.7	3
85	Derivation of toxicity thresholds for gas condensate oils protective of tropical species using experimental and modelling approaches. <i>Marine Pollution Bulletin</i> , 2021 , 172, 112899	6.7	O
84	The sensitivity of the deepsea species northern shrimp (Pandalus borealis) and the cold-water coral (Lophelia pertusa) to oil-associated aromatic compounds, dispersant, and Alaskan North Slope crude oil. <i>Marine Pollution Bulletin</i> , 2020 , 156, 111202	6.7	6
83	Recommendations for Improving Methods and Models for Aquatic Hazard Assessment of Ionizable Organic Chemicals. <i>Environmental Toxicology and Chemistry</i> , 2020 , 39, 269-286	3.8	24
82	Comparison of In Situ and Ex Situ Equilibrium Passive Sampling for Measuring Freely Dissolved Concentrations of Parent and Alkylated Polycyclic Aromatic Hydrocarbons in Sediments. <i>Environmental Toxicology and Chemistry</i> , 2020 , 39, 2169-2179	3.8	4
81	A Toxicokinetic Framework and Analysis Tool for Interpreting Organisation for Economic Co-operation and Development Guideline 305 Dietary Bioaccumulation Tests. <i>Environmental Toxicology and Chemistry</i> , 2020 , 39, 171-188	3.8	8
80	Modeling the toxicity of dissolved crude oil exposures to characterize the sensitivity of cod (Gadus morhua) larvae and role of individual and unresolved hydrocarbons. <i>Marine Pollution Bulletin</i> , 2019 , 138, 286-294	6.7	10
79	The treatment of biodegradation in models of sub-surface oil spills: A review and sensitivity study. <i>Marine Pollution Bulletin</i> , 2019 , 143, 204-219	6.7	25
78	Alternative Management of Oil and Gas Produced Water Requires More Research on Its Hazards and Risks. <i>Integrated Environmental Assessment and Management</i> , 2019 , 15, 677-682	2.5	11
77	Bioconcentration factors for hydrocarbons and petrochemicals: Understanding processes, uncertainty and predictive model performance. <i>Chemosphere</i> , 2019 , 226, 472-482	8.4	4
76	Review of Polycyclic Aromatic Hydrocarbons (PAHs) Sediment Quality Guidelines for the Protection of Benthic Life. <i>Integrated Environmental Assessment and Management</i> , 2019 , 15, 505-518	2.5	28
75	Re-evaluation of target lipid model-derived HC5 predictions for hydrocarbons. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 1579-1593	3.8	31
74	Technical basis for using passive sampling as a biomimetic extraction procedure to assess bioavailability and predict toxicity of petroleum substances. <i>Chemosphere</i> , 2018 , 199, 585-594	8.4	17
73	Aquatic exposures of chemical mixtures in urban environments: Approaches to impact assessment. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 703-714	3.8	11

(2014-2018)

72	of a regular series intended to sharpen understanding of current and emerging topics of interest to the scientific community. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 1235-1251	3.8	25
71	The sensitivity of a deep-sea fish species (Anoplopoma fimbria) to oil-associated aromatic compounds, dispersant, and Alaskan North Slope crude oil. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 2210-2221	3.8	22
70	Application of the Target Lipid Model and Passive Samplers to Characterize the Toxicity of Bioavailable Organics in Oil Sands Process-Affected Water. <i>Environmental Science & Environmental Science &</i>	10.3	18
69	Comparative Risk Assessment of spill response options for a deepwater oil well blowout: Part 1. Oil spill modeling. <i>Marine Pollution Bulletin</i> , 2018 , 133, 1001-1015	6.7	40
68	An evaluation of cumulative risks from offshore produced water discharges in the Bass Strait. <i>Marine Pollution Bulletin</i> , 2018 , 126, 610-621	6.7	9
67	Passive dosing yields dissolved aqueous exposures of crude oil comparable to the CROSERF (Chemical Response to Oil Spill: Ecological Effects Research Forum) water accommodated fraction method. <i>Environmental Toxicology and Chemistry</i> , 2018 , 37, 2810-2819	3.8	18
66	Response to Comment on "Assessing Aromatic-Hydrocarbon Toxicity to Fish Early Life Stages Using Passive-Dosing Methods and Target-Lipid and Chemical-Activity Models". <i>Environmental Science & Environmental Science</i>	10.3	
65	A re-evaluation of PETROTOX for predicting acute and chronic toxicity of petroleum substances. <i>Environmental Toxicology and Chemistry</i> , 2017 , 36, 2245-2252	3.8	23
64	Investigating the role of dissolved and droplet oil in aquatic toxicity using dispersed and passive dosing systems. <i>Environmental Toxicology and Chemistry</i> , 2017 , 36, 1020-1028	3.8	20
63	In Vivo Biotransformation Rates of Organic Chemicals in Fish: Relationship with Bioconcentration and Biomagnification Factors. <i>Environmental Science & Environmental Science </i>	10.3	24
62	Chronic toxicity of selected polycyclic aromatic hydrocarbons to algae and crustaceans using passive dosing. <i>Environmental Toxicology and Chemistry</i> , 2016 , 35, 2948-2957	3.8	24
61	Oil dispersants do facilitate biodegradation of spilled oil. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016 , 113, E1421	11.5	26
60	Assessing Aromatic-Hydrocarbon Toxicity to Fish Early Life Stages Using Passive-Dosing Methods and Target-Lipid and Chemical-Activity Models. <i>Environmental Science & Environmental Science & Environ</i>	193	23
59	Preparing the Hydrocarbon/Crude Oil. <i>Springer Protocols</i> , 2016 , 15-32	0.3	3
58	Water solubility of selected C9-C18 alkanes using a slow-stir technique: Comparison to structure - property models. <i>Chemosphere</i> , 2016 , 150, 416-423	8.4	14
57	Guidance for improving comparability and relevance of oil toxicity tests. <i>Marine Pollution Bulletin</i> , 2015 , 98, 156-70	6.7	66
56	Assessing the Fate of an Aromatic Hydrocarbon Fluid in Agricultural Spray Applications Using the Three-Stage ADVOCATE Model Framework. <i>Journal of Agricultural and Food Chemistry</i> , 2015 , 63, 6866-7	· 5 ·7	2
55	PETRORISK: a risk assessment framework for petroleum substances. <i>Integrated Environmental Assessment and Management</i> , 2014 , 10, 437-48	2.5	26

54	Passive sampling methods for contaminated sediments: risk assessment and management. <i>Integrated Environmental Assessment and Management</i> , 2014 , 10, 224-36	2.5	40
53	Evaluating toxicity of heavy fuel oil fractions using complementary modeling and biomimetic extraction methods. <i>Environmental Toxicology and Chemistry</i> , 2014 , 33, 2094-104	3.8	26
52	Use of passive samplers for improving oil toxicity and spill effects assessment. <i>Marine Pollution Bulletin</i> , 2014 , 86, 274-282	6.7	26
51	Comment on "Toxicity and mutagenicity of Gulf of Mexico waters during and after the deepwater horizon oil spill". <i>Environmental Science & Environmental Science & Environment</i>	10.3	6
50	Passive sampling in contaminated sediment assessment: building consensus to improve decision making. <i>Integrated Environmental Assessment and Management</i> , 2014 , 10, 163-6	2.5	15
49	Passive sampling methods for contaminated sediments: scientific rationale supporting use of freely dissolved concentrations. <i>Integrated Environmental Assessment and Management</i> , 2014 , 10, 197-209	2.5	122
48	Extension and validation of the target lipid model for deriving predicted no-effect concentrations for soils and sediments. <i>Environmental Toxicology and Chemistry</i> , 2014 , 33, 2679-87	3.8	15
47	Improving the quality and scientific understanding of trophic magnification factors (TMFs). <i>Environmental Science & amp; Technology</i> , 2013 , 47, 1186-7	10.3	40
46	A novel passive dosing system for determining the toxicity of phenanthrene to early life stages of zebrafish. <i>Science of the Total Environment</i> , 2013 , 463-464, 952-8	10.2	40
45	Modeling bioaccumulation in coupled pelagic-benthic food chains: past insights and future directions. <i>Environmental Toxicology and Chemistry</i> , 2013 , 32, 1931-4	3.8	1
44	Comparing laboratory- and field-measured biota-sediment accumulation factors. <i>Integrated Environmental Assessment and Management</i> , 2012 , 8, 32-41	2.5	24
43	Comparing laboratory and field measured bioaccumulation endpoints. <i>Integrated Environmental Assessment and Management</i> , 2012 , 8, 17-31	2.5	54
42	PETROTOX: an aquatic toxicity model for petroleum substances. <i>Environmental Toxicology and Chemistry</i> , 2012 , 31, 2498-506	3.8	75
41	A review of the tissue residue approach for organic and organometallic compounds in aquatic organisms. <i>Integrated Environmental Assessment and Management</i> , 2011 , 7, 50-74	2.5	44
40	Temporal patterns in the transcriptomic response of rainbow trout, Oncorhynchus mykiss, to crude oil. <i>Aquatic Toxicology</i> , 2010 , 99, 320-9	5.1	31
39	Multimedia modeling of human exposure to chemical substances: the roles of food web biomagnification and biotransformation. <i>Environmental Toxicology and Chemistry</i> , 2010 , 29, 45-55	3.8	35
38	Hepatic gene expression in rainbow trout (Oncorhynchus mykiss) exposed to different hydrocarbon mixtures. <i>Environmental Toxicology and Chemistry</i> , 2010 , 29, 2034-43	3.8	15
37	Environmental fate factors and human intake fractions for risk assessment of petroleum products. <i>Integrated Environmental Assessment and Management</i> , 2010 , 6, 135-44	2.5	6

(2003-2009)

36	Evaluation of bioaccumulation using in vivo laboratory and field studies. <i>Integrated Environmental Assessment and Management</i> , 2009 , 5, 598-623	2.5	68
35	Aqueous solubility and Daphnia magna chronic toxicity of di(2-ethylhexyl) adipate. <i>Bulletin of Environmental Contamination and Toxicology</i> , 2008 , 80, 539-43	2.7	3
34	A database of fish biotransformation rates for organic chemicals. <i>Environmental Toxicology and Chemistry</i> , 2008 , 27, 2263-70	3.8	105
33	Guidance for evaluating in vivo fish bioaccumulation data. <i>Integrated Environmental Assessment and Management</i> , 2008 , 4, 139-55	2.5	38
32	Development of a Multimedia Model for the Fate Prediction of Hydrocarbon Fluids in Agrochemical Formulations. <i>Journal of ASTM International</i> , 2008 , 5, 101637		1
31	The primary aerobic biodegradation of gasoline hydrocarbons. <i>Environmental Science & Environmental Sc</i>	10.3	63
30	Application of the target lipid model for deriving predicted no-effect concentrations for wastewater organisms. <i>Environmental Toxicology and Chemistry</i> , 2007 , 26, 2317-31	3.8	16
29	Assessing the aquatic hazard of commercial hydrocarbon resins. <i>Ecotoxicology and Environmental Safety</i> , 2007 , 66, 159-68	7	12
28	A kinetic model for predicting biodegradation. SAR and QSAR in Environmental Research, 2007, 18, 443-	53 .5	30
27	Hazard evaluation of diisononyl phthalate and diisodecyl phthalate in a Japanese medaka multigenerational assay. <i>Ecotoxicology and Environmental Safety</i> , 2006 , 65, 36-47	7	22
26	Base-line model for identifying the bioaccumulation potential of chemicals. <i>SAR and QSAR in Environmental Research</i> , 2005 , 16, 531-54	3.5	86
25	Five-stage environmental exposure assessment strategy for mixtures: gasoline as a case study. <i>Environmental Science & Environmental Environ</i>	10.3	23
24	Validation of the narcosis target lipid model for petroleum products: gasoline as a case study. <i>Environmental Toxicology and Chemistry</i> , 2005 , 24, 2382-94	3.8	64
23	A new biodegradation prediction model specific to petroleum hydrocarbons. <i>Environmental Toxicology and Chemistry</i> , 2005 , 24, 1847-60	3.8	54
22	Application of the narcosis target lipid model to algal toxicity and deriving predicted-no-effect concentrations. <i>Environmental Toxicology and Chemistry</i> , 2004 , 23, 2503-17	3.8	54
21	Applications of contaminant fate and bioaccumulation models in assessing ecological risks of chemicals: a case study for gasoline hydrocarbons. <i>Environmental Science & Environmental Science & Envir</i>	10.3	33
20	An Assessment of the Potential Environmental Risks Posed by Phthalates in Soil and Sediment. Handbook of Environmental Chemistry, 2003 , 317-349	0.8	4
19	Bioaccumulation of Phthalate Esters in Aquatic Food-Webs. <i>Handbook of Environmental Chemistry</i> , 2003 , 201-225	0.8	15

18	Physical-Chemical Properties and Evaluative Fate Modelling of Phthalate Esters. <i>Handbook of Environmental Chemistry</i> , 2003 , 57-84	0.8	23
17	Slow-stir water solubility measurements of selected alcohols and diesters. <i>Chemosphere</i> , 2002 , 48, 257-	65 .4	19
16	An assessment of the toxicity of phthalate esters to freshwater benthos. 1. Aqueous exposures. <i>Environmental Toxicology and Chemistry</i> , 2001 , 20, 1798-1804	3.8	39
15	An assessment of the toxicity of phthalate esters to freshwater benthos. 2. Sediment exposures. <i>Environmental Toxicology and Chemistry</i> , 2001 , 20, 1805-1815	3.8	22
14	Application of quantitative structureactivity relationships for assessing the aquatic toxicity of phthalate esters. <i>Ecotoxicology and Environmental Safety</i> , 2000 , 45, 61-78	7	56
13	A risk assessment of selected phthalate esters in North American and Western European surface waters. <i>Chemosphere</i> , 2000 , 40, 885-91	8.4	68
12	Polyacrylate-Coated SPME Fibers as a Tool To Simulate Body Residues and Target Concentrations of Complex Organic Mixtures for Estimation of Baseline Toxicity. <i>Environmental Science & Eamp; Technology</i> , 2000 , 34, 324-331	10.3	68
11	Assessing the aquatic toxicity of complex hydrocarbon mixtures using solid phase microextraction. <i>Toxicology Letters</i> , 2000 , 112-113, 273-82	4.4	50
10	A proposed multigeneration protocol for Japanese medaka (Oryzias latipes) to evaluate effects of endocrine disruptors. <i>Science of the Total Environment</i> , 1999 , 233, 211-20	10.2	75
9	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749	8.4	1123
9			1123 151
	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749		
8	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749 Aquatic toxicity of eighteen phthalate esters. <i>Environmental Toxicology and Chemistry</i> , 1997 , 16, 875-89 Sediment sorption coefficient measurements for four phthalate esters: Experimental results and	13.8	151
8	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749 Aquatic toxicity of eighteen phthalate esters. <i>Environmental Toxicology and Chemistry</i> , 1997 , 16, 875-89 Sediment sorption coefficient measurements for four phthalate esters: Experimental results and model theory. <i>Environmental Toxicology and Chemistry</i> , 1995 , 14, 1477-1486 Do aquatic effects or human health end points govern the development of sediment-quality	1 _{3.8}	151 25
8 7 6	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749 Aquatic toxicity of eighteen phthalate esters. <i>Environmental Toxicology and Chemistry</i> , 1997 , 16, 875-89 Sediment sorption coefficient measurements for four phthalate esters: Experimental results and model theory. <i>Environmental Toxicology and Chemistry</i> , 1995 , 14, 1477-1486 Do aquatic effects or human health end points govern the development of sediment-quality criteria for nonionic organic chemicals?. <i>Environmental Toxicology and Chemistry</i> , 1993 , 12, 507-523 An equilibrium model of organic chemical accumulation in aquatic food webs with sediment	13.8 3.8 3.8	151 25 22
8 7 6	The environmental fate of phthalate esters: A literature review. <i>Chemosphere</i> , 1997 , 35, 667-749 Aquatic toxicity of eighteen phthalate esters. <i>Environmental Toxicology and Chemistry</i> , 1997 , 16, 875-89 Sediment sorption coefficient measurements for four phthalate esters: Experimental results and model theory. <i>Environmental Toxicology and Chemistry</i> , 1995 , 14, 1477-1486 Do aquatic effects or human health end points govern the development of sediment-quality criteria for nonionic organic chemicals?. <i>Environmental Toxicology and Chemistry</i> , 1993 , 12, 507-523 An equilibrium model of organic chemical accumulation in aquatic food webs with sediment interaction. <i>Environmental Toxicology and Chemistry</i> , 1992 , 11, 615-629	13.8 3.8 3.8	151 25 22 276
8 7 6 5	The environmental fate of phthalate esters: A literature review. Chemosphere, 1997, 35, 667-749 Aquatic toxicity of eighteen phthalate esters. Environmental Toxicology and Chemistry, 1997, 16, 875-89 Sediment sorption coefficient measurements for four phthalate esters: Experimental results and model theory. Environmental Toxicology and Chemistry, 1995, 14, 1477-1486 Do aquatic effects or human health end points govern the development of sediment-quality criteria for nonionic organic chemicals?. Environmental Toxicology and Chemistry, 1993, 12, 507-523 An equilibrium model of organic chemical accumulation in aquatic food webs with sediment interaction. Environmental Toxicology and Chemistry, 1992, 11, 615-629 An equilibrium model of organic chemical accumulation in aquatic food webs with sediment interaction 1992, 11, 615	13.8 3.8 3.8 3.8	151 25 22 276 8