

Archana Sharma

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9452919/publications.pdf>

Version: 2024-02-01

25
papers

1,053
citations

471061

17
h-index

580395

25
g-index

25
all docs

25
docs citations

25
times ranked

1791
citing authors

#	ARTICLE	IF	CITATIONS
1	T cell factor 1 initiates the T helper type 2 fate by inducing the transcription factor GATA-3 and repressing interferon- γ . <i>Nature Immunology</i> , 2009, 10, 992-999.	7.0	179
2	The AGC kinase SGK1 regulates TH1 and TH2 differentiation downstream of the mTORC2 complex. <i>Nature Immunology</i> , 2014, 15, 457-464.	7.0	163
3	Cold-inducible RNA-binding protein causes endothelial dysfunction via activation of Nlrp3 inflammasome. <i>Scientific Reports</i> , 2016, 6, 26571.	1.6	81
4	TCF1 and β -catenin regulate T cell development and function. <i>Immunologic Research</i> , 2010, 47, 45-55.	1.3	74
5	Blocking Cold-Inducible RNA-Binding Protein Protects Liver From Ischemia-Reperfusion Injury. <i>Shock</i> , 2015, 43, 24-30.	1.0	72
6	T Cell Factor-1 Negatively Regulates Expression of IL-17 Family of Cytokines and Protects Mice from Experimental Autoimmune Encephalomyelitis. <i>Journal of Immunology</i> , 2011, 186, 3946-3952.	0.4	60
7	Differential alterations of tissue T-cell subsets after sepsis. <i>Immunology Letters</i> , 2015, 168, 41-50.	1.1	41
8	Mitigation of sepsis-induced inflammatory responses and organ injury through targeting Wnt/ β -catenin signaling. <i>Scientific Reports</i> , 2017, 7, 9235.	1.6	41
9	Cold-inducible RNA-binding protein activates splenic T cells during sepsis in a TLR4-dependent manner. <i>Cellular and Molecular Immunology</i> , 2018, 15, 38-47.	4.8	41
10	Receptor-interacting protein kinase 3 deficiency inhibits immune cell infiltration and attenuates organ injury in sepsis. <i>Critical Care</i> , 2014, 18, R142.	2.5	40
11	T Cell Factor-1 and β -Catenin Control the Development of Memory-like CD8 Thymocytes. <i>Journal of Immunology</i> , 2012, 188, 3859-3868.	0.4	27
12	Upregulation of GRAIL Is Associated with Impaired CD4 T Cell Proliferation in Sepsis. <i>Journal of Immunology</i> , 2014, 192, 2305-2314.	0.4	27
13	Pre-TCR-Induced β -Catenin Facilitates Traversal through β -Selection. <i>Journal of Immunology</i> , 2009, 182, 751-758.	0.4	26
14	Milk fat globule epidermal growth factor-factor 8-derived peptide attenuates organ injury and improves survival in sepsis. <i>Critical Care</i> , 2015, 19, 375.	2.5	24
15	Regulation of SATB1 during thymocyte development by TCR signaling. <i>Molecular Immunology</i> , 2016, 77, 34-43.	1.0	23
16	PYR-41, A Ubiquitin-Activating Enzyme E1 Inhibitor, Attenuates Lung Injury in Sepsis. <i>Shock</i> , 2018, 49, 442-450.	1.0	21
17	Specific and Randomly Derived Immunoactive Peptide Mimotopes of Mycobacterial Antigens. <i>Vaccine Journal</i> , 2006, 13, 1143-1154.	3.2	17
18	β -Catenin is required for the differentiation of iNKT2 and iNKT17 cells that augment IL-25-dependent lung inflammation. <i>BMC Immunology</i> , 2015, 16, 62.	0.9	17

#	ARTICLE	IF	CITATIONS
19	Sustained Expression of Pre-TCR Induced $\hat{\Gamma}^2$ -Catenin in Post- $\hat{\Gamma}^2$ -Selection Thymocytes Blocks T Cell Development. <i>Journal of Immunology</i> , 2009, 182, 759-765.	0.4	16
20	Molecular basis for the tissue specificity of $\hat{\Gamma}^2$ -catenin oncogenesis. <i>Oncogene</i> , 2013, 32, 1901-1909.	2.6	15
21	Receptor-Interacting Protein Kinase 3 Deficiency Delays Cutaneous Wound Healing. <i>PLoS ONE</i> , 2015, 10, e0140514.	1.1	13
22	T Cell Factor-1 Controls the Lifetime of CD4+ CD8+ Thymocytes In Vivo and Distal T Cell Receptor $\hat{\Gamma}^{\pm}$ -Chain Rearrangement Required for NKT Cell Development. <i>PLoS ONE</i> , 2014, 9, e115803.	1.1	12
23	Antagonists of Hsp16.3, a Low-Molecular-Weight Mycobacterial Chaperone and Virulence Factor, Derived from Phage-Displayed Peptide Libraries. <i>Applied and Environmental Microbiology</i> , 2005, 71, 7334-7344.	1.4	11
24	The protective role of human ghrelin in sepsis: Restoration of CD4 T cell proliferation. <i>PLoS ONE</i> , 2018, 13, e0201139.	1.1	10
25	IL-4 and IL-4 Receptor Expression Is Dispensable for the Development and Function of Natural Killer T Cells. <i>PLoS ONE</i> , 2013, 8, e71872.	1.1	2