

# Nonanzit PÃ©rez-HernÃ¡ndez

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9472820/publications.pdf>

Version: 2024-02-01

42  
papers

593  
citations

686830

13  
h-index

676716

22  
g-index

44  
all docs

44  
docs citations

44  
times ranked

1040  
citing authors

#	ARTICLE	IF	CITATIONS
1	Osteoprotegerin Gene Polymorphisms Are Associated with Subclinical Atherosclerosis in the Mexican Mestizo Population. <i>Diagnostics</i> , 2022, 12, 1433.	1.3	0
2	Evaluation of leptin levels in serum as a biomarker for suicide behavior: systematic review and meta-analysis. <i>International Journal of Neuroscience</i> , 2021, 131, 49-55.	0.8	11
3	Association between polymorphisms of FKBP5 gene and suicide attempt in a Mexican population: A case-control study. <i>Brain Research Bulletin</i> , 2021, 166, 37-43.	1.4	5
4	Are functional variants of the microRNA-146a gene associated with primary knee OA? Evidence in Mexican mestizo population. <i>Molecular Biology Reports</i> , 2021, 48, 1549-1557.	1.0	2
5	Variants of PCSK9 Gene Are Associated with Subclinical Atherosclerosis and Cardiometabolic Parameters in Mexicans. <i>The GEA Project. Diagnostics</i> , 2021, 11, 774.	1.3	7
6	Osteopontin Gene Polymorphisms Are Associated with Cardiovascular Risk Factors in Patients with Premature Coronary Artery Disease. <i>Biomedicines</i> , 2021, 9, 1600.	1.4	1
7	Quality of life of Latin-American individuals with type 2 diabetes mellitus: A systematic review. <i>Primary Care Diabetes</i> , 2020, 14, 317-334.	0.9	8
8	Risk of Wnt/ $\beta$ -catenin signalling pathway gene polymorphisms in primary SjÃ©gren's syndrome. <i>Rheumatology</i> , 2020, 59, 418-425.	0.9	9
9	DNA methyltransferase genes polymorphisms are associated with primary knee osteoarthritis: a matched case-control study. <i>Rheumatology International</i> , 2020, 40, 573-581.	1.5	8
10	Epstein-Barr virus-induced gene 3 (EBI3) single nucleotide polymorphisms and their association with central obesity and risk factors for cardiovascular disease: The GEA study. <i>Cytokine</i> , 2020, 135, 155225.	1.4	2
11	The role of peripheral cortisol levels in suicide behavior: A systematic review and meta-analysis of 30 studies. <i>Psychiatry Research</i> , 2020, 293, 113448.	1.7	12
12	Genetic Variants and Haplotypes in <i>OPG</i> Gene Are Associated with Premature Coronary Artery Disease and Traditional Cardiovascular Risk Factors in Mexican Population: The GEA Study. <i>DNA and Cell Biology</i> , 2020, 39, 2085-2094.	0.9	3
13	The rs46522 Polymorphism of the Ubiquitin-Conjugating Enzyme E2 Gene Is Associated with Abnormal Metabolic Parameters in Patients with Myocardial Infarction: The Genetics of Atherosclerosis Disease Mexican Study. <i>DNA and Cell Biology</i> , 2020, 39, 1155-1161.	0.9	2
14	Association between congenital heart disease and NKX2.5 gene polymorphisms: systematic review and meta-analysis. <i>Biomarkers in Medicine</i> , 2020, 14, 1747-1757.	0.6	1
15	Association between FKBP5 polymorphisms and depressive disorders or suicidal behavior: A systematic review and meta-analysis study. <i>Psychiatry Research</i> , 2019, 271, 658-668.	1.7	26
16	MRE11A Polymorphisms Are Associated With Subclinical Atherosclerosis and Cardiovascular Risk Factors. A Case-Control Study of the GEA Mexican Project. <i>Frontiers in Genetics</i> , 2019, 10, 530.	1.1	9
17	Association between polymorphisms of NOS1, NOS2 and NOS3 genes and suicide behavior: a systematic review and meta-analysis. <i>Metabolic Brain Disease</i> , 2019, 34, 967-977.	1.4	11
18	Interleukin 27 polymorphisms, their association with insulin resistance and their contribution to subclinical atherosclerosis. The GEA Mexican study. <i>Cytokine</i> , 2019, 114, 32-37.	1.4	14

#	ARTICLE	IF	CITATIONS
19	The $\hat{r}^2$ 44 C/G (rs1800972) polymorphism of the Î²â€defensin 1 is associated with increased risk of developing type 2 diabetes mellitus. <i>Molecular Genetics &amp; Genomic Medicine</i> , 2019, 7, e00509.	0.6	6
20	HHLPL-1 (rs2895811) gene polymorphism is associated with cardiovascular risk factors and cardiometabolic parameters in Mexicans patients with myocardial infarction. <i>Gene</i> , 2018, 663, 34-40.	1.0	10
21	Vitamin D Deficiency is not Associated with Fatty Liver in a Mexican Population. <i>Annals of Hepatology</i> , 2018, 17, 419-425.	0.6	6
22	Raet1e Polymorphisms Are Associated with Increased Risk of Developing Premature Coronary Artery Disease and with Some Cardiometabolic Parameters: The GEA Mexican Study. <i>Mediators of Inflammation</i> , 2018, 2018, 1-10.	1.4	3
23	Risk Factors and Prevalence of Suicide Attempt in Patients with Type 2 Diabetes in the Mexican Population. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 1198.	1.2	13
24	A haplotype of the phosphodiesterase 4D (PDE4D) gene is associated with myocardial infarction and with cardiometabolic parameters: the GEA study. <i>EXCLI Journal</i> , 2018, 17, 1182-1190.	0.5	1
25	Receptor-interacting protein 2 (RIP2) gene polymorphisms are associated with increased risk of subclinical atherosclerosis and clinical and metabolic parameters. The Genetics of Atherosclerotic Disease (GEA) Mexican study. <i>Experimental and Molecular Pathology</i> , 2017, 102, 1-6.	0.9	2
26	IL-15 polymorphisms are associated with subclinical atherosclerosis and cardiovascular risk factors. The Genetics of Atherosclerosis Disease (GEA) Mexican Study. <i>Cytokine</i> , 2017, 99, 173-178.	1.4	10
27	Association of the I148M/PNPLA3 (rs738409) polymorphism with premature coronary artery disease, fatty liver, and insulin resistance in type 2 diabetic patients and healthy controls. The GEA study. <i>Immunobiology</i> , 2017, 222, 960-966.	0.8	39
28	Vascular Calcification. <i>Chinese Medical Journal</i> , 2017, 130, 1113-1121.	0.9	25
29	Interleukin 35 Polymorphisms Are Associated with Decreased Risk of Premature Coronary Artery Disease, Metabolic Parameters, and IL-35 Levels: The Genetics of Atherosclerotic Disease (GEA) Study. <i>Mediators of Inflammation</i> , 2017, 2017, 1-10.	1.4	40
30	<i>HIF1A</i> (rs11549465) and <i>AKNA</i> (rs10817595) Gene Polymorphisms Are Associated with Primary SjÃ¶grenâ€™s Syndrome. <i>BioMed Research International</i> , 2017, 2017, 1-8.	0.9	10
31	Interleukin-27 polymorphisms are associated with premature coronary artery disease and metabolic parameters in the Mexican population: the genetics of atherosclerotic disease (GEA) Mexican study. <i>Oncotarget</i> , 2017, 8, 64459-64470.	0.8	31
32	PHACTR1 Gene Polymorphism Is Associated with Increased Risk of Developing Premature Coronary Artery Disease in Mexican Population. <i>International Journal of Environmental Research and Public Health</i> , 2016, 13, 803.	1.2	18
33	The Effect of Resveratrol and Quercetin Treatment on PPAR Mediated Uncoupling Protein (UCP-) 1, 2, and 3 Expression in Visceral White Adipose Tissue from Metabolic Syndrome Rats. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1069.	1.8	40
34	Possible role of intronic polymorphisms in the PHACTR1 gene on the development of cardiovascular disease. <i>Medical Hypotheses</i> , 2016, 97, 64-70.	0.8	10
35	Serum magnesium is inversely associated with coronary artery calcification in the Genetics of Atherosclerotic Disease (GEA) study. <i>Nutrition Journal</i> , 2015, 15, 22.	1.5	37
36	Interleukin-17A Gene Haplotypes Are Associated with Risk of Premature Coronary Artery Disease in Mexican Patients from the Genetics of Atherosclerotic Disease (GEA) Study. <i>PLoS ONE</i> , 2015, 10, e0114943.	1.1	21

#	ARTICLE	IF	CITATIONS
37	<i>Novel Mutations</i> in the Transcriptional Activator Domain of the Human TBX20 in Patients with Atrial Septal Defect. <i>BioMed Research International</i> , 2015, 2015, 1-7.	0.9	15
38	TIMP2 gene polymorphisms are associated with hypertension in patients with myocardial infarction. <i>Journal of Genetics</i> , 2014, 93, 517-522.	0.4	3
39	Protective role of DDAH2 (rs805304) gene polymorphism in patients with myocardial infarction. <i>Experimental and Molecular Pathology</i> , 2014, 97, 393-398.	0.9	7
40	The HIF1A rs2057482 polymorphism is associated with risk of developing premature coronary artery disease and with some metabolic and cardiovascular risk factors. The Genetics of Atherosclerotic Disease (GEA) Mexican Study. <i>Experimental and Molecular Pathology</i> , 2014, 96, 405-410.	0.9	18
41	The Matrix Metalloproteinase 2- <i>1575</i> gene Polymorphism is Associated with the Risk of Developing Myocardial Infarction in Mexican Patients. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 718-727.	0.9	27
42	Genetic admixture and diversity estimations in the Mexican Mestizo population from Mexico City using 15 STR polymorphic markers. <i>Forensic Science International: Genetics</i> , 2008, 2, e37-e39.	1.6	66