

Hyo-Jung Cho

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9475847/publications.pdf>

Version: 2024-02-01

30
papers

633
citations

623574

14
h-index

642610

23
g-index

30
all docs

30
docs citations

30
times ranked

909
citing authors

#	ARTICLE	IF	CITATIONS
1	Serum Exosomal MicroRNA, miR-10b-5p, as a Potential Diagnostic Biomarker for Early-Stage Hepatocellular Carcinoma. <i>Journal of Clinical Medicine</i> , 2020, 9, 281.	1.0	83
2	Nonalcoholic fatty liver disease as a sentinel marker for the development of diabetes mellitus in non-obese subjects. <i>Digestive and Liver Disease</i> , 2018, 50, 370-377.	0.4	45
3	Serum small extracellular vesicle-derived LINC00853 as a novel diagnostic marker for early hepatocellular carcinoma. <i>Molecular Oncology</i> , 2020, 14, 2646-2659.	2.1	45
4	High circulating microRNA-122 expression is a poor prognostic marker in patients with hepatitis B virus-related hepatocellular carcinoma who undergo radiofrequency ablation. <i>Clinical Biochemistry</i> , 2015, 48, 1073-1078.	0.8	40
5	Improvement of Nonalcoholic Fatty Liver Disease Reduces the Risk of Type 2 Diabetes Mellitus. <i>Gut and Liver</i> , 2019, 13, 440-449.	1.4	40
6	Low levels of circulating microRNA-26a/29a as poor prognostic markers in patients with hepatocellular carcinoma who underwent curative treatment. <i>Clinics and Research in Hepatology and Gastroenterology</i> , 2017, 41, 181-189.	0.7	37
7	Exosomal microRNA-4661-based serum panel as a potential diagnostic biomarker for early-stage hepatocellular carcinoma. <i>Cancer Medicine</i> , 2020, 9, 5459-5472.	1.3	35
8	Liver stiffness measured by MR elastography is a predictor of early HCC recurrence after treatment. <i>European Radiology</i> , 2020, 30, 4182-4192.	2.3	35
9	Accuracy and precision of proton density fat fraction measurement across field strengths and scan intervals: A phantom and human study. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 50, 305-314.	1.9	26
10	Plasma microRNA-122 as a predictive marker for treatment response following transarterial chemoembolization in patients with hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2017, 32, 199-207.	1.4	23
11	Early detection of hepatocellular carcinoma via liquid biopsy: panel of small extracellular vesicle-derived long noncoding RNAs identified as markers. <i>Molecular Oncology</i> , 2021, 15, 2715-2731.	2.1	22
12	Microbiome as a potential diagnostic and predictive biomarker in severe alcoholic hepatitis. <i>Alimentary Pharmacology and Therapeutics</i> , 2021, 53, 540-551.	1.9	22
13	Statin in combination with cisplatin makes favorable tumor-immune microenvironment for immunotherapy of head and neck squamous cell carcinoma. <i>Cancer Letters</i> , 2021, 522, 198-210.	3.2	20
14	Circulating Exosomal MicroRNA-1307-5p as a Predictor for Metastasis in Patients with Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 3819.	1.7	19
15	Plasma MicroRNA-21, 26a, and 29a-3p as Predictive Markers for Treatment Response Following Transarterial Chemoembolization in Patients with Hepatocellular Carcinoma. <i>Journal of Korean Medical Science</i> , 2018, 33, e6.	1.1	18
16	Role of Immune Cells in Patients with Hepatitis B Virus-Related Hepatocellular Carcinoma. <i>International Journal of Molecular Sciences</i> , 2021, 22, 8011.	1.8	18
17	Interleukin-8 level as a prognostic marker in patients with hepatitis B virus-associated hepatocellular carcinoma treated with transarterial chemoembolization. <i>Cytokine</i> , 2015, 76, 449-457.	1.4	16
18	Prediction of Post-Endoscopic Retrograde Cholangiopancreatography Pancreatitis Using 4-Hour Post-Endoscopic Retrograde Cholangiopancreatography Serum Amylase and Lipase Levels. <i>Journal of Korean Medical Science</i> , 2017, 32, 1814.	1.1	15

#	ARTICLE	IF	CITATIONS
19	Analysis of endoscopic features for histologic discrepancies between biopsy and endoscopic submucosal dissection in gastric neoplasms: 10-year results. <i>Digestive and Liver Disease</i> , 2019, 51, 79-85.	0.4	15
20	Liver stiffness in magnetic resonance elastography is prognostic for sorafenib-treated advanced hepatocellular carcinoma. <i>European Radiology</i> , 2021, 31, 2507-2517.	2.3	9
21	Novel Gene Signatures as Prognostic Biomarkers for Predicting the Recurrence of Hepatocellular Carcinoma. <i>Cancers</i> , 2022, 14, 865.	1.7	9
22	Hepatic Arterial Infusion Chemotherapy for Advanced Hepatocellular Carcinoma: Radiologic and Clinical Factors Predictive of Survival. <i>American Journal of Roentgenology</i> , 2021, 216, 1566-1573.	1.0	8
23	Independent Risk Factors for Hepatocellular Carcinoma Recurrence after Direct-Acting Antiviral Therapy in Patients with Chronic Hepatitis C. <i>Gut and Liver</i> , 2021, 15, 410-419.	1.4	7
24	Prognostic Value of Alpha-Fetoprotein in Patients Who Achieve a Complete Response to Transarterial Chemoembolization for Hepatocellular Carcinoma. <i>Yonsei Medical Journal</i> , 2021, 62, 12.	0.9	6
25	Overexpressed Proteins in HCC Cell-Derived Exosomes, CCT8, and Cofilin-1 Are Potential Biomarkers for Patients with HCC. <i>Diagnostics</i> , 2021, 11, 1221.	1.3	6
26	Risk Prediction Model Based on Magnetic Resonance Elastography-Assessed Liver Stiffness for Predicting Posthepatectomy Liver Failure in Patients with Hepatocellular Carcinoma. <i>Gut and Liver</i> , 2022, 16, 277-289.	1.4	5
27	Relationship between the dynamics of non-alcoholic fatty liver disease and incident diabetes mellitus. <i>Scientific Reports</i> , 2022, 12, 2538.	1.6	3
28	Effect of direct-acting antivirals for hepatitis C virus-related hepatocellular carcinoma recurrence and death after curative treatment. <i>Journal of Liver Cancer</i> , 2022, 22, 125-135.	0.3	3
29	Combining hepatic surface nodularity and serum tests better predicts hepatic fibrosis stages in chronic liver disease. <i>Abdominal Radiology</i> , 2021, 46, 4189-4199.	1.0	2
30	Infiltrative hepatocellular carcinoma with multiple lung metastasis completely cured using nivolumab: a case report. <i>Journal of Liver Cancer</i> , 2021, 21, 169-176.	0.3	1