

# Yutaka Fujiwara

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/948043/publications.pdf>

Version: 2024-02-01

119  
papers

3,077  
citations

147566

31  
h-index

197535

49  
g-index

119  
all docs

119  
docs citations

119  
times ranked

5463  
citing authors

#	ARTICLE	IF	CITATIONS
1	A Novel Serum Metabolomics-Based Diagnostic Approach to Pancreatic Cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2013, 22, 571-579.	1.1	157
2	High sensitivity detection of epidermal growth factor receptor mutations in the pleural effusion of non-small cell lung cancer patients. <i>Cancer Science</i> , 2006, 97, 642-648.	1.7	138
3	Phase I Dose-Escalation Study and Biomarker Analysis of E7080 in Patients with Advanced Solid Tumors. <i>Clinical Cancer Research</i> , 2011, 17, 2528-2537.	3.2	137
4	Association of antithyroglobulin antibodies with the development of thyroid dysfunction induced by nivolumab. <i>Cancer Science</i> , 2018, 109, 3583-3590.	1.7	118
5	Reliability of Small Biopsy Samples Compared With Resected Specimens for the Determination of Programmed Death-Ligand 1 Expression in Non-Small-Cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2015, 16, 385-390.	1.1	115
6	Foretinib (GSK1363089), a multi-kinase inhibitor of MET and VEGFRs, inhibits growth of gastric cancer cell lines by blocking inter-receptor tyrosine kinase networks. <i>Investigational New Drugs</i> , 2012, 30, 1352-1360.	1.2	100
7	PD-L1 expression in neuroendocrine tumors of the lung. <i>Lung Cancer</i> , 2017, 108, 115-120.	0.9	98
8	Expression of programmed death 1 (PD-1) and its ligand (PD-L1) in thymic epithelial tumors: Impact on treatment efficacy and alteration in expression after chemotherapy. <i>Lung Cancer</i> , 2016, 99, 4-10.	0.9	81
9	Phase 1 study of galunisertib, a TGF-beta receptor I kinase inhibitor, in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 76, 1143-1152.	1.1	73
10	Phase I and Pharmacokinetic Study of HER2-targeted rhuMAb 2C4 (Pertuzumab, RO4368451) in Japanese Patients with Solid Tumors. <i>Japanese Journal of Clinical Oncology</i> , 2009, 39, 260-266.	0.6	57
11	Management of axitinib (AG-013736)-induced fatigue and thyroid dysfunction, and predictive biomarkers of axitinib exposure: results from phase I studies in Japanese patients. <i>Investigational New Drugs</i> , 2012, 30, 1055-1064.	1.2	56
12	Change in the lymphocyte-to-monocyte ratio is an early surrogate marker of the efficacy of nivolumab monotherapy in advanced non-small-cell lung cancer. <i>Lung Cancer</i> , 2018, 124, 179-188.	0.9	56
13	Epidermal Growth Factor Receptor Mutation Is Associated With Longer Local Control After Definitive Chemoradiotherapy in Patients With Stage III Nonsquamous Non-Small-Cell Lung Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2015, 91, 140-148.	0.4	53
14	Phase 1 study of abemaciclib, an inhibitor of CDK 4 and 6, as a single agent for Japanese patients with advanced cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 281-288.	1.1	51
15	Picoliter-Droplet Digital Polymerase Chain Reaction-Based Analysis of Cell-Free Plasma DNA to Assess EGFR Mutations in Lung Adenocarcinoma That Confer Resistance to Tyrosine-Kinase Inhibitors. <i>Oncologist</i> , 2016, 21, 156-164.	1.9	50
16	Phase II study of nab-paclitaxel+carboplatin for patients with non-small cell lung cancer and interstitial lung disease. <i>Cancer Science</i> , 2019, 110, 3738-3745.	1.7	49
17	First-in-Human Phase I Study of an Oral HSP90 Inhibitor, TAS-116, in Patients with Advanced Solid Tumors. <i>Molecular Cancer Therapeutics</i> , 2019, 18, 531-540.	1.9	49
18	Phase I, dose escalation and pharmacokinetic study of cediranib (RECENTIN <sup>®</sup> ), a highly potent and selective VEGFR signaling inhibitor, in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 64, 1165-1172.	1.1	46

#	ARTICLE	IF	CITATIONS
19	Phase I study of ipilimumab in phased combination with paclitaxel and carboplatin in Japanese patients with non-small-cell lung cancer. <i>Investigational New Drugs</i> , 2015, 33, 881-889.	1.2	46
20	The genomic and epigenomic landscape in thymic carcinoma. <i>Carcinogenesis</i> , 2017, 38, 1084-1091.	1.3	46
21	Phase I dose-finding study of monotherapy with atezolizumab, an engineered immunoglobulin monoclonal antibody targeting PD-L1, in Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2016, 34, 596-603.	1.2	43
22	An overview of the recent progress in irinotecan pharmacogenetics. <i>Pharmacogenomics</i> , 2010, 11, 391-406.	0.6	41
23	Metabolomics Evaluation of Serum Markers for Cachexia and Their Intra-Day Variation in Patients with Advanced Pancreatic Cancer. <i>PLoS ONE</i> , 2014, 9, e113259.	1.1	40
24	Phase I and pharmacokinetic study of vorinostat (suberoylanilide hydroxamic acid) in Japanese patients with solid tumors. <i>Cancer Science</i> , 2009, 100, 1728-1734.	1.7	39
25	Circulating Endothelial Cells in Non-small Cell Lung Cancer Patients Treated with Carboplatin and Paclitaxel. <i>Journal of Thoracic Oncology</i> , 2009, 4, 208-213.	0.5	38
26	Retrospective analysis of the efficacy of chemotherapy and molecular targeted therapy for advanced pulmonary pleomorphic carcinoma. <i>BMC Research Notes</i> , 2015, 8, 800.	0.6	38
27	Phase I dose-escalating study of panobinostat (LBH589) Administered intravenously to Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2012, 30, 1950-1957.	1.2	37
28	A phase 1 and dose-finding study of LY2523355 (litronesib), an Eg5 inhibitor, in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2014, 74, 15-23.	1.1	37
29	Randomized trial of standard pain control with or without gabapentin for pain related to radiation-induced mucositis in head and neck cancer. <i>Auris Nasus Larynx</i> , 2016, 43, 677-684.	0.5	37
30	First-in-Human Phase 1 Study of MORAb-202, an Antibody-Drug Conjugate Comprising Farletuzumab Linked to Eribulin Mesylate, in Patients with Folate Receptor-1-Positive Advanced Solid Tumors. <i>Clinical Cancer Research</i> , 2021, 27, 3905-3915.	3.2	37
31	Radiologic features of pneumonitis associated with nivolumab in non-small-cell lung cancer and malignant melanoma. <i>Future Oncology</i> , 2019, 15, 1911-1920.	1.1	36
32	Effect of Platinum Combined with Irinotecan or Paclitaxel against Large Cell Neuroendocrine Carcinoma of the Lung. <i>Japanese Journal of Clinical Oncology</i> , 2007, 37, 482-486.	0.6	34
33	Safety and pharmacokinetics of milademetan, a MDM2 inhibitor, in Japanese patients with solid tumors: A phase I study. <i>Cancer Science</i> , 2021, 112, 2361-2370.	1.7	33
34	Safety and pharmacokinetics of DS-6051b in Japanese patients with non-small cell lung cancer harboring ROS1 fusions: a phase I study. <i>Oncotarget</i> , 2018, 9, 23729-23737.	0.8	33
35	Cytotoxic chemotherapy may overcome the development of acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors (EGFR-TKIs) therapy. <i>Lung Cancer</i> , 2015, 89, 287-293.	0.9	29
36	Phase I study of adjuvant gemcitabine or S-1 in patients with biliary tract cancers undergoing major hepatectomy: KHBO1003 study. <i>Cancer Chemotherapy and Pharmacology</i> , 2014, 74, 699-709.	1.1	27

#	ARTICLE	IF	CITATIONS
37	Phase I clinical and pharmacokinetic study of 3-weekly, 3-h infusion of ixabepilone (BMS-247550), an epothilone B analog, in Japanese patients with refractory solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2008, 61, 751-758.	1.1	26
38	Inhibition of the mTOR/S6K signal is necessary to enhance fluorouracil-induced apoptosis in gastric cancer cells with HER2 amplification. <i>International Journal of Oncology</i> , 2012, 41, 551-558.	1.4	26
39	Phase Ia/Ib study of the pan-class I PI3K inhibitor pictilisib (GDC-0941) administered as a single agent in Japanese patients with solid tumors and in combination in Japanese patients with non-squamous non-small cell lung cancer. <i>Investigational New Drugs</i> , 2017, 35, 37-46.	1.2	26
40	Phase 1 Study of Cabozantinib in Japanese Patients With Expansion Cohorts in Non-Small-Cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2019, 20, e317-e328.	1.1	26
41	A phase 1 study evaluating the pharmacokinetics and preliminary efficacy of veliparib (ABT-888) in combination with carboplatin/paclitaxel in Japanese subjects with non-small cell lung cancer (NSCLC). <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 76, 1063-1072.	1.1	25
42	Appearance of a BRAF Mutation Conferring Resistance to Crizotinib in Non-Small Cell Lung Cancer Harboring Oncogenic ROS1 Fusion. <i>Journal of Thoracic Oncology</i> , 2018, 13, e66-e69.	0.5	24
43	Validity of new methods to evaluate renal function in cancer patients treated with cisplatin. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 281-288.	1.1	23
44	Excessive MET signaling causes acquired resistance and addiction to MET inhibitors in the MKN45 gastric cancer cell line. <i>Investigational New Drugs</i> , 2013, 31, 1158-1168.	1.2	22
45	Phase 1/2 study assessing the safety and efficacy of dabrafenib and trametinib combination therapy in Japanese patients with BRAF V600 mutation-positive advanced cutaneous melanoma. <i>Journal of Dermatology</i> , 2018, 45, 397-407.	0.6	22
46	Efficacy of adjuvant chemotherapy for non-small cell lung cancer assessed by metastatic potential associated with ACTN4. <i>Oncotarget</i> , 2016, 7, 33165-33178.	0.8	22
47	Successful EGFR-TKI Rechallenge of Leptomeningeal Carcinomatosis after Gefitinib-induced Interstitial Lung Disease. <i>Japanese Journal of Clinical Oncology</i> , 2013, 43, 422-425.	0.6	20
48	Pharmacokinetic profiles of significant adverse events with crizotinib in Japanese patients with ABCB1 polymorphism. <i>Cancer Science</i> , 2016, 107, 1117-1123.	1.7	20
49	Malignant pleural effusion as a predictor of the efficacy of anti-PD-1 antibody in patients with non-small cell lung cancer. <i>Thoracic Cancer</i> , 2019, 10, 815-822.	0.8	20
50	Efficacy of Taltrectinib (AB-106/DS-6051b) in ROS1+ NSCLC: An Updated Pooled Analysis of U.S. and Japan Phase 1 Studies. <i>JTO Clinical and Research Reports</i> , 2021, 2, 100108.	0.6	20
51	Prediction of Glomerular Filtration Rate in Cancer Patients by an Equation for Japanese Estimated Glomerular Filtration Rate. <i>Japanese Journal of Clinical Oncology</i> , 2013, 43, 271-277.	0.6	19
52	A Minimum Of 100 Tumor Cells in a Single Biopsy Sample Is Required to Assess Programmed Cell Death Ligand 1 Expression in Predicting Patient Response to Nivolumab Treatment in Nonsquamous Non-Small Cell Lung Carcinoma. <i>Journal of Thoracic Oncology</i> , 2019, 14, 1818-1827.	0.5	18
53	Predictive value of serum VEGF levels for elderly patients or for patients with poor performance status receiving anti-PD-1 antibody therapy for advanced non-small-cell lung cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1229-1236.	2.0	18
54	Phase 1 study of the investigational, oral angiogenesis inhibitor motesanib in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2010, 66, 935-943.	1.1	17

#	ARTICLE	IF	CITATIONS
55	A phase I study of resminostat in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 75, 1155-1161.	1.1	17
56	Comparison of the pharmacokinetics of erlotinib administered in complete fasting and 2h after a meal in patients with lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 76, 125-132.	1.1	17
57	Comparison of Radiotherapy and Chemoradiotherapy for Locoregional Recurrence of Non-small-cell Lung Cancer Developing After Surgery. <i>Clinical Lung Cancer</i> , 2017, 18, e441-e448.	1.1	17
58	Mechanisms of acquired resistance to insulin-like growth factor 1 receptor inhibitor in MCF-7 breast cancer cell line. <i>Investigational New Drugs</i> , 2013, 31, 293-303.	1.2	16
59	Rapid improvement of glucagonoma-related necrolytic migratory erythema with octreotide. <i>Clinical Journal of Gastroenterology</i> , 2014, 7, 255-259.	0.4	16
60	Phase I, multicenter, open-label, dose-escalation study of sonidegib in Asian patients with advanced solid tumors. <i>Cancer Science</i> , 2016, 107, 1477-1483.	1.7	16
61	First-in-human phase I study of E7090, a novel selective fibroblast growth factor receptor inhibitor, in patients with advanced solid tumors. <i>Cancer Science</i> , 2020, 111, 571-579.	1.7	16
62	Pharmacokinetic Study of Adjuvant Gemcitabine Therapy for Biliary Tract Cancer following Major Hepatectomy (KHBO1101). <i>PLoS ONE</i> , 2015, 10, e0143072.	1.1	16
63	Impact of KRAS mutation on response and outcome of patients with stage III non-squamous non-small cell lung cancer. <i>Cancer Science</i> , 2015, 106, 1402-1407.	1.7	15
64	Phase I trial of volasertib, a Polo-like kinase inhibitor, in Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2016, 34, 66-74.	1.2	15
65	Sequential Use of Anaplastic Lymphoma Kinase Inhibitors in Japanese Patients With ALK-Rearranged Non-small-Cell Lung Cancer: A Retrospective Analysis. <i>Clinical Lung Cancer</i> , 2017, 18, e251-e258.	1.1	15
66	Actinin-4 protein overexpression as a predictive biomarker in adjuvant chemotherapy for resected lung adenocarcinoma. <i>Biomarkers in Medicine</i> , 2017, 11, 721-731.	0.6	15
67	Circulating cell-free plasma tumour DNA shows a higher incidence of EGFR mutations in patients with extrathoracic disease progression. <i>ESMO Open</i> , 2018, 3, e000292.	2.0	15
68	Left atrial extension of metastatic lung tumor via pulmonary vein: report on the first case of Ewing sarcoma. <i>Rare Tumors</i> , 2010, 2, 151-153.	0.3	14
69	Treatment and relapse of interstitial lung disease in nivolumab-treated patients with non-small cell lung cancer. <i>Cancer Science</i> , 2021, 112, 1506-1513.	1.7	14
70	Medical treatment involving investigational drugs and genetic profile of thymic carcinoma. <i>Lung Cancer</i> , 2016, 93, 77-81.	0.9	13
71	The first case of SMARCB1 (INI1) - deficient squamous cell carcinoma of the pleura: a case report. <i>BMC Cancer</i> , 2018, 18, 398.	1.1	13
72	Lenvatinib in combination with everolimus in patients with advanced or metastatic renal cell carcinoma: A phase I study. <i>International Journal of Urology</i> , 2018, 25, 922-928.	0.5	13

#	ARTICLE	IF	CITATIONS
73	Phase II trial of Sâ€1 treatment as palliativeâ€intent chemotherapy for previously treated advanced thymic carcinoma. <i>Cancer Medicine</i> , 2020, 9, 7418-7427.	1.3	13
74	Fiveâ€year safety and efficacy data from a phase Ib study of nivolumab and chemotherapy in advanced nonâ€smallâ€cell lung cancer. <i>Cancer Science</i> , 2020, 111, 1933-1942.	1.7	13
75	Clinicopathological Features in Young Patients Treated for Small-Cell Lung Cancer: Significance of Immunohistological and Molecular Analyses. <i>Clinical Lung Cancer</i> , 2014, 15, 244-247.	1.1	12
76	Tremelimumab-associated tumor regression following after initial progression: two case reports. <i>Immunotherapy</i> , 2016, 8, 9-15.	1.0	12
77	Efficacy and safety of osimertinib in a Japanese compassionate use program. <i>Japanese Journal of Clinical Oncology</i> , 2017, 47, 625-629.	0.6	12
78	Efficacy and safety of crizotinib in patients with ROS1 rearranged non-small cell lung cancer: a retrospective analysis. <i>Journal of Thoracic Disease</i> , 2019, 11, 2965-2972.	0.6	11
79	Risk factors for pneumonitis in patients with nonâ€small cell lung cancer treated with immune checkpoint inhibitors plus chemotherapy: A retrospective analysis. <i>Thoracic Cancer</i> , 2022, 13, 724-731.	0.8	10
80	Effects of Aprepitant on the Pharmacokinetics of Controlled-Release Oral Oxycodone in Cancer Patients. <i>PLoS ONE</i> , 2014, 9, e104215.	1.1	9
81	Phase II study of amrubicin at a dose of 45 mg/m <sup>2</sup> in patients with previously treated small-cell lung cancer. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 941-946.	0.6	9
82	Current Status of Single-Agent Phase I Trials in Japan: Toward Globalization. <i>Journal of Clinical Oncology</i> , 2015, 33, 2051-2061.	0.8	9
83	Phase I and pharmacokinetics/pharmacodynamics study of the MEK inhibitor RO4987655 in Japanese patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2015, 33, 641-651.	1.2	8
84	Candidates for Intensive Local Treatment in cIIIA-N2 Non-Small Cell Lung Cancer: Deciphering the Heterogeneity. <i>International Journal of Radiation Oncology Biology Physics</i> , 2016, 94, 155-162.	0.4	8
85	Safety, tolerability, and pharmacokinetic profile of dabrafenib in Japanese patients with BRAF V600 mutation-positive solid tumors: a phase 1 study. <i>Investigational New Drugs</i> , 2018, 36, 259-268.	1.2	8
86	Long-term survival without surgery in NSCLC patients with synchronous brain oligometastasis: systemic chemotherapy revisited. <i>Journal of Thoracic Disease</i> , 2018, 10, 1696-1702.	0.6	8
87	Instrumental evaluation sensitively detects subclinical skin changes by the epidermal growth factor receptor inhibitors and risk factors for severe acneiform eruption. <i>Journal of Dermatology</i> , 2019, 46, 18-25.	0.6	8
88	Radiographic features and poor prognostic factors of interstitial lung disease with nivolumab for nonâ€small cell lung cancer. <i>Cancer Science</i> , 2021, 112, 1495-1505.	1.7	8
89	A phase I study of tasisulam sodium using an albumin-tailored dose in Japanese patients with advanced solid tumors. <i>Cancer Chemotherapy and Pharmacology</i> , 2013, 71, 991-998.	1.1	6
90	Hypothyroidism in patients with colorectal carcinoma treated with fluoropyrimidines. <i>Oncology Reports</i> , 2013, 30, 1802-1806.	1.2	6

#	ARTICLE	IF	CITATIONS
91	Comparison of Amrubicin and Weekly Cisplatin/Etoposide/Irinotecan in Patients With Relapsed Small-cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2017, 18, 234-240.e2.	1.1	6
92	Evaluation of time to failure of strategy as an alternative surrogate endpoint in patients with lung cancer with EGFR mutations. <i>ESMO Open</i> , 2018, 3, e000399.	2.0	6
93	Pharmacokinetic study of the oral fluorouracil antitumor agent Sâ€1 in patients with impaired renal function. <i>Cancer Science</i> , 2019, 110, 1987-1994.	1.7	6
94	Mixed response to osimertinib and the beneficial effects of additional local therapy. <i>Thoracic Cancer</i> , 2019, 10, 738-743.	0.8	6
95	Exploration of germline variants responsible for adverse events of crizotinib in anaplastic lymphoma kinase-positive non-small cell lung cancer by target-gene panel sequencing. <i>Lung Cancer</i> , 2019, 128, 20-25.	0.9	6
96	Dose exploration results from Phase 1 study of cemiplimab, a human monoclonal programmed death (PD)-1 antibody, in Japanese patients with advanced malignancies. <i>Cancer Chemotherapy and Pharmacology</i> , 2021, 87, 53-64.	1.1	6
97	Firstâ€inâ€human study of the cancer peptide vaccine TAS0313 in patients with advanced solid tumors. <i>Cancer Science</i> , 2021, 112, 1514-1523.	1.7	6
98	Infection risk with PI3K-AKT-mTOR pathway inhibitors and immune checkpoint inhibitors in patients with advanced solid tumours in phase I clinical trials. <i>ESMO Open</i> , 2020, 5, e000653.	2.0	5
99	Phase 1 study of telisotuzumab vedotin in Japanese patients with advanced solid tumors. <i>Cancer Medicine</i> , 2021, 10, 2350-2358.	1.3	5
100	Effect of sequential chemoradiotherapy in patients with limited-disease small-cell lung cancer who were ineligible for concurrent therapy: a retrospective study at two institutions. <i>Japanese Journal of Clinical Oncology</i> , 2018, 48, 82-88.	0.6	4
101	Global trends in the distribution of cancer types among patients in oncology phase I trials, 1991â€2015. <i>Investigational New Drugs</i> , 2019, 37, 166-174.	1.2	4
102	Phase I study to evaluate the safety and tolerability of MEDI4736, an anti-programmed cell death ligand-1 (PD-L1) antibody, in Japanese patients with advanced solid tumors.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3039-3039.	0.8	4
103	Individual optimal dose of amrubicin to prevent severe neutropenia in Japanese patients with lung cancer. <i>Cancer Science</i> , 2019, 110, 3573-3583.	1.7	3
104	Dose Escalation Data from the Phase 1 Study of the Liposomal Formulation of Eribulin (E7389-LF) in Japanese Patients with Advanced Solid Tumors. <i>Clinical Cancer Research</i> , 2022, 28, 1783-1791.	3.2	3
105	Serum Total Bilirubin as a Predictive Factor for Severe Neutropenia in Lung Cancer Patients Treated with Cisplatin and Irinotecan. <i>Japanese Journal of Clinical Oncology</i> , 2007, 37, 358-364.	0.6	2
106	Regulation of MET Kinase Inhibitor Resistance by Copy Number of <I>MET</I> in Gastric Carcinoma Cells. <i>Oncology Research</i> , 2014, 21, 287-293.	0.6	2
107	Secondary Osteosarcoma Developing 10 Years after Chemoradiotherapy for Non-small-cell Lung Cancer. <i>Japanese Journal of Clinical Oncology</i> , 2014, 44, 191-194.	0.6	2
108	Surveillance of protocol deviations in Japanese oncology registration trials: a single institute experience. <i>Investigational New Drugs</i> , 2017, 35, 392-396.	1.2	2

#	ARTICLE	IF	CITATIONS
109	Next-Generation Sequencer Analysis of Pulmonary Pleomorphic Carcinoma With a CD74-ROS1 Fusion Successfully Treated With Crizotinib. <i>Journal of Thoracic Oncology</i> , 2019, 14, e106-e108.	0.5	2
110	Phase I Study of Tremelimumab Monotherapy or in Combination With Durvalumab in Japanese Patients With Advanced Solid Tumors or Malignant Mesothelioma. <i>Oncologist</i> , 2022, 27, e703-e722.	1.9	2
111	Successful neutrophil engraftment by reduced use of granulocyte colony-stimulating factor after allogeneic hematopoietic stem cell transplantation with mycophenolate mofetil for graft-versus-host disease prophylaxis. <i>International Journal of Hematology</i> , 2011, 93, 765-770.	0.7	1
112	Improved survival among patients enrolled in oncology phase 1 trials in recent decades. <i>Cancer Chemotherapy and Pharmacology</i> , 2020, 85, 449-459.	1.1	1
113	Phase I study of adjuvant chemotherapy with gemcitabine plus cisplatin in patients with biliary tract cancer undergoing curative resection without major hepatectomy (KHBO1004).. <i>Journal of Clinical Oncology</i> , 2014, 32, 347-347.	0.8	1
114	Adjuvant Chemotherapy in Patients with Completely Resected Small Cell Lung Cancer: A Retrospective Analysis of 26 Consecutive Cases. <i>Japanese Journal of Clinical Oncology</i> , 2014, 44, 835-840.	0.6	0
115	Do all patients in the phase I oncology trials need to be hospitalized? Domestic but outstanding issues for globalization of drug development in Japan. <i>International Journal of Clinical Oncology</i> , 2017, 22, 780-785.	1.0	0
116	Efficacy of surgery for skin cancers initially suspected to be carcinoma of unknown primary: a retrospective observational study. <i>International Journal of Dermatology</i> , 2021, , .	0.5	0
117	Programmed cell death 1 (PD-1) and its ligand (PD-L1) expression in thymic epithelial tumors (TETs): Impact on the treatment efficacy and alteration in expression after chemotherapy (C).. <i>Journal of Clinical Oncology</i> , 2015, 33, 7515-7515.	0.8	0
118	Possible utility of actinin-4 as a predictive biomarker of the efficacy of postoperative adjuvant chemotherapy for completely resected early stage lung adenocarcinoma.. <i>Journal of Clinical Oncology</i> , 2016, 34, e20003-e20003.	0.8	0
119	Successful <sc>IMRT</sc> and concurrent chemotherapy for a patient with intrathoracic extensiveâ€stage small cell lung cancer. <i>Respirology Case Reports</i> , 2022, 10, e0919.	0.3	0