

Gary W Saunders

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9485837/publications.pdf>

Version: 2024-02-01

195
papers

8,336
citations

53660

45
h-index

58464

82
g-index

199
all docs

199
docs citations

199
times ranked

4260
citing authors

#	ARTICLE	IF	CITATIONS
1	Ten people-centered rules for socially sustainable ecosystem restoration. <i>Restoration Ecology</i> , 2022, 30, e13574.	1.4	25
2	Molecular Analysis Resolves the Taxonomy of the <i>Laurencia</i> Complex (Rhodomelaceae, Ceramiales) in Bermuda and Uncover Novel Species of <i>Chondrophyucus</i> and <i>Laurenciella</i> . <i>Cryptogamie, Algologie</i> , 2022, 43, .	0.3	2
3	Three new species of <i>Asteromenia</i> (Hymenocladaceae, Rhodophyta) from Australia. <i>Botanica Marina</i> , 2022, .	0.6	0
4	Taxonomic investigation of <i>Ralfsia</i> -like (Ralfsiales, Phaeophyceae) taxa in the North Atlantic Ocean based on molecular and morphological data, with descriptions of <i>Pseudoralfsiaceae</i> fam. nov., <i>Pseudoralfsia azorica</i> gen. et sp. nov. and <i>Nuchella vesicularis</i> gen. et sp. nov.. <i>European Journal of Phycology</i> , 2021, 56, 12-23.	0.9	7
5	The Acrotylaceae (Gigartinales) revisited: molecular data indicate family-level differences in one of the most enigmatic red-algal families. <i>Australian Systematic Botany</i> , 2021, 34, 305-326.	0.3	1
6	Global biogeography and diversification of a group of brown seaweeds (Phaeophyceae) driven by clade-specific evolutionary processes. <i>Journal of Biogeography</i> , 2021, 48, 703-715.	1.4	19
7	First record of <i>Scinaia</i> cf. <i>johnstoniae</i> (Nemaliales, Rhodophyta) in Gwaii Haanas, British Columbia, Canada. <i>BiolInvasions Records</i> , 2021, 10, 270-276.	0.4	0
8	Chiixuu Tli iinasdli: Indigenous Ethics and Values Lead to Ecological Restoration for People and Place in Gwaii Haanas. <i>Ecological Restoration</i> , 2021, 39, 45-51.	0.5	9
9	On the nomenclatural reinstatement and lectotypification of <i>Spyridia americana</i> Durant (1850). <i>Botanica Marina</i> , 2021, 64, 221-225.	0.6	0
10	Revisiting a DNA barcode survey of Haida Gwaii kelp: the quest for <i>Eisenia arborea</i> (Arthrothamnaceae, Tj ETQq0 0,0rgBT /Oyerlock 10	0.5	1
11	<i>Lithothamnion</i> (Hapalidiales, Rhodophyta) in the changing Arctic and Subarctic: DNA sequencing of type and recent specimens provides a systematics foundation*. <i>European Journal of Phycology</i> , 2021, 56, 468-493.	0.9	13
12	Reinstatement of Indian Ocean <i>Porolithon coarctatum</i> and <i>P. Agardineri</i> based on sequencing type specimens, and <i>P. Âepiphyticum</i> <i>sp. nov.</i> (Corallinales, Rhodophyta), with comments on subfamilies Hydrolithoideae and Metagoniolithoideae. <i>Botanica Marina</i> , 2021, 64, 363-377.	0.6	3
13	Resurrection of <i>Plocamium pusillum</i> Sonder (Plocamiaceae, Rhodophyta) from Australia. <i>Cryptogamie, Algologie</i> , 2021, 42, .	0.3	1
14	Unique biodiversity in Arctic marine forests is shaped by diverse recolonization pathways and far northern glacial refugia. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 22590-22596.	3.3	33
15	<i>Commanderella</i> gen. nov. and new insights into foliose Kallymeniaceae (Rhodophyta) from the Russian Pacific coast based on molecular studies. <i>Phycologia</i> , 2020, 59, 200-207.	0.6	6
16	Two new species of Solieriaceae (Rhodophyta, Gigartinales) from the euphotic and mesophotic zones off Bermuda, <i>Meristotheca odontoloma</i> and <i>Tepoztequiella muriamans</i> . <i>Phycologia</i> , 2020, 59, 177-185.	0.6	4
17	<i>Eucheumatopsis sanibelensis</i> sp. nov. from the Gulf coast of Florida, USA. <i>Phytotaxa</i> , 2020, 440, 215-224.	0.1	1
18	Reassessment of Tristan da Cunha <i>Gelidium</i> (Gelidiales, Rhodophyta) species. <i>Botanica Marina</i> , 2020, 63, 455-462.	0.6	1

#	ARTICLE	IF	CITATIONS
19	Taxonomic study of the brown algal genus <i>Chorda</i> (Chordaceae, Laminariales) with description of the new species <i>Chorda borealis</i> from Alaska and northern Canada. <i>European Journal of Phycology</i> , 2019, 54, 509-517.	0.9	3
20	New species of <i>Galene</i> and <i>Howella</i> gen. nov. (Halymeniaceae, Rhodophyta) from the mesophotic zone off Bermuda. <i>Phycologia</i> , 2019, 58, 690-697.	0.6	10
21	A molecular survey of <i>Ralfsia</i> sensu stricto (Ralfsiales, Phaeophyceae) in Canada uncovers three new species: <i>R.</i> <i>robertii</i> sp. nov., <i>R.</i> <i>tenebris</i> sp. nov., and <i>R.</i> <i>unimaculata</i> sp. nov.. <i>Botany</i> , 2019, 97, 135-147.	0.5	4
22	Trans-Arctic speciation of Florideophyceae (Rhodophyta) since the opening of the Bering Strait, with consideration of the 'species pump' hypothesis. <i>Journal of Biogeography</i> , 2019, 46, 694-705.	1.4	15
23	DNA barcoding of the marine macroalgae from Nome, Alaska (Northern Bering Sea) reveals many trans-Arctic species. <i>Polar Biology</i> , 2019, 42, 851-864.	0.5	25
24	A DNA barcode survey of marine macroalgae from Bergen (Norway). <i>Marine Biology Research</i> , 2019, 15, 580-589.	0.3	15
25	Collections from the mesophytic zone off Bermuda reveal three species of Kallymeniaceae (Gigartinales, Rhodophyta) in genera with transoceanic distributions. <i>Journal of Phycology</i> , 2019, 55, 415-424.	1.0	9
26	A molecular assessment of species diversity and generic boundaries in the red algal tribes Polysiphoniae and Streblocladiae (Rhodomelaceae, Rhodophyta) in Canada. <i>European Journal of Phycology</i> , 2019, 54, 1-25.	0.9	30
27	The phylogeographic history of amphitropical <i>Callophyllis variegata</i> (Florideophyceae, Rhodophyta) in the Pacific Ocean. <i>Algae</i> , 2019, 34, 91-97.	0.9	4
28	Preliminary DNA Barcode Report on the Marine Red Algae (Rhodophyta) from the British Overseas Territory of Tristan da Cunha. <i>Cryptogamie, Algologie</i> , 2019, 40, 105.	0.3	9
29	DNA Barcoding Sheds Light on Novel Records in the Tunisian Red Algal Flora. <i>Cryptogamie, Algologie</i> , 2019, 40, 5.	0.3	13
30	Glacial vicariance drives phylogeographic diversification in the amphi-boreal kelp <i>Saccharina latissima</i> . <i>Scientific Reports</i> , 2018, 8, 1112.	1.6	61
31	<i>Ottia meiospora</i> (Ottiaceae, Rhodophyta), a new genus and family endophytic within the thallus of <i>Nothocladus</i> (Batrachospermales, Rhodophyta). <i>Journal of Phycology</i> , 2018, 54, 79-84.	1.0	7
32	Phylogenetic analyses of transcriptome data resolve familial assignments for genera of the red-algal Acrochaetiales-Palmariales Complex (Nemaliophycidae). <i>Molecular Phylogenetics and Evolution</i> , 2018, 119, 151-159.	1.2	31
33	Detecting <i>Alaria esculenta</i> and <i>Laminaria digitata</i> (Laminariales, Phaeophyceae) gametophytes in red algae, with consideration of distribution patterns in the intertidal zone. <i>Phycologia</i> , 2018, 57, 1-8.	0.6	6
34	Assessment of the order Rhodymeniales (Rhodophyta) from British Columbia using an integrative taxonomic approach reveals overlooked and cryptic species diversity. <i>Botany</i> , 2018, 96, 359-383.	0.5	5
35	Patterns and drivers of species diversity in the Indo-Pacific red seaweed <i>Portieria</i> . <i>Journal of Biogeography</i> , 2018, 45, 2299-2313.	1.4	46
36	A revision of the genus <i>Cryptonemia</i> (Halymeniaceae, Rhodophyta) in Bermuda, western Atlantic Ocean, including five new species and <i>C. bermudensis</i> (Collins & M. Howe) comb. nov.. <i>European Journal of Phycology</i> , 2018, 53, 350-368.	0.9	10

#	ARTICLE	IF	CITATIONS
37	A new species of Digenea (Rhodomelaceae, Ceramiales) based upon a molecular assessment and morphological observations of plants historically known as <i>D. simplex</i> in Bermuda. <i>Phytotaxa</i> , 2018, 338, 90.	0.1	8
38	Intensive land-based production of red and green macroalgae for human consumption in the Pacific Northwest: an evaluation of seasonal growth, yield, nutritional composition, and contaminant levels. <i>Algae</i> , 2018, 33, 109-125.	0.9	17
39	<i>Calliblepharis rammediorum</i> sp. nov. (Gigartinales, Rhodophyta) from the Israeli Levant Mediterranean Sea. <i>Cryptogamie, Algologie</i> , 2018, 39, 109-121.	0.3	3
40	Notes on the Marine Algae of the Bermudas. 16. Two New Epiphytic Species of <i>Champia</i> (Champiaceae, Tj ETQq0 0.0 rgBT /Overlock 10	0.3	3
41	PCR fishing for red endophytes in British Columbia Kallymeniaceae (Gigartinales, Tj ETQq1 1 0.784314 rgBT /Overlock 10	1.0	5
42	A contaminant DNA barcode sequence reveals a new red algal order, Corynodactylales (Nemaliophycidae, Florideophyceae). <i>Botany</i> , 2017, 95, 561-566.	0.5	4
43	Genetic barcoding resolves the historically known red alga <i>Champia parvula</i> from southern New England, USA, as <i>C. farlowii</i> sp. nov. (Champiaceae, Rhodymeniales). <i>Phytotaxa</i> , 2017, 302, 77.	0.1	7
44	Key Kamchatkan collections provide new taxonomic and distributional insights for reportedly panâ€œNorth Pacific species of Rhodymeniophycidae (Rhodophyta). <i>Phycologia</i> , 2017, 56, 296-302.	0.6	4
45	A molecular-assisted investigation of diversity, biogeography and phylogenetic relationships for species of <i>Neoptilota</i> and <i>Ptilota</i> (Wrangeliaceae, Rhodophyta) reported along Canadian coasts. <i>Phycologia</i> , 2017, 56, 36-53.	0.6	4
46	Mychodea and the Mychodeaceae (Gigartinales, Rhodophyta) revisited: molecular analyses shed light on interspecies relationships in Australiaâ€™s largest endemic algal genus and family. <i>Australian Systematic Botany</i> , 2017, 30, 230.	0.3	4
47	A molecular investigation of Canadian Scytosiphonaceae (Phaeophyceae) including descriptions of <i>Planosiphon</i> gen. nov. and <i>Scytosiphon promiscuus</i> sp. nov.. <i>Botany</i> , 2017, 95, 653-671.	0.5	19
48	Kelp transcriptomes provide robust support for interfamilial relationships and revision of the little known Arthrothamnaceae (Laminariales). <i>Journal of Phycology</i> , 2017, 53, 1-6.	1.0	28
49	Updates to the Marine Algal Flora of the Boulder Patch in the Beaufort Sea off Northern Alaska as Revealed by DNA Barcoding + Supplementary Appendix 1 (See Article Tools). <i>Arctic</i> , 2017, 70, .	0.2	8
50	Phylogenetic Analyses Support Recognition of Ten New Genera, Ten New Species and 16 New Combinations in the Family Kallymeniaceae (Gigartinales, Rhodophyta). <i>Cryptogamie, Algologie</i> , 2017, 38, 79.	0.3	30
51	Divergence time estimates and the evolution of major lineages in the florideophyte red algae. <i>Scientific Reports</i> , 2016, 6, 21361.	1.6	139
52	Molecular-assisted alpha taxonomy of the genus <i>Rhodymenia</i> (Rhodymeniaceae, Rhodymeniales) from Australia reveals overlooked species diversity. <i>European Journal of Phycology</i> , 2016, 51, 354-367.	0.9	10
53	A molecular phylogenetic and DNA barcode assessment of the tribe Pterosiphonieae (Ceramiales, Tj ETQq1 1 0.784314 rgBT /Overlock 0.5 25	0.5	25
54	Endemic or introduced? Phylogeography of <i>Asparagopsis</i> (Florideophyceae) in Australia reveals multiple introductions and a new mitochondrial lineage. <i>Journal of Phycology</i> , 2016, 52, 141-147.	1.0	26

#	ARTICLE	IF	CITATIONS
55	A new monotypic family for the enigmatic crustose red alga <i>Plagiospora gracilis</i> . Botanical Journal of the Linnean Society, 2016, 182, 1-13.	0.8	8
56	Multigene analyses resolve early diverging lineages in the Rhodymeniophycidae (Florideophyceae). Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50	1.0	10
57	<i>Rhytymenia</i> , a new genus of red algae based on the rare <i>Kallymenia maculata</i> (Kallymeniaceae). Tj ETQq1 1 0.784314 rgBT/Overlock 1	0.8	6
58	Application of multigene phylogenetics and siteâ€stripping to resolve intraordinal relationships in the Rhodymeniales (Rhodophyta). Journal of Phycology, 2016, 52, 339-355.	1.0	14
59	Characterization of the putatively introduced red alga <i>Acrochaetium secundatum</i> (Acrochaetiales, Rhodophyta) growing epizoically on the pelage of southern sea otters (<i>Enhydra</i>). Tj ETQq1 1 0.784314 rgBT/Overlock	1.0	14
60	Multigene phylogeny of the red algal subclass Nemaliophycidae. Molecular Phylogenetics and Evolution, 2016, 94, 730-736.	1.2	21
61	A re-examination of the genus <i>Leptofaucheia</i> (Faucheaceae, Rhodymeniales) with clarification of species in Australia and the northwest Pacific. Phycologia, 2015, 54, 375-384.	0.6	5
62	Reproductive morphology and <i>scp>DNA</scp></i> sequences of the brown alga <i>Platysiphon verticillatus</i> support the new combination <i>Platysiphon glacialis</i> . Journal of Phycology, 2015, 51, 910-917.	1.0	15
63	Evidence for the introduction of the Asian red alga <i>Neosiphonia japonica</i> and its introgression with <i>Neosiphonia harveyi</i> (Ceramiales, Rhodophyta) in the Northwest Atlantic. Molecular Ecology, 2015, 24, 5927-5937.	2.0	25
64	Etheliaceae fam. nov. (Gigartinales, Rhodophyta), with a clarification of the generitype of <i>Ethelia</i> and the addition of six novel species from warm waters. Journal of Phycology, 2015, 51, 1158-1171.	1.0	13
65	Population genetic analyses are consistent with the introduction of <i>Ceramium secundatum</i> (Ceramiaceae, Rhodophyta) to Narragansett Bay, Rhode Island, USA. Ecology and Evolution, 2015, 5, 5088-5095.	0.8	3
66	A DNA barcode survey of <i>Schizymenia</i> (Nemastomatales, Rhodophyta) in Australia and British Columbia reveals overlooked diversity including <i>S. tenuis</i> sp. nov. and <i>Predaea borealis</i> sp. nov. Botany, 2015, 93, 859-871.	0.5	10
67	On the utility of mucilage ducts as a taxonomic character in <i>Laminaria</i> and <i>Saccharina</i> (Phaeophyceae) â€“ the conundrum of <i>S. groenlandica</i> . Phycologia, 2015, 54, 440-450.	0.6	19
68	A DNA barcode survey of the red algal genus <i>Mazzaella</i> in British Columbia reveals overlooked diversity and new distributional records: descriptions of <i>M. dewreedei</i> sp. nov. and <i>M. macrocarpa</i> sp. nov.. Botany, 2014, 92, 223-231.	0.5	18
69	A floristic survey of marine tube-forming diatoms reveals unexpected diversity and extensive co-habitation among genetic lines of the <i>Berkeleya rutilans</i> complex (Bacillariophyceae). European Journal of Phycology, 2014, 49, 47-59.	0.9	16
70	A study of two <i>Acrochaetium</i> complexes in Canada with distinction of <i>Rhododrewia</i> gen. nov. (Acrochaetiales, Rhodophyta). Phycologia, 2014, 53, 221-232.	0.6	9
71	<i>Crebradomus</i> and <i>Dissimularia</i> , new genera in the family Chondrymeniaceae (Gigartinales, Rhodophyta) from the central, southern and western Pacific Ocean. Phycologia, 2014, 53, 146-166.	0.6	11
72	A DNA barcode survey of Haida Gwaii kelp (Laminariales, Phaeophyceae) reveals novel ecological and distributional observations and <i>Saccharina druehlii</i> sp. nov.. Botany, 2014, 92, 821-826.	0.5	10

#	ARTICLE	IF	CITATIONS
73	The monospecific genus <i>Meredithia</i> (<i>Kallymeniaceae</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 75 <i>atlantic</i> , <i>Pacific</i> , and <i>Indian Oceans</i> . <i>Journal of Phycology</i> , 2014, 50, 167-186.	1.0	17
74	Long distance kelp rafting impacts seaweed biogeography in the Northeast Pacific: the kelp conveyor hypothesis. <i>Journal of Phycology</i> , 2014, 50, 968-974.	1.0	33
75	Evidence for genotypic differentiation between marine snails (<i>Littorina sitkana</i>) from the upper- and lower-intertidal zone in Bamfield Inlet (British Columbia, Canada). <i>Journal of Experimental Marine Biology and Ecology</i> , 2014, 461, 389-396.	0.7	7
76	Molecular-assisted alpha taxonomy reveals pseudocryptic diversity among species of <i>Bossiella</i> (<i>Corallinales</i> , <i>Rhodophyta</i>) in the eastern Pacific Ocean. <i>Phycologia</i> , 2014, 53, 443-456.	0.6	44
77	Molecular markers from three organellar genomes unravel complex taxonomic relationships within the coralline algal genus <i>Chiharaea</i> (<i>Corallinales</i> , <i>Rhodophyta</i>). <i>Molecular Phylogenetics and Evolution</i> , 2013, 67, 529-540.	1.2	20
78	DNA barcoding unmask overlooked diversity improving knowledge on the composition and origins of the Churchill algal flora. <i>BMC Ecology</i> , 2013, 13, 9.	3.0	32
79	Notes on the Marine Algae of the Bermudas. 13. <i>Helminthocladia kempii</i> sp. nov. (<i>Nemaliales</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 75 <i>atlantic</i> . <i>Cryptogamie, Algologie</i> , 2013, 34, 229-244.	0.3	4
80	<i>Fredericia deveauniensis</i> , gen. et sp. nov. (<i>Phyllophoraceae</i> , <i>Rhodophyta</i>), a New Cryptogenic Species. <i>Cryptogamie, Algologie</i> , 2013, 34, 273-296.	0.3	15
81	Resolving species diversity in the red algal genus <i>Callophyllis</i> (<i>Kallymeniaceae</i> , <i>Gigartinales</i>) in Canada using molecular assisted alpha taxonomy. <i>European Journal of Phycology</i> , 2013, 48, 27-46.	0.9	19
82	DNA barcoding and phylogenetics of <i>Ramicrusta</i> and <i>Incendia</i> gen. nov., two early diverging lineages of the <i>Peyssonneliaceae</i> (<i>Rhodophyta</i>). <i>Phycologia</i> , 2013, 52, 82-108.	0.6	20
83	<i>Pseudopolyides furcellarioides</i> gen. et sp. nov. (<i>Gigartinales</i> , <i>Rhodophyta</i>) an erect member of the <i>Cruoriaceae</i> based on morphological and molecular evidence. <i>Phycologia</i> , 2013, 52, 191-203.	0.6	7
84	A Molecular Phylogenetic Study of the Tribe <i>Corallineae</i> (<i>Corallinales</i> , <i>Rhodophyta</i>) with an Assessment of Genus-Level Taxonomic Features and Descriptions of Novel Genera. <i>Journal of Phycology</i> , 2013, 49, 103-114.	1.0	70
85	<i>Entwisleia bella</i> , gen. et sp. nov., a novel marine "batrachospermaceous" red alga from southeastern Tasmania representing a new family and order in the <i>Nemaliophycidae</i> . <i>European Journal of Phycology</i> , 2013, 48, 398-410.	0.9	14
86	A Molecular Survey of <i>Ulva</i> (<i>Chlorophyta</i>) in Temperate Australia Reveals Enhanced Levels of Cosmopolitanism. <i>Journal of Phycology</i> , 2013, 49, 69-81.	1.0	86
87	A Comparison of Morphological and Molecular-Based Surveys to Estimate the Species Richness of <i>Chaetoceros</i> and <i>Thalassiosira</i> (<i>Bacillariophyta</i>), in the Bay of Fundy. <i>PLoS ONE</i> , 2013, 8, e73521.	1.1	24
88	First record of the invasive red alga <i>Heterosiphonia japonica</i> (<i>Ceramiales</i> , <i>Rhodophyta</i>) in Canada. <i>BioInvasions Records</i> , 2013, 2, 27-32.	0.4	15
89	A survey of Sam Orr's Pond (New Brunswick, Canada) uncovers the invasive green alga <i>Codium fragile</i> (<i>Chlorophyta</i>) and the orange-striped green anemone <i>Diadumene lineata</i> (<i>Cnidaria</i>), first records for the Bay of Fundy and Canada, respectively. <i>BioInvasions Records</i> , 2013, 2, 185-189.	0.4	4
90	Refinements for the amplification and sequencing of red algal DNA barcode and RedToL phylogenetic markers: a summary of current primers, profiles and strategies. <i>Algae</i> , 2013, 28, 31-43.	0.9	193

#	ARTICLE	IF	CITATIONS
91	CBOL Protist Working Group: Barcoding Eukaryotic Richness beyond the Animal, Plant, and Fungal Kingdoms. <i>PLoS Biology</i> , 2012, 10, e1001419.	2.6	488
92	Acquiring DNA sequence data from dried archival red algae (Florideophyceae) for the purpose of applying available names to contemporary genetic species: a critical assessment. <i>Botany</i> , 2012, 90, 191-203.	0.5	53
93	DNA barcoding of Canadian Ahnfeltiales (Rhodophyta) reveals a new species " <i>Ahnfeltia borealis</i> sp. nov.. <i>Phycologia</i> , 2012, 51, 247-259.	0.6	18
94	An examination of the red algal genus <i>Pugetia</i> (Kallymeniaceae, Gigartinales), with descriptions of <i>Salishia firma</i> gen. & comb. nov., <i>Pugetia cryptica</i> sp. nov. and <i>Beringia wynnei</i> sp. nov.. <i>Phycologia</i> , 2012, 51, 33-61.	0.6	29
95	Methods for DNA Barcoding Photosynthetic Protists Emphasizing the Macroalgae and Diatoms. <i>Methods in Molecular Biology</i> , 2012, 858, 207-222.	0.4	183
96	A SURVEY OF BANGIALES (RHODOPHYTA) BASED ON MULTIPLE MOLECULAR MARKERS REVEALS CRYPTIC DIVERSITY. <i>Journal of Phycology</i> , 2012, 48, 869-882.	1.0	65
97	First report of <i>Halopeltis</i> (Rhodophyta, Rhodymeniaceae) from the non-tropical Northern Hemisphere: <i>H. adnata</i> (Okamura) comb. nov. from Korea, and <i>H. pellucida</i> sp. nov. and <i>H. willisii</i> sp. nov. from the North Atlantic. <i>Algae</i> , 2012, 27, 95-108.	0.9	7
98	Taxonomic assessment of North American species of the genera <i>Cumathamnion</i> , <i>Delesseria</i> , <i>Membranoptera</i> and <i>Pantoneura</i> (Delesseriaceae, Rhodophyta) using molecular data. <i>Algae</i> , 2012, 27, 155-173.	0.9	10
99	Notes on the Marine Algae of the Bermudas. 12. A phylogenetic Assessment of <i>Nemastoma gelatinosum</i> M. Howe (Rhodophyta, Nemastomatales) from its Type Locality. <i>Cryptogamie, Algologie</i> , 2011, 32, 313-325.	0.3	4
100	A new genus and species from the North Atlantic, <i>Archestenogramma profundum</i> (Phyllophoraceae, Rhodophyta), with taxonomic resolution of the orphaned <i>Leptofauchea brasiliensis</i> . <i>European Journal of Phycology</i> , 2011, 46, 442-452.	0.9	7
101	A new taxonomic interpretation of the type of <i>Plocamium cartilagineum</i> (Plocamiales), <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10</i>	0.9	18
102	A multigene phylogenetic assessment of the <i>Dilsea/Neodilsea</i> species complex (Dumontiaceae), <i>Tj ETQq0 0 0 rgBT /Overlock 10</i> <i>Marina</i> , 2011, 54, 481-486.	0.6	4
103	Taxonomic and molecular studies of the family <i>Sebdeniaceae</i> (Sebdeniales, Rhodophyta): new species of <i>Lesleigha</i> gen. nov. and <i>Crassitegula</i> from Hawaii, east Asia and Lord Howe Island. <i>European Journal of Phycology</i> , 2011, 46, 416-441.	0.9	7
104	Barcoding Diatoms: Exploring Alternatives to COI-5P. <i>Protist</i> , 2011, 162, 405-422.	0.6	165
105	Data mining approach identifies research priorities and data requirements for resolving the red algal tree of life. <i>BMC Evolutionary Biology</i> , 2010, 10, 16.	3.2	101
106	Multigene phylogenetic analyses support recognition of the <i>Sporolithales</i> ord. nov.. <i>Molecular Phylogenetics and Evolution</i> , 2010, 54, 302-305.	1.2	77
107	DNA BARCODING IS A POWERFUL TOOL TO UNCOVER ALGAL DIVERSITY: A CASE STUDY OF THE PHYLLOPHORACEAE (GIGARTINALES, RHODOPHYTA) IN THE CANADIAN FLORA. <i>Journal of Phycology</i> , 2010, 46, 374-389.	1.0	180
108	Further to the Occurrence of Red Abalone, <i>Haliotis rufescens</i> , in British Columbia. <i>Canadian Field-Naturalist</i> , 2010, 124, 238.	0.0	2

#	ARTICLE	IF	CITATIONS
109	DNA barcoding reveals multiple overlooked Australian species of the red algal order Rhodymeniales (Florideophyceae), with resurrection of <i>Halopeltis</i> J.ÂAgardh and description of <i>Pseudohalopeltis</i> gen. nov.. Botany, 2010, 88, 639-667.	0.5	39
110	A comparison of two DNA barcode markers for species discrimination in the red algal family Kallymeniaceae (Gigartinales, Florideophyceae), with a description of <i>Euthora timburtonii</i> sp.â€%nov.. Botany, 2010, 88, 119-131.	0.5	75
111	A DNA barcode examination of the Laminariaceae (Phaeophyceae) in Canada reveals novel biogeographical and evolutionary insights. Phycologia, 2010, 49, 235-248.	0.6	93
112	Notes on the marine algae of the Bermudas. 11. More additions to the benthic flora and a phylogenetic assessment of <i>Halymenia pseudofloresii</i> (Halymeniales, Rhodophyta) from its type locality. Phycologia, 2010, 49, 154-168.	0.6	29
113	Recognition of <i>Rubrointrusa membranacea</i> gen. et comb. nov., <i>Rhodonematella subimmersa</i> gen. et comb. nov. (with a reinterpretation of the life history) and the Meiodiscaceae fam. nov. within the Palmariales (Rhodophyta). Phycologia, 2010, 49, 283-300.	0.6	25
114	Providing a valid epithet for the species widely known as Halosaccocolax kjellmanii S. Lund (Palmariales, Rhodophyta) â€“ Rhodophysema kjellmanii sp. nov. Phycologia, 2010, 49, 628-628.	0.6	5
115	On the utility of DNA barcoding for species differentiation among brown macroalgae (Phaeophyceae) including a novel extraction protocol. Phycological Research, 2009, 57, 131-141.	0.8	81
116	Historical versus contemporary measures of seaweed biodiversity in the Bay of Fundy. Botany, 2009, 87, 1066-1076.	0.5	9
117	<i>Ramicrusta textilis</i> sp. nov. (Peyssonneliaceae, Rhodophyta), an anatomically complex Caribbean alga that overgrows corals. Phycologia, 2009, 48, 480-491.	0.6	39
118	Routine DNA barcoding of Canadian Gracilariales (Rhodophyta) reveals the invasive species <i>Gracilaria vermiculophylla</i> in British Columbia. Molecular Ecology Resources, 2009, 9, 140-150.	2.2	114
119	RESOLVING EVOLUTIONARY RELATIONSHIPS AMONG THE BROWN ALGAE USING CHLOROPLAST AND NUCLEAR GENES ¹ . Journal of Phycology, 2008, 44, 394-405.	1.0	58
120	PHYLOGENETIC RELATIONSHIPS AMONG LINEAGES OF THE CERAMIACEAE (CERAMIALES, RHODOPHYTA) BASED ON NUCLEAR SMALL SUBUNIT rDNA SEQUENCE DATA ¹ . Journal of Phycology, 2008, 44, 1033-1048.	1.0	25
121	PHYLOGENETIC ANALYSES OF THE RED ALGAL ORDER RHODYMENIALES SUPPORTS RECOGNITION OF THE HYMENOCLADIACEAE FAM. NOV., FRYEELLACEAE FAM. NOV., AND <i>NEOGASTROCLONIUM</i> GEN. NOV. ¹ . Journal of Phycology, 2008, 44, 1556-1571.	1.0	40
122	Assigning morphological variants of <i>Fucus</i> (Fucales, Phaeophyceae) in Canadian waters to recognized species using DNA barcoding. Botany, 2008, 86, 1065-1079.	0.5	82
123	A DNA barcode examination of the red algal family Dumontiaceae in Canadian waters reveals substantial cryptic species diversity. 1. The foliose <i>Dilsea</i> â€“ <i>Neodilsea</i> complex and <i>Weeksia</i> This paper is one of a selection of papers published in the Special Issue on Systematics Research.. Botany, 2008, 86, 773-789.	0.5	145
124	<i>Rhodachlya madagascarensis</i> gen. et sp. nov.: a distinct acrochaetioid represents a new order and family (Rhodachlyales ord. Nov., Rhodachlyaceae fam. Nov.) of the Florideophyceae (Rhodophyta). Phycologia, 2008, 47, 203-212.	0.6	21
125	Resurrecting the red algal genus <i>Grania</i> within the order Acrochaetiales (Florideophyceae, Rhodophyta). European Journal of Phycology, 2008, 43, 151-160.	0.9	10
126	A review of the red algal genus <i>Leptofauchea</i> (Faucheaceae, Rhodymeniales) including a description of <i>L. chilensis</i> sp. nov. Phycologia, 2007, 46, 198-213.	0.6	16

#	ARTICLE	IF	CITATIONS
127	Establishment of the onset of host specificity in four phyllobothriid tapeworm species (Cestoda): Tj ETQq1 1 0.784314 rgBT /Overlock 18	0.7	18
128	A nuclear phylogeny of the Florideophyceae (Rhodophyta) inferred from combined EF2, small subunit and large subunit ribosomal DNA: Establishing the new red algal subclass Corallinophycidae. <i>Molecular Phylogenetics and Evolution</i> , 2007, 43, 1118-1130.	1.2	141
129	A molecular assessment of northeast Pacific <i>Alaria</i> species (Laminariales, Phaeophyceae) with reference to the utility of DNA barcoding. <i>Molecular Phylogenetics and Evolution</i> , 2007, 44, 634-648.	1.2	172
130	Large subunit rDNA and rbcL gene sequence data place <i>Petrohwa bernabei</i> gen. et sp. nov. in the Batrachospermales (Rhodophyta), but do not provide further resolution among taxa in this order. <i>Phycological Research</i> , 2007, 55, 103-112.	0.8	30
131	<i>Halichrysis corallinarius</i> sp. nov. (Rhodymeniaceae, Rhodophyta) from Puerto Rico, Caribbean Sea. <i>Phycological Research</i> , 2007, 55, 240-248.	0.8	7
132	Combining small and large subunit ribosomal DNA genes to resolve relationships among orders of the Rhodymeniophycidae (Rhodophyta): recognition of the Acrosymphytales ord. nov. and Sebdeniales ord. nov.. <i>European Journal of Phycology</i> , 2006, 41, 379-394.	0.9	37
133	Unraveling the <i>Asteromenia peltata</i> species complex with clarification of the genera <i>Halichrysis</i> and <i>Drouetia</i> (Rhodymeniaceae, Rhodophyta). <i>Canadian Journal of Botany</i> , 2006, 84, 1581-1607.	1.2	31
134	Collections of the invasive species <i>Grateloupia turuturu</i> (Halymeniales, Rhodophyta) from Tasmania, Australia. <i>Phycologia</i> , 2006, 45, 711-714.	0.6	26
135	A MULTI-GENE MOLECULAR INVESTIGATION OF THE KELP (LAMINARIALES, PHAEOPHYCEAE) SUPPORTS SUBSTANTIAL TAXONOMIC RE-ORGANIZATION1. <i>Journal of Phycology</i> , 2006, 42, 493-512.	1.0	262
136	<i>Crassitegula walsinghamii</i> (Sebdeniaceae, Halymeniales), a new red algal genus and species from Bermuda based upon morphology and SSU rDNA sequence analyses. <i>European Journal of Phycology</i> , 2006, 41, 115-124.	0.9	17
137	AN EVALUATION OF METHODS USED TO ASSESS INTERGENERIC HYBRIDIZATION IN KELP USING PACIFIC LAMINARIALES (PHAEOPHYCEAE)1. <i>Journal of Phycology</i> , 2005, 41, 250-262.	1.0	49
138	Molecular divergence and morphological diversity among four cryptic species of <i>Plocamium</i> (Plocamiales, Florideophyceae) in northern Europe. <i>European Journal of Phycology</i> , 2005, 40, 293-312.	0.9	68
139	An assessment of two taxonomic distinctness indices for detecting seaweed assemblage responses to environmental stress. <i>Botanica Marina</i> , 2005, 48, .	0.6	13
140	Molecular investigation reveals epi/endophytic extrageneric kelp (Laminariales, Phaeophyceae) gametophytes colonizing <i>Lessoniopsis littoralis</i> thalli. <i>Botanica Marina</i> , 2005, 48, .	0.6	16
141	Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future applications. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2005, 360, 1879-1888.	1.8	644
142	Phylogenetic study of the Nemaliales (Rhodophyta) based on large-subunit ribosomal DNA sequences supports segregation of the Scinaiaceae fam. nov. and resurrection of <i>Dichotomaria</i> Lamarck. <i>Phycological Research</i> , 2004, 52, 224-234.	0.8	28
143	A UNIQUELY CALCIFIED BROWN ALGA FROM HAWAII: NEWHOUSIA IMBRICATA GEN. ET SP. NOV. (DICTYOTALES, PHAEOPHYCEAE)1. <i>Journal of Phycology</i> , 2004, 40, 383-394.	1.0	29
144	THE INVASIVE GENUS <i>ASPARAGOPSIS</i> (BONNEMAISONIACEAE, RHODOPHYTA): MOLECULAR SYSTEMATICS, MORPHOLOGY, AND ECOPHYSIOLOGY OFFALKENBERGIAISOLATES. <i>Journal of Phycology</i> , 2004, 40, 1112-1126.	1.0	75

#	ARTICLE	IF	CITATIONS
145	Sexing Arctic Terns in the Field and Laboratory. <i>Waterbirds</i> , 2004, 27, 314-320.	0.2	40
146	Small-subunit rDNA sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta). 3. Delineating the Gigartinales sensu stricto. <i>Canadian Journal of Botany</i> , 2004, 82, 43-74.	1.2	83
147	Assessing red algal supraordinal diversity and taxonomy in the context of contemporary systematic data. <i>American Journal of Botany</i> , 2004, 91, 1494-1507.	0.8	156
148	Phylogenetic study of the Nemaliales (Rhodophyta) based on large-subunit ribosomal DNA sequences supports segregation of the Scinaiaceae fam. nov. and resurrection of <i>Dichotomaria</i> Lamarck. <i>Phycological Research</i> , 2004, 52, 224-234.	0.8	29
149	Nuclear rDNA sequences from <i>Ballia prieurii</i> support recognition of <i>Balliopsis</i> gen. nov. in the Batrachospermales (Florideophyceae, Rhodophyta). <i>Phycologia</i> , 2002, 41, 61-67.	0.6	18
150	Phylogenetic analyses of anatomical and nuclear SSU rDNA sequence data indicate that the Dasyaceae and Delesseriaceae (Ceramiales, Rhodophyta) are polyphyletic. <i>European Journal of Phycology</i> , 2002, 37, 551-569.	0.9	36
151	A re-classification of the Acrochaetiales based on molecular and morphological data, and establishment of the Colaconematales ord. nov. (Florideophyceae, Rhodophyta). <i>European Journal of Phycology</i> , 2002, 37, 463-476.	0.9	60
152	CRYPTOMONAD EVOLUTION: NUCLEAR 18S rDNA PHYLOGENY VERSUS CELL MORPHOLOGY AND PIGMENTATION1. <i>Journal of Phycology</i> , 2002, 38, 1236-1244.	1.0	84
153	TWO NEW AUSTRALIAN SPECIES OF PREDAEA (NEMASTOMATACEAE, RHODOPHYTA) WITH TAXONOMIC RECOMMENDATIONS FOR AN EMENDED NEMASTOMATALES AND EXPANDED HALYMENIALES1. <i>Journal of Phycology</i> , 2002, 38, 1245-1260.	1.0	34
154	Using molecular data to resolve the taxonomic limits of the genera <i>Callophyllis</i> , <i>Euthora</i> and <i>Pugetia</i> (Kallymeniaceae, Rhodophyta). <i>Phycological Research</i> , 2002, 50, 275-281.	0.8	13
155	Phylogenetic relationships of <i>Polysiphonia</i> (Rhodomelaceae, Rhodophyta) and its relatives based on anatomical and nuclear small-subunit rDNA sequence data. <i>Canadian Journal of Botany</i> , 2001, 79, 1465-1476.	1.2	17
156	Affinities of the freshwater red alga <i>Audouinella macrospora</i> (Florideophyceae, Rhodophyta) and related forms based on ssu rRNA gene sequence analysis and pit plug ultrastructure. <i>Journal of Phycology</i> , 2001, 36, 433-440.	1.0	24
157	MOLECULAR SYSTEMATICS OF THE FLORIDEOPHYCEAE (RHODOPHYTA) USING NUCLEAR LARGE AND SMALL SUBUNIT rDNA SEQUENCE DATA. <i>Journal of Phycology</i> , 2001, 37, 1073-1082.	1.0	106
158	PHYLOGENY OF THE DUMONTIACEAE (GIGARTINALES, RHODOPHYTA) AND ASSOCIATED FAMILIES BASED ON SSU rDNA AND INTERNAL TRANSCRIBED SPACER SEQUENCE DATA. <i>Journal of Phycology</i> , 2001, 37, 184-196.	1.0	71
159	Phylogenetic relationships of <i>Polysiphonia</i> (Rhodomelaceae, Rhodophyta) and its relatives based on anatomical and nuclear small-subunit rDNA sequence data. <i>Canadian Journal of Botany</i> , 2001, 79, 1465-1476.	1.2	43
160	The morphology, taxonomy, and molecular phylogeny of <i>Heterocladia</i> and <i>Trigenea</i> (rhodomelaceae,) <i>TJ ETQq0 0 0 rgBT /Overlock 10 Tf</i> 36, 199-219.	1.0	14
161	Title is missing!. <i>Journal of Applied Phycology</i> , 2000, 12, 25-35.	1.5	33
162	Nuclear small-subunit rDNA sequences from <i>Ballia</i> spp. (Rhodophyta): proposal of the Balliales ord. nov., Balliaceae fam. nov., <i>Ballia nana</i> sp. nov. and <i>Inkyuleea</i> gen. nov. (Ceramiales). <i>Phycologia</i> , 2000, 39, 272-287.	0.6	28

#	ARTICLE	IF	CITATIONS
163	Bringing order to red algal families: taxonomists ask the jurists 'Who's in charge here?'. <i>Phycologia</i> , 2000, 39, 358-361.	0.6	7
164	MOLECULAR SYSTEMATIC ANALYSES INDICATE THAT THE ENIGMATIC APOPHLAEA IS A MEMBER OF THE HILDENBRANDIALES (RHODOPHYTA, FLORIDEOPHYCIDAE). <i>Journal of Phycology</i> , 1999, 35, 171-175.	1.0	23
165	The families of the order Rhodymeniales (Rhodophyta): a molecular-systematic investigation with a description of Faucheaceae fam. nov.. <i>Phycologia</i> , 1999, 38, 23-40.	0.6	59
166	Amplified Fragment Length Polymorphism (AFLP) as a source of genetic markers for red algae. <i>Journal of Applied Phycology</i> , 1998, 10, 365-370.	1.5	25
167	PHYLOGENY OF THE BATRACHOSPERMALES (RHODOPHYTA) INFERRED FROM <i>rbcL</i> AND 18S RIBOSOMAL DNA GENE SEQUENCES. <i>Journal of Phycology</i> , 1998, 34, 341-350.	1.0	101
168	A molecular systematic investigation of the Acrochaetiales (Florideophycidae, Rhodophyta) and related taxa based on nuclear small-subunit ribosomal DNA sequence data. <i>European Journal of Phycology</i> , 1998, 33, 221-229.	0.9	24
169	Phylogenetic relationships of the Raphidophyceae and Xanthophyceae as inferred from nucleotide sequences of the 18s ribosomal RNA gene. <i>American Journal of Botany</i> , 1997, 84, 966-972.	0.8	62
170	Phylogenesis of pit-plug-associated features in the Rhodophyta: inferences from molecular systematic data. <i>Canadian Journal of Botany</i> , 1997, 75, 1436-1447.	1.2	54
171	Phylogenetic relationships of the "golden algae"™ (haptophytes, heterokont chromophytes) and their plastids. <i>Plant Systematics and Evolution Supplementum = Entwicklungsgeschichte Und Systematik Der Pflanzen Supplementum</i> , 1997, , 187-219.	1.5	114
172	PHYLOGENETIC AFFINITIES OF CHRYSONEPHELE PALUSTRIS (CHRYSOPHYCEAE) BASED ON INFERRED NUCLEAR SMALL-SUBUNIT RIBOSOMAL RNA SEQUENCE1. <i>Journal of Phycology</i> , 1997, 33, 132-134.	1.0	8
173	A PHYLOGENETIC ANALYSIS OF THE SYNUROPHYCEAE USING MOLECULAR DATA AND SCALE CASE MORPHOLOGY1. <i>Journal of Phycology</i> , 1997, 33, 135-151.	1.0	31
174	PHYLOGENETIC AFFINITIES OF THE SARCIINOCHRYSIDALES AND CHRYSOMERIDALES (HETEROKONTA) BASED ON ANALYSES OF MOLECULAR AND COMBINED DATA1. <i>Journal of Phycology</i> , 1997, 33, 310-318.	1.0	67
175	Nuclear small-subunit ribosomal RNA gene sequences from representative Ceramiaceae (Ceramiiales,) Tj ETQq1 1 0.784314 rgBT /Ove	0.9	28
176	The morphology, reproduction and small-subunit rRNA gene sequence of <i>Cephalocystis</i> (Rhodymeniaceae, Rhodophyta), a new genus based on <i>Cordylecladia furcellata</i> J. Agardh from Australia. <i>Phycologia</i> , 1996, 35, 48-60.	0.6	18
177	Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta). 2. Recognition of the Halymeniales ord.nov.. <i>Canadian Journal of Botany</i> , 1996, 74, 694-707.	1.2	112
178	Cladistic analyses of combined traditional and molecular data sets reveal an algal lineage.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 244-248.	3.3	95
179	A REVISION OF THE SYSTEMATICS OF THE NIZYMENIACEAE(GIGARTINALES, RHODOPHYTA) BASED ON POLYSACCHARIDES, ANATOMY, AND NUCLEOTIDE SEQUENCES1. <i>Journal of Phycology</i> , 1995, 31, 153-166.	1.0	33
180	PHYLOGENETIC RELATIONSHIPS OF SPECIES OF UNCERTAIN TAXONOMIC POSITION WITHIN THE ACROCHAETIALES-PALMARIALES COMPLEX (RHODOPHYTA): INFERENCES FROM PHENOTYPIC AND 18S rDNA SEQUENCE DATA1. <i>Journal of Phycology</i> , 1995, 31, 601-611.	1.0	34

#	ARTICLE	IF	CITATIONS
181	The phylogenetic affinities of <i>Notheia anomala</i> (Fucales, Phaeophyceae) as determined from partial small-subunit rRNA gene sequences. <i>Phycologia</i> , 1995, 34, 383-389.	0.6	35
182	Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta). 1. Evidence for the Plocamiales ord.nov.. <i>Canadian Journal of Botany</i> , 1994, 72, 1250-1263.	1.2	194
183	Revision of the kelp family Alariaceae and the taxonomic affinities of <i>Lessoniopsis Reinke</i> (Laminariales), Tj ETQq1 1,0,784314,rgBT /O	1.0	27
184	GEL PURIFICATION OF RED ALGAL GENOMIC DNA: AN INEXPENSIVE AND RAPID METHOD FOR THE ISOLATION OF POLYMERASE CHAIN REACTION-FRIENDLY DNA1. <i>Journal of Phycology</i> , 1993, 29, 251-254.	1.0	200
185	ULTRASTRUCTURE AND 18SRRNA GENE SEQUENCE FORPELAGOMONAS CALCEOLATAGEN. ET SP. NOV. AND THE DESCRIPTION OF A NEW ALGAL CLASS, THE PELAGOPHYCEAE CLASSIS NOV.1. <i>Journal of Phycology</i> , 1993, 29, 701-715.	1.0	217
186	When is a family not a family?. <i>BioSystems</i> , 1992, 28, 109-116.	0.9	17
187	NUCLEOTIDE SEQUENCES OF THE SMALL-SUBUNIT RIBOSOMAL RNA GENES FROM SELECTED LAMINARIALES (PHAEOPHYTA): IMPLICATIONS FOR KELP EVOLUTION1. <i>Journal of Phycology</i> , 1992, 28, 544-549.	1.0	103
188	DNA EXTRACTION METHODS FOR KELP (LAMINARIALES) TISSUE1. <i>Journal of Phycology</i> , 1992, 28, 712-716.	1.0	45
189	Morphology and reproduction of <i>Meiodiscus spetsbergensis</i> (Kjellman) gen. et comb. nov., a new genus of Rhodophysemataceae (Rhodophyta). <i>Phycologia</i> , 1991, 30, 272-286.	0.6	14
190	Restriction enzyme mapping of the nuclear ribosomal cistron in selected Laminariales (Phaeophyta): a phylogenetic assessment. <i>Canadian Journal of Botany</i> , 1991, 69, 2647-2654.	1.2	14
191	Biology of <i>Furcellaria lumbricalis</i> (Hudson) Lamouroux (Rhodophyta: Gigartinales), a commercial carrageenophyte. <i>Journal of Applied Phycology</i> , 1991, 3, 61-82.	1.5	27
192	Considerations of life history, morphology and taxonomy in <i>Rhodophysema georgii</i> Batters (Rhodophyta, Palmariales). <i>British Phycological Journal</i> , 1989, 24, 63-71.	1.3	7
193	Life-history variation in <i>Rhodophysema elegans</i> (Palmariales, Rhodophyta) from the North Atlantic and crustose <i>Rhodophysema</i> spp. from the North Pacific. <i>Canadian Journal of Botany</i> , 1989, 67, 2857-2872.	1.2	13
194	INFRAGENERIC TAXONOMY OF AHNFELTIA (AHNFELTIALES, RHODOPHYTA). <i>Journal of Phycology</i> , 1989, 25, 351-368.	1.0	22
195	The pseudodichotomous <i>Dasya sylviae</i> sp.Ânov. (Delesseriaceae, Ceramiales) from 60â€“90 m mesophotic reefs off Bermuda. <i>European Journal of Taxonomy</i> , 0, 751, 24-37.	0.6	3