

Jan-Ingvar Jansson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9486493/publications.pdf>

Version: 2024-02-01

42
papers

1,813
citations

361296

20
h-index

265120

42
g-index

43
all docs

43
docs citations

43
times ranked

3276
citing authors

#	ARTICLE	IF	CITATIONS
1	Multiplexed single-cell mass cytometry reveals distinct inhibitory effects on intracellular phosphoproteins by midostaurin in combination with chemotherapy in AML cells. <i>Experimental Hematology and Oncology</i> , 2021, 10, 7.	2.0	4
2	The Critical Role of Dysregulated RhoB Signaling Pathway in Radioresistance of Colorectal Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019, 104, 1153-1164.	0.4	17
3	Putative Role of Nuclear Factor-Kappa B But Not Hypoxia-Inducible Factor-1 β in Hypoxia-Dependent Regulation of Oxidative Stress in Hematopoietic Stem and Progenitor Cells. <i>Antioxidants and Redox Signaling</i> , 2019, 31, 211-226.	2.5	7
4	Switchable presentation of cytokines on electroactive polypyrrole surfaces for hematopoietic stem and progenitor cells. <i>Journal of Materials Chemistry B</i> , 2018, 6, 4665-4675.	2.9	6
5	The stem cell regulator PEDF is dispensable for maintenance and function of hematopoietic stem cells. <i>Scientific Reports</i> , 2017, 7, 10134.	1.6	4
6	Hypoxia Mediates Differential Response to Anti-EGFR Therapy in HNSCC Cells. <i>International Journal of Molecular Sciences</i> , 2017, 18, 943.	1.8	20
7	Pyruvate dehydrogenase kinase 1 is essential for transplantable mouse bone marrow hematopoietic stem cell and progenitor function. <i>PLoS ONE</i> , 2017, 12, e0171714.	1.1	19
8	Coexpression of hyperactivated AKT1 with additional genes activated in leukemia drives hematopoietic progenitor cells to cell cycle block and apoptosis. <i>Experimental Hematology</i> , 2015, 43, 554-564.	0.2	6
9	miR-20b regulates expression of proteinase-activated receptor-1 (PAR-1) thrombin receptor in melanoma cells. <i>Pigment Cell and Melanoma Research</i> , 2014, 27, 431-441.	1.5	27
10	Single-nucleotide polymorphisms of ABCG2 increase the efficacy of tyrosine kinase inhibitors in the K562 chronic myeloid leukemia cell line. <i>Pharmacogenetics and Genomics</i> , 2014, 24, 52-61.	0.7	34
11	Interleukin-1 β induced activation of the hypothalamus-pituitary-adrenal axis is dependent on interleukin-1 receptors on non-hematopoietic cells. <i>Brain, Behavior, and Immunity</i> , 2014, 40, 166-173.	2.0	34
12	Interleukin-6 primarily produced by non-hematopoietic cells mediates the lipopolysaccharide-induced febrile response. <i>Brain, Behavior, and Immunity</i> , 2013, 33, 123-130.	2.0	18
13	ABCB1 haplotypes do not influence transport or efficacy of tyrosine kinase inhibitors in vitro. <i>Pharmacogenomics and Personalized Medicine</i> , 2013, 6, 63.	0.4	18
14	Hypoxic and normoxic in vitro cultures maintain similar numbers of long-term reconstituting hematopoietic stem cells from mouse bone marrow. <i>Experimental Hematology</i> , 2012, 40, 879-881.	0.2	1
15	Lipopolysaccharide-Induced Fever Depends on Prostaglandin E2 Production Specifically in Brain Endothelial Cells. <i>Endocrinology</i> , 2012, 153, 4849-4861.	1.4	87
16	The Q705K Polymorphism in NLRP3 Is a Gain-of-Function Alteration Leading to Excessive Interleukin-1 β and IL-18 Production. <i>PLoS ONE</i> , 2012, 7, e34977.	1.1	127
17	The pan-ErbB tyrosine kinase inhibitor canertinib induces caspase-mediated cell death in human T-cell leukemia (Jurkat) cells. <i>Biochemical and Biophysical Research Communications</i> , 2011, 410, 422-427.	1.0	8
18	IL-7 mediates Ebf-1-dependent lineage restriction in early lymphoid progenitors. <i>Blood</i> , 2011, 118, 1283-1290.	0.6	80

#	ARTICLE	IF	CITATIONS
19	Irreversible pan-ERBB inhibitor canertinib elicits anti-leukaemic effects and induces the regression of FLT3-ITD transformed cells in mice. <i>British Journal of Haematology</i> , 2011, 155, 198-208.	1.2	7
20	Hypoxia mediates low cell-cycle activity and increases the proportion of long-term-reconstituting hematopoietic stem cells during in vitro culture. <i>Experimental Hematology</i> , 2010, 38, 301-310.e2.	0.2	143
21	The hematopoietic stem cell niche: Low in oxygen but a nice place to be. <i>Journal of Cellular Physiology</i> , 2010, 222, 17-22.	2.0	394
22	Interleukin-7-induced Stat-5 Acts in Synergy with Flt-3 Signaling to Stimulate Expansion of Hematopoietic Progenitor Cells. <i>Journal of Biological Chemistry</i> , 2010, 285, 36275-36284.	1.6	28
23	The pan-ErbB receptor tyrosine kinase inhibitor canertinib induces ErbB-independent apoptosis in human leukemia (HL-60 and U-937) cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 393, 6-10.	1.0	9
24	Deficiency of Activating Fcγ3-Receptors Reduces Hepatic Clearance and Deposition of IC and Increases CIC Levels in Mercury-Induced Autoimmunity. <i>PLoS ONE</i> , 2010, 5, e13413.	1.1	8
25	Systemic Reduction of Functionally Suppressive CD4 ^{dim} CD25 ^{high} Foxp3 ⁺ Tregs in Human Second Trimester Pregnancy Is Induced by Progesterone and 17β-Estradiol. <i>Journal of Immunology</i> , 2009, 183, 759-769.	0.4	136
26	BH3-only protein Bim more critical than Puma in tyrosine kinase inhibitor-induced apoptosis of human leukemic cells and transduced hematopoietic progenitors carrying oncogenic FLT3. <i>Blood</i> , 2009, 113, 2302-2311.	0.6	31
27	The Lim-only protein LMO2 acts as a positive regulator of erythroid differentiation. <i>Biochemical and Biophysical Research Communications</i> , 2007, 364, 675-681.	1.0	15
28	The BH3-only protein Puma plays an essential role in cytokine deprivation-induced apoptosis of mast cells. <i>Blood</i> , 2007, 110, 3209-3217.	0.6	103
29	Bcl11b mutations identified in murine lymphomas increase the proliferation rate of hematopoietic progenitor cells. <i>BMC Cancer</i> , 2007, 7, 195.	1.1	12
30	Absence of hot spot mutations of the PIK3CA gene in acute myeloid leukaemia. <i>European Journal of Haematology</i> , 2006, 77, 86-87.	1.1	12
31	Stem cell factor promotes mast cell survival via inactivation of FOXO3a-mediated transcriptional induction and MEK-regulated phosphorylation of the proapoptotic protein Bim. <i>Blood</i> , 2005, 106, 1330-1336.	0.6	109
32	Characterization of the Mouse Myeloid-associated Differentiation Marker (Myadm) Gene: Promoter Analysis and Protein Localization. <i>Molecular Biology Reports</i> , 2005, 32, 149-157.	1.0	7
33	FLT3 ligand regulates apoptosis through AKT-dependent inactivation of transcription factor FoxO3. <i>Biochemical and Biophysical Research Communications</i> , 2004, 318, 899-903.	1.0	32
34	Inactivation of the forkhead transcription factor FoxO3 is essential for PKB-mediated survival of hematopoietic progenitor cells by kit ligand. <i>Experimental Hematology</i> , 2003, 31, 316-323.	0.2	40
35	The basic helix-loop-helix transcription factor TAL1/SCL inhibits the expression of the p16INK4A and pT± genes. <i>Biochemical and Biophysical Research Communications</i> , 2003, 312, 1073-1081.	1.0	20
36	The Lim-only protein LMO4 modulates the transcriptional activity of HEN1. <i>Biochemical and Biophysical Research Communications</i> , 2003, 307, 891-899.	1.0	39

#	ARTICLE	IF	CITATIONS
37	Phosphatidylinositol 3-kinase is essential for kit ligand-mediated survival, whereas interleukin-3 and flt3 ligand induce expression of antiapoptotic Bcl-2 family genes. <i>Journal of Leukocyte Biology</i> , 2003, 74, 923-931.	1.5	25
38	Distinct and regulated expression of Notch receptors in hematopoietic lineages and during myeloid differentiation. <i>European Journal of Immunology</i> , 2001, 31, 3240-3247.	1.6	49
39	Isolation of MYADM, a novel hematopoietic-associated marker gene expressed in multipotent progenitor cells and up-regulated during myeloid differentiation. <i>Journal of Leukocyte Biology</i> , 2000, 67, 423-431.	1.5	25
40	Characterization, Chromosomal Localization, and Expression during Hematopoietic Differentiation of the Gene Encoding Arl6ip, ADP-Ribosylation-like Factor-6 Interacting Protein (ARL6). <i>Genomics</i> , 2000, 68, 351-354.	1.3	30
41	Concentration-Dependent Effects of Hematopoietic Growth Factors during In Vitro Expansion of Mouse Stem Cells and Progenitor Cells. <i>Growth Factors</i> , 1997, 14, 59-66.	0.5	3
42	Interleukin-7 Responsiveness of B220+ B Cell Precursors from Bone Marrow Decreases in Aging Mice. <i>Cellular Immunology</i> , 1993, 147, 267-278.	1.4	19