

Jan-Ingvar Jnsson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9486493/jan-ingvar-jonsson-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

42
papers

1,539
citations

20
h-index

39
g-index

43
ext. papers

1,706
ext. citations

4.3
avg, IF

4.37
L-index

#	Paper	IF	Citations
42	Multiplexed single-cell mass cytometry reveals distinct inhibitory effects on intracellular phosphoproteins by midostaurin in combination with chemotherapy in AML cells. <i>Experimental Hematology and Oncology</i> , 2021 , 10, 7	7.8	0
41	Putative Role of Nuclear Factor-Kappa B But Not Hypoxia-Inducible Factor-1 α in Hypoxia-Dependent Regulation of Oxidative Stress in Hematopoietic Stem and Progenitor Cells. <i>Antioxidants and Redox Signaling</i> , 2019 , 31, 211-226	8.4	6
40	The Critical Role of Dysregulated RhoB Signaling Pathway in Radioresistance of Colorectal Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019 , 104, 1153-1164	4	7
39	Switchable presentation of cytokines on electroactive polypyrrole surfaces for hematopoietic stem and progenitor cells. <i>Journal of Materials Chemistry B</i> , 2018 , 6, 4665-4675	7.3	6
38	Hypoxia Mediates Differential Response to Anti-EGFR Therapy in HNSCC Cells. <i>International Journal of Molecular Sciences</i> , 2017 , 18,	6.3	20
37	The stem cell regulator PEDF is dispensable for maintenance and function of hematopoietic stem cells. <i>Scientific Reports</i> , 2017 , 7, 10134	4.9	2
36	Pyruvate dehydrogenase kinase 1 is essential for transplantable mouse bone marrow hematopoietic stem cell and progenitor function. <i>PLoS ONE</i> , 2017 , 12, e0171714	3.7	13
35	Coexpression of hyperactivated AKT1 with additional genes activated in leukemia drives hematopoietic progenitor cells to cell cycle block and apoptosis. <i>Experimental Hematology</i> , 2015 , 43, 554-64	3.1	4
34	Interleukin-1 α -induced activation of the hypothalamus-pituitary-adrenal axis is dependent on interleukin-1 receptors on non-hematopoietic cells. <i>Brain, Behavior, and Immunity</i> , 2014 , 40, 166-73	16.6	23
33	miR-20b regulates expression of proteinase-activated receptor-1 (PAR-1) thrombin receptor in melanoma cells. <i>Pigment Cell and Melanoma Research</i> , 2014 , 27, 431-41	4.5	24
32	Single-nucleotide polymorphisms of ABCG2 increase the efficacy of tyrosine kinase inhibitors in the K562 chronic myeloid leukemia cell line. <i>Pharmacogenetics and Genomics</i> , 2014 , 24, 52-61	1.9	28
31	Interleukin-6 primarily produced by non-hematopoietic cells mediates the lipopolysaccharide-induced febrile response. <i>Brain, Behavior, and Immunity</i> , 2013 , 33, 123-30	16.6	15
30	ABCB1 haplotypes do not influence transport or efficacy of tyrosine kinase inhibitors in vitro. <i>Pharmacogenomics and Personalized Medicine</i> , 2013 , 6, 63-72	2.1	16
29	Hypoxic and normoxic in vitro cultures maintain similar numbers of long-term reconstituting hematopoietic stem cells from mouse bone marrow. <i>Experimental Hematology</i> , 2012 , 40, 879-81	3.1	1
28	Lipopolysaccharide-induced fever depends on prostaglandin E2 production specifically in brain endothelial cells. <i>Endocrinology</i> , 2012 , 153, 4849-61	4.8	71
27	The Q705K polymorphism in NLRP3 is a gain-of-function alteration leading to excessive interleukin-1 β and IL-18 production. <i>PLoS ONE</i> , 2012 , 7, e34977	3.7	109
26	The pan-ErbB tyrosine kinase inhibitor canertinib induces caspase-mediated cell death in human T-cell leukemia (Jurkat) cells. <i>Biochemical and Biophysical Research Communications</i> , 2011 , 410, 422-7	3.4	7

25	IL-7 mediates Ebf-1-dependent lineage restriction in early lymphoid progenitors. <i>Blood</i> , 2011 , 118, 1283-90		58
24	Irreversible pan-ERBB inhibitor canertinib elicits anti-leukaemic effects and induces the regression of FLT3-ITD transformed cells in mice. <i>British Journal of Haematology</i> , 2011 , 155, 198-208	4.5	7
23	Interleukin-7-induced Stat-5 acts in synergy with Flt-3 signaling to stimulate expansion of hematopoietic progenitor cells. <i>Journal of Biological Chemistry</i> , 2010 , 285, 36275-84	5.4	16
22	The pan-ErbB receptor tyrosine kinase inhibitor canertinib induces ErbB-independent apoptosis in human leukemia (HL-60 and U-937) cells. <i>Biochemical and Biophysical Research Communications</i> , 2010 , 393, 6-10	3.4	9
21	Hypoxia mediates low cell-cycle activity and increases the proportion of long-term-reconstituting hematopoietic stem cells during in vitro culture. <i>Experimental Hematology</i> , 2010 , 38, 301-310.e2	3.1	125
20	The hematopoietic stem cell niche: low in oxygen but a nice place to be. <i>Journal of Cellular Physiology</i> , 2010 , 222, 17-22	7	331
19	Deficiency of activating Fcγ receptors reduces hepatic clearance and deposition of IC and increases CIC levels in mercury-induced autoimmunity. <i>PLoS ONE</i> , 2010 , 5, e13413	3.7	8
18	Systemic reduction of functionally suppressive CD4 ^{dim} CD25 ^{high} Foxp3 ⁺ Tregs in human second trimester pregnancy is induced by progesterone and 17β-estradiol. <i>Journal of Immunology</i> , 2009 , 183, 759-69	5.3	116
17	BH3-only protein Bim more critical than Puma in tyrosine kinase inhibitor-induced apoptosis of human leukemic cells and transduced hematopoietic progenitors carrying oncogenic FLT3. <i>Blood</i> , 2009 , 113, 2302-11	2.2	29
16	Bcl11b mutations identified in murine lymphomas increase the proliferation rate of hematopoietic progenitor cells. <i>BMC Cancer</i> , 2007 , 7, 195	4.8	11
15	The Lim-only protein LMO2 acts as a positive regulator of erythroid differentiation. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 364, 675-81	3.4	13
14	The BH3-only protein Puma plays an essential role in cytokine deprivation induced apoptosis of mast cells. <i>Blood</i> , 2007 , 110, 3209-17	2.2	94
13	Absence of hot spot mutations of the PIK3CA gene in acute myeloid leukaemia. <i>European Journal of Haematology</i> , 2006 , 77, 86-7	3.8	11
12	Stem cell factor promotes mast cell survival via inactivation of FOXO3a-mediated transcriptional induction and MEK-regulated phosphorylation of the proapoptotic protein Bim. <i>Blood</i> , 2005 , 106, 1330-6	2.2	99
11	Characterization of the mouse myeloid-associated differentiation marker (MYADM) gene: promoter analysis and protein localization. <i>Molecular Biology Reports</i> , 2005 , 32, 149-57	2.8	3
10	FLT3 ligand regulates apoptosis through AKT-dependent inactivation of transcription factor FoxO3. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 318, 899-903	3.4	26
9	Inactivation of the forkhead transcription factor FoxO3 is essential for PKB-mediated survival of hematopoietic progenitor cells by kit ligand. <i>Experimental Hematology</i> , 2003 , 31, 316-23	3.1	37
8	The basic helix-loop-helix transcription factor TAL1/SCL inhibits the expression of the p16INK4A and pTalpha genes. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 312, 1073-81	3.4	19

7	The LIM-only protein LMO4 modulates the transcriptional activity of HEN1. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 307, 891-9	3-4	37
6	Phosphatidylinositol 3-kinase is essential for kit ligand-mediated survival, whereas interleukin-3 and flt3 ligand induce expression of antiapoptotic Bcl-2 family genes. <i>Journal of Leukocyte Biology</i> , 2003 , 74, 923-31	6.5	23
5	Distinct and regulated expression of Notch receptors in hematopoietic lineages and during myeloid differentiation. <i>European Journal of Immunology</i> , 2001 , 31, 3240-7	6.1	48
4	Isolation of MYADM, a novel hematopoietic-associated marker gene expressed in multipotent progenitor cells and up-regulated during myeloid differentiation. <i>Journal of Leukocyte Biology</i> , 2000 , 67, 423-31	6.5	21
3	Characterization, chromosomal localization, and expression during hematopoietic differentiation of the gene encoding Arl6ip, ADP-ribosylation-like factor-6 interacting protein (ARL6). <i>Genomics</i> , 2000 , 68, 351-4	4.3	25
2	Concentration-dependent effects of hematopoietic growth factors during in vitro expansion of mouse stem cells and progenitor cells. <i>Growth Factors</i> , 1997 , 14, 59-66	1.6	3
1	Interleukin-7 responsiveness of B220+ B cell precursors from bone marrow decreases in aging mice. <i>Cellular Immunology</i> , 1993 , 147, 267-78	4.4	18