

Aline Meirhaeghe

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/949079/aline-meirhaeghe-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

116
papers

13,376
citations

36
h-index

115
g-index

118
ext. papers

16,257
ext. citations

7.5
avg, IF

7.2
L-index

#	Paper	IF	Citations
116	Comparison of clinical profiles and care for patients with incident versus recurrent acute coronary syndromes in France: Data from the MONICA registries.. <i>PLoS ONE</i> , 2022 , 17, e0263589	3.7	
115	Interplay of physical activity and genetic variants of the endothelial lipase on cardiovascular disease risk factors. <i>Pediatric Research</i> , 2021 ,	3.2	1
114	Large disparities in 28-day case fatality by stroke subtype: data from a French stroke registry between 2008 and 2017. <i>European Journal of Neurology</i> , 2021 , 28, 2208-2217	6	1
113	The relationship between neighbourhood walkability and cardiovascular risk factors in northern France. <i>Science of the Total Environment</i> , 2021 , 772, 144877	10.2	1
112	Development of a Genetic Risk Score to predict the risk of overweight and obesity in European adolescents from the HELENA study. <i>Scientific Reports</i> , 2021 , 11, 3067	4.9	3
111	Mendelian randomization analysis does not support causal associations of birth weight with hypertension risk and blood pressure in adulthood. <i>European Journal of Epidemiology</i> , 2020 , 35, 685-697	12.1	2
110	Coronary heart disease incidence still decreased between 2006 and 2014 in France, except in young age groups: Results from the French MONICA registries. <i>European Journal of Preventive Cardiology</i> , 2020 , 27, 1178-1186	3.9	6
109	Association between lipoprotein lipase gene polymorphisms and cardiovascular disease risk factors in European adolescents: The Healthy Lifestyle in Europe by Nutrition in Adolescence study. <i>Pediatric Diabetes</i> , 2020 , 21, 747-757	3.6	1
108	Interplay between the Mediterranean diet and C-reactive protein genetic polymorphisms towards inflammation in adolescents. <i>Clinical Nutrition</i> , 2020 , 39, 1919-1926	5.9	7
107	PCSK9 post-transcriptional regulation: Role of a 3'RUTR microRNA-binding site variant in linkage disequilibrium with c.1420G. <i>Atherosclerosis</i> , 2020 , 314, 63-70	3.1	4
106	Interaction Effect of the Mediterranean Diet and an Obesity Genetic Risk Score on Adiposity and Metabolic Syndrome in Adolescents: The HELENA Study. <i>Nutrients</i> , 2020 , 12,	6.7	2
105	Association of Birth Weight With Type 2 Diabetes and Glycemic Traits: A Mendelian Randomization Study. <i>JAMA Network Open</i> , 2019 , 2, e1910915	10.4	14
104	Association between UCP1, UCP2, and UCP3 gene polymorphisms with markers of adiposity in European adolescents: The HELENA study. <i>Pediatric Obesity</i> , 2019 , 14, e12504	4.6	5
103	Sex Differences in Stroke Attack, Incidence, and Mortality Rates in Northern France. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2018 , 27, 1368-1374	2.8	11
102	Contributions of mean and shape of blood pressure distribution to worldwide trends and variations in raised blood pressure: a pooled analysis of 1018 population-based measurement studies with 88.6 million participants. <i>International Journal of Epidemiology</i> , 2018 , 47, 872-883i	7.8	40
101	Associations between REV-ERB α sleep duration and body mass index in European adolescents. <i>Sleep Medicine</i> , 2018 , 46, 56-60	4.6	7
100	Dietary linoleic acid interacts with FADS1 genetic variability to modulate HDL-cholesterol and obesity-related traits. <i>Clinical Nutrition</i> , 2018 , 37, 1683-1689	5.9	10

99	Associations between long-term exposure to air pollution, glycosylated hemoglobin, fasting blood glucose and diabetes mellitus in northern France. <i>Environment International</i> , 2018 , 120, 121-129	12.9	33
98	Comparison of the rates of stroke and acute coronary events in northern France. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 1534-1542	3.9	3
97	Low-grade systemic inflammation: a partial mediator of the relationship between diabetes and lung function. <i>Annals of Epidemiology</i> , 2018 , 28, 26-32	6.4	12
96	Identification of a functional FADS1 3R/7R variant associated with erythrocyte n-6 polyunsaturated fatty acids levels. <i>Journal of Clinical Lipidology</i> , 2018 , 12, 1280-1289	4.9	5
95	Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.6 million children, adolescents, and adults. <i>Lancet, The</i> , 2017 , 390, 2627-2642	40	2980
94	Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. <i>Lancet, The</i> , 2017 , 389, 37-55	40	1100
93	Dietary fat intake modifies the influence of the FTO rs9939609 polymorphism on adiposity in adolescents: The HELENA cross-sectional study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 937-43	4.5	14
92	Multiple microRNA regulation of lipoprotein lipase gene abolished by 3R/7R polymorphisms in a triglyceride-lowering haplotype harboring p.Ser474Ter. <i>Atherosclerosis</i> , 2016 , 246, 280-6	3.1	19
91	How obesity relates to socio-economic status: identification of eating behavior mediators. <i>International Journal of Obesity</i> , 2016 , 40, 1794-1801	5.5	31
90	Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. <i>Lancet, The</i> , 2016 , 387, 1513-1530	40	2039
89	Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. <i>Lancet, The</i> , 2016 , 387, 1377-1396	40	2787
88	Age- and sex-specific causal effects of adiposity on cardiovascular risk factors. <i>Diabetes</i> , 2015 , 64, 1841-52	5.9	50
87	Effects of diabetes definition on global surveillance of diabetes prevalence and diagnosis: a pooled analysis of 96 population-based studies with 331,288 participants. <i>Lancet Diabetes and Endocrinology, the</i> , 2015 , 3, 624-37	18.1	109
86	Is the adiposity-associated FTO gene variant related to all-cause mortality independent of adiposity? Meta-analysis of data from 169,551 Caucasian adults. <i>Obesity Reviews</i> , 2015 , 16, 327-340	10.6	7
85	Examination of the brain natriuretic peptide rs198389 single-nucleotide polymorphism on type 2 diabetes mellitus and related phenotypes in an Algerian population. <i>Gene</i> , 2015 , 567, 159-63	3.8	3
84	The n-3 long-chain PUFAs modulate the impact of the GCKR Pro446Leu polymorphism on triglycerides in adolescents. <i>Journal of Lipid Research</i> , 2015 , 56, 1774-80	6.3	9
83	Effects of established blood pressure loci on blood pressure values and hypertension risk in an Algerian population sample. <i>Journal of Human Hypertension</i> , 2015 , 29, 296-302	2.6	5
82	Cardiorespiratory fitness and ideal cardiovascular health in European adolescents. <i>Heart</i> , 2015 , 101, 766-73	5.3	61

81	Associations of early life and sociodemographic factors with menarcheal age in European adolescents. <i>European Journal of Pediatrics</i> , 2015 , 174, 271-8	4.1	4
80	Associations of common SNPs in the SORT1, GCKR, LPL, APOA1, CETP, LDLR, APOE genes with lipid trait levels in an Algerian population sample. <i>International Journal of Clinical and Experimental Pathology</i> , 2015 , 8, 7358-63	1.4	6
79	Associations of genetic variants in/near body mass index-associated genes with type 2 diabetes: a systematic meta-analysis. <i>Clinical Endocrinology</i> , 2014 , 81, 702-10	3.4	27
78	Effects of established BMI-associated loci on obesity-related traits in a French representative population sample. <i>BMC Genetics</i> , 2014 , 15, 62	2.6	18
77	Physical activity modifies the associations between genetic variants and blood pressure in European adolescents. <i>Journal of Pediatrics</i> , 2014 , 165, 1046-9.e1-2	3.6	3
76	Combined effect of established BMI loci on obesity-related traits in an Algerian population sample. <i>BMC Genetics</i> , 2014 , 15, 128	2.6	5
75	Hypomethylation of the promoter of the catalytic subunit of protein phosphatase 2A in response to hyperglycemia. <i>Physiological Reports</i> , 2014 , 2, e12076	2.6	4
74	The TCF7L2 rs7903146 polymorphism, dietary intakes and type 2 diabetes risk in an Algerian population. <i>BMC Genetics</i> , 2014 , 15, 134	2.6	17
73	Impact of REV-ERB alpha gene polymorphisms on obesity phenotypes in adult and adolescent samples. <i>International Journal of Obesity</i> , 2013 , 37, 666-72	5.5	34
72	Thyroid hormone receptor alpha gene variants increase the risk of developing obesity and show gene-diet interactions. <i>International Journal of Obesity</i> , 2013 , 37, 1499-505	5.5	12
71	Impact of APOE gene polymorphisms on the lipid profile in an Algerian population. <i>Lipids in Health and Disease</i> , 2013 , 12, 155	4.4	24
70	CD36 and SR-BI are involved in cellular uptake of provitamin A carotenoids by Caco-2 and HEK cells, and some of their genetic variants are associated with plasma concentrations of these micronutrients in humans. <i>Journal of Nutrition</i> , 2013 , 143, 448-56	4.1	85
69	Genetic and molecular insights into the role of PROX1 in glucose metabolism. <i>Diabetes</i> , 2013 , 62, 1738-45.	9	29
68	New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. <i>Nature Genetics</i> , 2013 , 45, 76-82	36.3	232
67	Intake and dietary sources of haem and non-haem iron among European adolescents and their association with iron status and different lifestyle and socio-economic factors. <i>European Journal of Clinical Nutrition</i> , 2013 , 67, 765-72	5.2	20
66	A genome-wide association meta-analysis identifies new childhood obesity loci. <i>Nature Genetics</i> , 2012 , 44, 526-31	36.3	292
65	Intake and dietary sources of haem and non-haem iron in Flemish preschoolers. <i>European Journal of Clinical Nutrition</i> , 2012 , 66, 806-12	5.2	6
64	Body size at birth modifies the effect of fat mass and obesity associated (FTO) rs9939609 polymorphism on adiposity in adolescents: the Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) study. <i>British Journal of Nutrition</i> , 2012 , 107, 1498-504	3.6	11

63	Study of thyroid hormone receptor alpha gene polymorphisms on Alzheimer's disease. <i>Neurobiology of Aging</i> , 2011 , 32, 624-30	5.6	12
62	Associations between common genetic polymorphisms in the liver X receptor alpha and its target genes with the serum HDL-cholesterol concentration in adolescents of the HELENA Study. <i>Atherosclerosis</i> , 2011 , 216, 166-9	3.1	16
61	The effect of ponderal index at birth on the relationships between common LEP and LEPR polymorphisms and adiposity in adolescents. <i>Obesity</i> , 2011 , 19, 2038-45	8	16
60	Association between the FTO rs9939609 polymorphism and leptin in European adolescents: a possible link with energy balance control. The HELENA study. <i>International Journal of Obesity</i> , 2011 , 35, 66-71	5.5	35
59	Common polymorphisms in six genes of the methyl group metabolism pathway and obesity in European adolescents. <i>Pediatric Obesity</i> , 2011 , 6, e336-44		8
58	Association between a thyroid hormone receptor- β gene polymorphism and blood pressure but not with coronary heart disease risk. <i>American Journal of Hypertension</i> , 2011 , 24, 1027-34	2.3	9
57	FADS1 genetic variability interacts with dietary linolenic acid intake to affect serum non-HDL-cholesterol concentrations in European adolescents. <i>Journal of Nutrition</i> , 2011 , 141, 1247-53	4.1	35
56	Study of estrogen receptor- α and receptor- β gene polymorphisms on Alzheimer's disease. <i>Journal of Alzheimer's Disease</i> , 2011 , 26, 431-9	4.3	15
55	Polymorphisms in the CD36/FAT gene are associated with plasma vitamin E concentrations in humans. <i>American Journal of Clinical Nutrition</i> , 2011 , 93, 644-51	7	36
54	Physical activity attenuates the effect of low birth weight on insulin resistance in adolescents: findings from two observational studies. <i>Diabetes</i> , 2011 , 60, 2295-9	0.9	25
53	Mendelian randomization study of B-type natriuretic peptide and type 2 diabetes: evidence of causal association from population studies. <i>PLoS Medicine</i> , 2011 , 8, e1001112	11.6	79
52	Physical activity attenuates the influence of FTO variants on obesity risk: a meta-analysis of 218,166 adults and 19,268 children. <i>PLoS Medicine</i> , 2011 , 8, e1001116	11.6	379
51	Single-nucleotide polymorphism of CD36 locus and obesity in European adolescents. <i>Obesity</i> , 2010 , 18, 1398-403	8	47
50	Influence of maternal educational level on the association between the rs3809508 neuromedin B gene polymorphism and the risk of obesity in the HELENA study. <i>International Journal of Obesity</i> , 2010 , 34, 478-86	5.5	14
49	Attenuation of the effect of the FTO rs9939609 polymorphism on total and central body fat by physical activity in adolescents: the HELENA study. <i>JAMA Pediatrics</i> , 2010 , 164, 328-33		85
48	Single nucleotide polymorphisms in the FADS gene cluster are associated with delta-5 and delta-6 desaturase activities estimated by serum fatty acid ratios. <i>Journal of Lipid Research</i> , 2010 , 51, 2325-33	6.3	134
47	Breast-feeding modulates the influence of the peroxisome proliferator-activated receptor-gamma (PPARG2) Pro12Ala polymorphism on adiposity in adolescents: The Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) cross-sectional study. <i>Diabetes Care</i> , 2010 , 33, 190-6	14.6	17
46	Concordance of two multiple analytical approaches demonstrate that interaction between BMI and ADIPOQ haplotypes is a determinant of LDL cholesterol in a general French population. <i>Journal of Human Genetics</i> , 2010 , 55, 227-31	4.3	3

45	No association between polymorphisms in the INSIG1 gene and the risk of type 2 diabetes and related traits. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 252-7	7	8
44	Triglyceride-mediated pathways and coronary heart disease. <i>Lancet, The</i> , 2010 , 376, 956-7; author reply 957-8	4.0	4
43	Suggestive evidence of associations between liver X receptor polymorphisms with type 2 diabetes mellitus and obesity in three cohort studies: HUNT2 (Norway), MONICA (France) and HELENA (Europe). <i>BMC Medical Genetics</i> , 2010 , 11, 144	2.1	23
42	Peroxisome proliferator-activated receptor gamma polymorphisms and coronary heart disease. <i>PPAR Research</i> , 2009 , 2009, 543746	4.3	21
41	Early life programming of abdominal adiposity in adolescents: The HELENA Study. <i>Diabetes Care</i> , 2009 , 32, 2120-2	14.6	41
40	Associations between common genetic polymorphisms in angiotensin-like proteins 3 and 4 and lipid metabolism and adiposity in European adolescents and adults. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 5070-7	5.6	28
39	Effect of an FTO polymorphism on fat mass, obesity, and type 2 diabetes mellitus in the French MONICA Study. <i>Metabolism: Clinical and Experimental</i> , 2009 , 58, 971-5	12.7	56
38	Association between angiotensin-like 6 (ANGPTL6) gene polymorphisms and metabolic syndrome-related phenotypes in the French MONICA Study. <i>Diabetes and Metabolism</i> , 2009 , 35, 287-92	5.4	8
37	In obese and non-obese adults, the cis-regulatory rs361072 promoter variant of PIK3CB is associated with insulin resistance not with type 2 diabetes. <i>Molecular Genetics and Metabolism</i> , 2009 , 96, 129-32	3.7	8
36	Study of the genetic variability of ZAC1 (PLAGL1) in French population-based samples. <i>Journal of Hypertension</i> , 2009 , 27, 314-21	1.9	5
35	Association between liver X receptor alpha gene polymorphisms and risk of metabolic syndrome in French populations. <i>International Journal of Obesity</i> , 2008 , 32, 421-8	5.5	27
34	The APOA5 Trp19 allele is associated with metabolic syndrome via its association with plasma triglycerides. <i>BMC Medical Genetics</i> , 2008 , 9, 84	2.1	24
33	Association between the T-381C polymorphism of the brain natriuretic peptide gene and risk of type 2 diabetes in human populations. <i>Human Molecular Genetics</i> , 2007 , 16, 1343-50	5.6	64
32	A possible role for the PPARG Pro12Ala polymorphism in preterm birth. <i>Diabetes</i> , 2007 , 56, 494-8	0.9	34
31	Ablation of PGC-1beta results in defective mitochondrial activity, thermogenesis, hepatic function, and cardiac performance. <i>PLoS Biology</i> , 2006 , 4, e369	9.7	217
30	A study of the relationships between KLF2 polymorphisms and body weight control in a French population. <i>BMC Medical Genetics</i> , 2006 , 7, 26	2.1	2
29	Study of the impact of perilipin polymorphisms in a French population. <i>Journal of Negative Results in BioMedicine</i> , 2006 , 5, 10		13
28	Study of a new PPARgamma2 promoter polymorphism and haplotype analysis in a French population. <i>Molecular Genetics and Metabolism</i> , 2005 , 85, 140-8	3.7	22

27	Lack of association between certain candidate gene polymorphisms and the metabolic syndrome. <i>Molecular Genetics and Metabolism</i> , 2005 , 86, 293-9	3.7	39
26	The APOA4 Thr347->Ser347 polymorphism is not a major risk factor of obesity. <i>Obesity</i> , 2005 , 13, 2132-8		11
25	A dominant negative human peroxisome proliferator-activated receptor (PPAR){alpha} is a constitutive transcriptional corepressor and inhibits signaling through all PPAR isoforms. <i>Endocrinology</i> , 2005 , 146, 1871-82	4.8	38
24	Association between peroxisome proliferator-activated receptor gamma haplotypes and the metabolic syndrome in French men and women. <i>Diabetes</i> , 2005 , 54, 3043-8	0.9	68
23	Genetic variants in human sterol regulatory element binding protein-1c in syndromes of severe insulin resistance and type 2 diabetes. <i>Diabetes</i> , 2004 , 53, 842-6	0.9	46
22	Impact of genetic variation of PPARgamma in humans. <i>Molecular Genetics and Metabolism</i> , 2004 , 83, 93-102	3.7	119
21	The Gly16-->Arg16 and Gln27-->Glu27 polymorphisms of beta2-adrenergic receptor are associated with metabolic syndrome in men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003 , 88, 4862-6	5.6	63
20	Characterization of the human, mouse and rat PGC1 beta (peroxisome-proliferator-activated receptor-gamma co-activator 1 beta) gene in vitro and in vivo. <i>Biochemical Journal</i> , 2003 , 373, 155-65	3.8	167
19	A functional polymorphism in a STAT5B site of the human PPAR gamma 3 gene promoter affects height and lipid metabolism in a French population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2003 , 23, 289-94	9.4	85
18	Digenic inheritance of severe insulin resistance in a human pedigree. <i>Nature Genetics</i> , 2002 , 31, 379-84	36.3	121
17	The 5A6A polymorphism in the promoter of the stromelysin-1 (MMP3) gene as a risk factor for restenosis. <i>European Heart Journal</i> , 2002 , 23, 721-5	9.5	48
16	Impact of sulfonylurea receptor 1 genetic variability on non-insulin-dependent diabetes mellitus prevalence and treatment: a population study. <i>American Journal of Medical Genetics Part A</i> , 2001 , 101, 4-8		36
15	Polymorphisms in the insulin response element of APOC-III gene promoter influence the correlation between insulin and triglycerides or triglyceride-rich lipoproteins in humans. <i>International Journal of Obesity</i> , 2001 , 25, 1012-7	5.5	11
14	The human G-protein beta3 subunit C825T polymorphism is associated with coronary artery vasoconstriction. <i>European Heart Journal</i> , 2001 , 22, 845-8	9.5	28
13	The effect of the Gly16Arg polymorphism of the beta(2)-adrenergic receptor gene on plasma free fatty acid levels is modulated by physical activity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 5881-7	5.6	30
12	Polymorphisms of the tissue factor pathway inhibitor gene and the risk of restenosis after coronary angioplasty. <i>Blood Coagulation and Fibrinolysis</i> , 2001 , 12, 317-23	1	23
11	The Effect of the Gly16Arg Polymorphism of the β -Adrenergic Receptor Gene on Plasma Free Fatty Acid Levels Is Modulated by Physical Activity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 5881-5887	5.6	24
10	Impact of polymorphisms of the human beta2-adrenoceptor gene on obesity in a French population. <i>International Journal of Obesity</i> , 2000 , 24, 382-7	5.5	69

9	Impact of the Peroxisome Proliferator Activated Receptor gamma2 Pro12Ala polymorphism on adiposity, lipids and non-insulin-dependent diabetes mellitus. <i>International Journal of Obesity</i> , 2000 , 24, 195-9	5.5	131
8	An uncoupling protein 3 gene polymorphism associated with a lower risk of developing Type II diabetes and with atherogenic lipid profile in a French cohort. <i>Diabetologia</i> , 2000 , 43, 1424-8	10.3	70
7	Intronic polymorphism in the fatty acid transport protein 1 gene is associated with increased plasma triglyceride levels in a French population. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000 , 20, 1330-4	9.4	32
6	Gender related association between genetic variations of APOC-III gene and lipid and lipoprotein variables in northern France. <i>Atherosclerosis</i> , 2000 , 150, 149-57	3.1	35
5	Beta2-adrenoceptor gene polymorphism, body weight, and physical activity. <i>Lancet, The</i> , 1999 , 353, 896-900	4.0	121
4	Paraoxonase (Gln192-Arg) polymorphism in French type 2 diabetics. <i>Atherosclerosis</i> , 1999 , 147, 415-6	3.1	8
3	A genetic polymorphism of the peroxisome proliferator-activated receptor gamma gene influences plasma leptin levels in obese humans. <i>Human Molecular Genetics</i> , 1998 , 7, 435-40	5.6	164
2	Role of Lipid Binding Proteins in Disease397-400		
1	ACE inhibitors, AT1 receptor blockers and COVID-19: clinical epidemiology evidences for a continuation of treatments. The ACER-COVID study		8