

# Kobi Rosenblum

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9492255/publications.pdf>

Version: 2024-02-01

68  
papers

5,056  
citations

117571

34  
h-index

98753

67  
g-index

77  
all docs

77  
docs citations

77  
times ranked

5228  
citing authors

#	ARTICLE	IF	CITATIONS
1	Brain-Derived Neurotrophic Factor Induces Long-Term Potentiation in Intact Adult Hippocampus: Requirement for ERK Activation Coupled to CREB and Upregulation of <i>Arc</i> Synthesis. <i>Journal of Neuroscience</i> , 2002, 22, 1532-1540.	1.7	699
2	eIF2 $\pm$ Phosphorylation Bidirectionally Regulates the Switch from Short- to Long-Term Synaptic Plasticity and Memory. <i>Cell</i> , 2007, 129, 195-206.	13.5	437
3	Glycogen synthase kinase-3 inhibition is integral to long-term potentiation. <i>European Journal of Neuroscience</i> , 2007, 25, 81-86.	1.2	300
4	Taste memory: The role of protein synthesis in gustatory cortex. <i>Behavioral and Neural Biology</i> , 1993, 59, 49-56.	2.3	279
5	Specific and Differential Activation of Mitogen-Activated Protein Kinase Cascades by Unfamiliar Taste in the Insular Cortex of the Behaving Rat. <i>Journal of Neuroscience</i> , 1998, 18, 10037-10044.	1.7	276
6	NMDA Receptor and the Tyrosine Phosphorylation of Its 2B Subunit in Taste Learning in the Rat Insular Cortex. <i>Journal of Neuroscience</i> , 1997, 17, 5129-5135.	1.7	217
7	PKR: A Kinase to Remember. <i>Frontiers in Molecular Neuroscience</i> , 2018, 11, 480.	1.4	172
8	ERK1/II Regulation by the Muscarinic Acetylcholine Receptors in Neurons. <i>Journal of Neuroscience</i> , 2000, 20, 977-985.	1.7	161
9	The Role of Extracellular Regulated Kinases I/II in Late-Phase Long-Term Potentiation. <i>Journal of Neuroscience</i> , 2002, 22, 5432-5441.	1.7	144
10	Removal of S6K1 and S6K2 leads to divergent alterations in learning, memory, and synaptic plasticity. <i>Learning and Memory</i> , 2008, 15, 29-38.	0.5	132
11	The roles of protein expression in synaptic plasticity and memory consolidation. <i>Frontiers in Molecular Neuroscience</i> , 2014, 7, 86.	1.4	125
12	Insular cortex neurons encode and retrieve specific immune responses. <i>Cell</i> , 2021, 184, 5902-5915.e17.	13.5	124
13	Molecular Mechanisms of Long-Term Potentiation in the Insular Cortex <i>In Vivo</i> . <i>Journal of Neuroscience</i> , 1999, 19, RC36-RC36.	1.7	103
14	The role of eEF2 pathway in learning and synaptic plasticity. <i>Neurobiology of Learning and Memory</i> , 2013, 105, 100-106.	1.0	94
15	Consolidation and translation regulation: Figure 1.. <i>Learning and Memory</i> , 2012, 19, 410-422.	0.5	77
16	Local Inhibition of PERK Enhances Memory and Reverses Age-Related Deterioration of Cognitive and Neuronal Properties. <i>Journal of Neuroscience</i> , 2018, 38, 648-658.	1.7	74
17	eIF2 $\pm$ controls memory consolidation via excitatory and somatostatin neurons. <i>Nature</i> , 2020, 586, 412-416.	13.7	74
18	Molecular Mechanisms Underlying Memory Consolidation of Taste Information in the Cortex. <i>Frontiers in Behavioral Neuroscience</i> , 2011, 5, 87.	1.0	68

#	ARTICLE	IF	CITATIONS
19	Blocking the eIF2 $\pm$ Kinase (PKR) Enhances Positive and Negative Forms of Cortex-Dependent Taste Memory. <i>Journal of Neuroscience</i> , 2013, 33, 2517-2525.	1.7	68
20	Calcium/Calmodulin-Dependent Protein Kinase II and Eukaryotic Elongation Factor 2 Kinase Pathways Mediate the Antidepressant Action of Ketamine. <i>Biological Psychiatry</i> , 2018, 84, 65-75.	0.7	68
21	ERK-dependent PSD-95 induction in the gustatory cortex is necessary for taste learning, but not retrieval. <i>Nature Neuroscience</i> , 2008, 11, 1149-1151.	7.1	66
22	Encoding of Conditioned Taste Aversion in Cortico-Amygdala Circuits. <i>Cell Reports</i> , 2018, 24, 278-283.	2.9	66
23	Biphasic Activation of the mTOR Pathway in the Gustatory Cortex Is Correlated with and Necessary for Taste Learning. <i>Journal of Neuroscience</i> , 2009, 29, 7424-7431.	1.7	65
24	Impaired associative taste learning and abnormal brain activation in kinase-defective eEF2K mice. <i>Learning and Memory</i> , 2012, 19, 116-125.	0.5	61
25	ApoE $\mu$ 4 is associated with eIF2 $\pm$ phosphorylation and impaired learning in young mice. <i>Neurobiology of Aging</i> , 2013, 34, 863-872.	1.5	61
26	Genetic or Pharmacological Reduction of PERK Enhances Cortical-Dependent Taste Learning. <i>Journal of Neuroscience</i> , 2014, 34, 14624-14632.	1.7	57
27	eEF2K/eEF2 Pathway Controls the Excitation/Inhibition Balance and Susceptibility to Epileptic Seizures. <i>Cerebral Cortex</i> , 2017, 27, bhw075.	1.6	57
28	A Novel Role for Extracellular Signal-Regulated Kinase in Maintaining Long-Term Memory-Relevant Excitability Changes. <i>Journal of Neuroscience</i> , 2007, 27, 12584-12589.	1.7	55
29	Activity of Insula to Basolateral Amygdala Projecting Neurons is Necessary and Sufficient for Taste Valence Representation. <i>Journal of Neuroscience</i> , 2019, 39, 9369-9382.	1.7	55
30	Facilitation of taste memory acquisition by experiencing previous novel taste is protein-synthesis dependent. <i>Learning and Memory</i> , 2008, 15, 501-507.	0.5	52
31	PKR Inhibition Rescues Memory Deficit and ATF4 Overexpression in ApoE $\mu$ 4 Human Replacement Mice. <i>Journal of Neuroscience</i> , 2015, 35, 12986-12993.	1.7	51
32	The Insula and Taste Learning. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 335.	1.4	51
33	Different signal transduction cascades are activated simultaneously in the rat insular cortex and hippocampus following novel taste learning. <i>European Journal of Neuroscience</i> , 2006, 24, 1434-1442.	1.2	47
34	Tyrosine Phosphorylation of the 2B Subunit of the NMDA Receptor Is Necessary for Taste Memory Formation. <i>Journal of Neuroscience</i> , 2009, 29, 9219-9226.	1.7	45
35	Multi-input Synapses, but Not LTP-Strengthened Synapses, Correlate with Hippocampal Memory Storage in Aged Mice. <i>Current Biology</i> , 2019, 29, 3600-3610.e4.	1.8	39
36	A Novel Role for Protein Synthesis in Long-Term Neuronal Plasticity: Maintaining Reduced Postburst Afterhyperpolarization. <i>Journal of Neuroscience</i> , 2010, 30, 4338-4342.	1.7	30

#	ARTICLE	IF	CITATIONS
37	Differential Contribution of Hippocampal Subfields to Components of Associative Taste Learning. <i>Journal of Neuroscience</i> , 2014, 34, 11007-11015.	1.7	30
38	A molecular mechanism underlying gustatory memory trace for an association in the insular cortex. <i>ELife</i> , 2015, 4, e07582.	2.8	29
39	Behavioral interference and C/EBP $\alpha$ expression in the insular-cortex reveal a prolonged time period for taste memory consolidation. <i>Learning and Memory</i> , 2006, 13, 571-574.	0.5	27
40	NMDA and Dopamine Converge on the NMDA-Receptor to Induce ERK Activation and Synaptic Depression in Mature Hippocampus. <i>PLoS ONE</i> , 2006, 1, e138.	1.1	27
41	Taste Familiarity Is Inversely Correlated with Arc/Arg3.1 Hemispheric Lateralization. <i>Journal of Neuroscience</i> , 2013, 33, 11734-11743.	1.7	25
42	Expression of Quinone Reductase-2 in the Cortex Is a Muscarinic Acetylcholine Receptor-Dependent Memory Consolidation Constraint. <i>Journal of Neuroscience</i> , 2015, 35, 15568-15581.	1.7	25
43	Memory of Conditioned Taste Aversion Is Erased by Inhibition of PI3K in the Insular Cortex. <i>Neuropsychopharmacology</i> , 2013, 38, 1143-1153.	2.8	24
44	The differential role of cortical protein synthesis in taste memory formation and persistence. <i>Npj Science of Learning</i> , 2016, 1, 16001.	1.5	21
45	eEF2/eEF2K Pathway in the Mature Dentate Gyrus Determines Neurogenesis Level and Cognition. <i>Current Biology</i> , 2020, 30, 3507-3521.e7.	1.8	21
46	The Role of Protein Phosphorylation in the Gustatory Cortex and Amygdala During Taste Learning. <i>Experimental Neurobiology</i> , 2012, 21, 37-51.	0.7	20
47	NMDAR-dependent proteasome activity in the gustatory cortex is necessary for conditioned taste aversion. <i>Neurobiology of Learning and Memory</i> , 2016, 130, 7-16.	1.0	19
48	D1 Dopamine Receptor Activation Induces Neuronal eEF2 Pathway-Dependent Protein Synthesis. <i>Frontiers in Molecular Neuroscience</i> , 2020, 13, 67.	1.4	19
49	Dopamine-induced tyrosine phosphorylation of NR2B (Tyr1472) is essential for ERK1/2 activation and processing of novel taste information. <i>Frontiers in Molecular Neuroscience</i> , 2014, 7, 66.	1.4	18
50	Concurrence of High Fat Diet and APOE Gene Induces Allele Specific Metabolic and Mental Stress Changes in a Mouse Model of Alzheimer's Disease. <i>Frontiers in Behavioral Neuroscience</i> , 2016, 10, 170.	1.0	17
51	mAChR-dependent decrease in proteasome activity in the gustatory cortex is necessary for novel taste learning. <i>Neurobiology of Learning and Memory</i> , 2016, 135, 115-124.	1.0	17
52	Virally mediated gene manipulation in the adult CNS. <i>Frontiers in Molecular Neuroscience</i> , 2011, 4, 57.	1.4	16
53	MAPK activation in the hippocampus in vivo is correlated with experimental setting. <i>Neurobiology of Learning and Memory</i> , 2007, 88, 58-64.	1.0	15
54	Olfactory learning abilities are correlated with the rate by which intrinsic neuronal excitability is modulated in the piriform cortex. <i>European Journal of Neuroscience</i> , 2009, 30, 1339-1348.	1.2	15

#	ARTICLE	IF	CITATIONS
55	Measuring mRNA translation in neuronal processes and somata by tRNA-FRET. <i>Nucleic Acids Research</i> , 2020, 48, e32-e32.	6.5	15
56	Parvalbumin interneuron inhibition onto anterior insula neurons projecting to the basolateral amygdala drives aversive taste memory retrieval. <i>Current Biology</i> , 2021, 31, 2770-2784.e6.	1.8	15
57	Persistent ERK activation maintains learning-induced long-lasting modulation of synaptic connectivity. <i>Learning and Memory</i> , 2008, 15, 756-761.	0.5	12
58	Fluid consumption and taste novelty determines transcription temporal dynamics in the gustatory cortex. <i>Molecular Brain</i> , 2016, 9, 13.	1.3	12
59	Insula to mPFC reciprocal connectivity differentially underlies novel taste neophobic response and learning in mice. <i>ELife</i> , 2021, 10, .	2.8	12
60	4E-BP2-dependent translation in parvalbumin neurons controls epileptic seizure threshold. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	3.3	10
61	Design and synthesis of novel protein kinase R (PKR) inhibitors. <i>Molecular Diversity</i> , 2016, 20, 805-819.	2.1	8
62	Trace Fear Conditioning: Procedure for Assessing Complex Hippocampal Function in Mice. <i>Bio-protocol</i> , 2018, 8, e2475.	0.2	8
63	Dopamine-Dependent QR2 Pathway Activation in CA1 Interneurons Enhances Novel Memory Formation. <i>Journal of Neuroscience</i> , 2020, 40, 8698-8714.	1.7	7
64	Muscarinic-Dependent miR-182 and QR2 Expression Regulation in the Anterior Insula Enables Novel Taste Learning. <i>ENeuro</i> , 2020, 7, ENEURO.0067-20.2020.	0.9	6
65	Somatostatin Interneurons of the Insula Mediate QR2-Dependent Novel Taste Memory Enhancement. <i>ENeuro</i> , 2021, 8, ENEURO.0152-21.2021.	0.9	5
66	Rho-associated kinase in the gustatory cortex is involved in conditioned taste aversion memory formation but not in memory retrieval or relearning. <i>Neurobiology of Learning and Memory</i> , 2012, 97, 1-6.	1.0	4
67	Editorial. <i>Neurobiology of Learning and Memory</i> , 2015, 124, 1-2.	1.0	1
68	The Role of the Eukaryotic Elongation Factor 2 (eEF2) Pathway in Neuronal Function. , 0, , 63-80.		0