

Tanja Schmitz-HÃ¼bsch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9495361/publications.pdf>

Version: 2024-02-01

16
papers

1,020
citations

933447

10
h-index

940533

16
g-index

19
all docs

19
docs citations

19
times ranked

1923
citing authors

#	ARTICLE	IF	CITATIONS
1	Proposal for Post Hoc Quality Control in Instrumented Motion Analysis Using Markerless Motion Capture: Development and Usability Study. <i>JMIR Human Factors</i> , 2022, 9, e26825.	2.0	2
2	Cultural bias in motor function patterns: Potential relevance for predictive, preventive, and personalized medicine. <i>EPMA Journal</i> , 2021, 12, 91-101.	6.1	4
3	Spinocerebellar ataxia type 14: refining clinicogenetic diagnosis in a rare adult-onset disorder. <i>Annals of Clinical and Translational Neurology</i> , 2021, 8, 774-789.	3.7	13
4	Neural Processes of Psychological Stress and Relaxation Predict the Future Evolution of Quality of Life in Multiple Sclerosis. <i>Frontiers in Neurology</i> , 2021, 12, 753107.	2.4	7
5	Instrumental Assessment of Stepping in Place Captures Clinically Relevant Motor Symptoms of Parkinson's Disease. <i>Sensors</i> , 2020, 20, 5465.	3.8	8
6	Investigation of Visual System Involvement in Spinocerebellar Ataxia Type 14. <i>Cerebellum</i> , 2020, 19, 469-482.	2.5	3
7	Association Between Fatigue and Motor Exertion in Patients With Multiple Sclerosis—a Prospective Study. <i>Frontiers in Neurology</i> , 2020, 11, 208.	2.4	18
8	Survival in patients with spinocerebellar ataxia types 1, 2, 3, and 6 (EUROSCA): a longitudinal cohort study. <i>Lancet Neurology</i> , The, 2018, 17, 327-334.	10.2	69
9	Less Is More — Estimation of the Number of Strides Required to Assess Gait Variability in Spatially Confined Settings. <i>Frontiers in Aging Neuroscience</i> , 2018, 10, 435.	3.4	41
10	Subjective and objective assessment of physical activity in multiple sclerosis and their relation to health-related quality of life. <i>BMC Neurology</i> , 2017, 17, 10.	1.8	18
11	Connectivity Predicts deep brain stimulation outcome in Parkinson disease. <i>Annals of Neurology</i> , 2017, 82, 67-78.	5.3	514
12	Maximum walking speed in multiple sclerosis assessed with visual perceptive computing. <i>PLoS ONE</i> , 2017, 12, e0189281.	2.5	29
13	Accuracy and repeatability of two methods of gait analysis — GaitRite, and Mobility Lab, in subjects with cerebellar ataxia. <i>Gait and Posture</i> , 2016, 48, 194-201.	1.4	59
14	Validity of visual perceptive computing for static posturography in patients with multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2016, 22, 1596-1606.	3.0	39
15	Accuracy and Reliability of the Kinect Version 2 for Clinical Measurement of Motor Function. <i>PLoS ONE</i> , 2016, 11, e0166532.	2.5	183
16	Cerebellar neurochemical alterations in spinocerebellar ataxia type 14 appear to include glutathione deficiency. <i>Journal of Neurology</i> , 2015, 262, 1927-1935.	3.6	13