
Mariana Resende

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9496190/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Myeloid HIFâ€1α regulates pulmonary inflammation during experimental Mycobacterium tuberculosis infection. Immunology, 2020, 159, 121-129. | 2.0 | 17 |
| 2 | <i>Leishmania infantum</i> Enhances Migration of Macrophages via a Phosphoinositide 3-Kinase Î ³ -Dependent Pathway. ACS Infectious Diseases, 2020, 6, 1643-1649. | 1.8 | 6 |
| 3 | TNF-Mediated Compensatory Immunity to <i>Mycobacterium avium</i> in the Absence of Macrophage Activation by IFN-γ. Journal of Immunology, 2019, 203, 2451-2458. | 0.4 | 5 |
| 4 | IFN-γ–Dependent Reduction of Erythrocyte Life Span Leads to Anemia during Mycobacterial Infection. Journal of Immunology, 2019, 203, 2485-2496. | 0.4 | 27 |
| 5 | Infection of hematopoietic stem cells by Leishmania infantum increases erythropoiesis and alters the phenotypic and functional profiles of progeny. Cellular Immunology, 2018, 326, 77-85. | 1.4 | 10 |
| 6 | Innate IFN-γ–Producing Cells Developing in the Absence of IL-2 Receptor Common γ-Chain. Journal of Immunology, 2017, 199, 1429-1439. | 0.4 | 9 |
| 7 | The Warburg effect in mycobacterial granulomas is dependent on the recruitment and activation of macrophages by interferonâ€ <i>î³</i> . Immunology, 2015, 145, 498-507. | 2.0 | 45 |
| 8 | Lack of the Transcription Factor Hypoxia-Inducible Factor 1α (HIF-1α) in Macrophages Accelerates the Necrosis of Mycobacterium avium-Induced Granulomas. Infection and Immunity, 2015, 83, 3534-3544. | 1.0 | 27 |
| 9 | <i>Leishmania</i> -Infected MHC Class Ilhigh Dendritic Cells Polarize CD4+ T Cells toward a Nonprotective T-bet+ IFN-γ+ IL-10+ Phenotype. Journal of Immunology, 2013, 191, 262-273. | 0.4 | 37 |
| 10 | Characterization and evaluation of BNIPDaoct-loaded PLGA nanoparticles for visceral leishmaniasis: <i>in vitro</i> and <i>in vivo</i> studies. Nanomedicine, 2012, 7, 1839-1849. | 1.7 | 35 |
| 11 | Rapamycin Combined with TGF-Î ² Converts Human Invariant NKT Cells into Suppressive Foxp3+ Regulatory Cells. Journal of Immunology, 2012, 188, 624-631. | 0.4 | 59 |
| 12 | Proinflammatory Environment Dictates the IL-17–Producing Capacity of Human Invariant NKT Cells. Journal of Immunology, 2011, 186, 5758-5765. | 0.4 | 90 |
| 13 | Activation of Phosphatidylinositol 3-Kinase/Akt and Impairment of Nuclear Factor-κB. American Journal of Pathology, 2010, 177, 2898-2911. | 1.9 | 48 |