

Wael Agur

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9496834/publications.pdf>

Version: 2024-02-01

40
papers

758
citations

567144

15
h-index

526166

27
g-index

43
all docs

43
docs citations

43
times ranked

794
citing authors

#	ARTICLE	IF	CITATIONS
1	The importance of overcoming the challenges in delivering the Proper Understanding of Recurrent Stress Urinary Incontinence Treatment (PURSUIT) study. , 2022, 1, 100020.		1
2	Treating Women with Recurrent Stress Urinary Incontinence: A Hornetâ€™s Nest Still Needing Proper Clinical Evidence. European Urology, 2021, 79, 6-7.	0.9	5
3	Vesico-vaginal fistula presenting as overactive bladder in a case of Gellhorn pessary for vault prolapse. BMJ Case Reports, 2021, 14, e240331.	0.2	1
4	Long-term efficacy and complications of a multicentre randomised controlled trial comparing Retropubic and Transobturator Mid-urethral Slings: a prospective observational study. BJOG: an International Journal of Obstetrics and Gynaecology, 2021, , .	1.1	1
5	Letter to Editor: â€œUtility of patient decision aids (PDA) in stress urinary incontinence surgeryâ€•by Jha and Duckett 2019. International Urogynecology Journal, 2020, 31, 419-419.	0.7	1
6	Effectiveness of pelvic floor muscle training with and without electromyographic biofeedback for urinary incontinence in women: multicentre randomised controlled trial. BMJ, The, 2020, 371, m3719.	3.0	50
7	The TOPSY pessary self-management intervention for pelvic organ prolapse: a study protocol for the process evaluation. Trials, 2020, 21, 836.	0.7	3
8	Clinical and cost-effectiveness of vaginal pessary self-management compared to clinic-based care for pelvic organ prolapse: protocol for the TOPSY randomised controlled trial. Trials, 2020, 21, 837.	0.7	7
9	Mesh inlay, mesh kit or native tissue repair for women having repeat anterior or posterior prolapse surgery: randomised controlled trial (PROSPECT). BJOG: an International Journal of Obstetrics and Gynaecology, 2020, 127, 1002-1013.	1.1	13
10	Basic versus biofeedback-mediated intensive pelvic floor muscle training for women with urinary incontinence: the OPAL RCT. Health Technology Assessment, 2020, 24, 1-144.	1.3	19
11	Development, validation and initial evaluation of patient-decision aid (SUI-PDAÂ©) for women considering stress urinary incontinence surgery. International Urogynecology Journal, 2019, 30, 2013-2022.	0.7	9
12	Do we need reminders of the precautionary principle?. BJOG: an International Journal of Obstetrics and Gynaecology, 2019, 126, 1423-1423.	1.1	0
13	There is still a place for vaginal mesh use in urogynaecology. BJOG: an International Journal of Obstetrics and Gynaecology, 2019, 126, 1075.	1.1	2
14	Effectiveness and cost-effectiveness of basic versus biofeedback-mediated intensive pelvic floor muscle training for female stress or mixed urinary incontinence: protocol for the OPAL randomised trial. BMJ Open, 2019, 9, e024153.	0.8	14
15	Re: Renaissance of the autologous pubovaginal sling. International Urogynecology Journal, 2018, 29, 615-615.	0.7	0
16	Management of female stress urinary incontinence. Maturitas, 2018, 110, 124.	1.0	0
17	Re: Ferdinando Fusco, Mohamed Abdel-Fattah, Christopher R. Chapple, et al. Updated Systematic Review and Meta-analysis of the Comparative Data on Colposuspensions, Pubovaginal Slings, and Midurethral Tapes in the Surgical Treatment of Female Stress Urinary Incontinence. Eur Urol 2017;72:567â€“91. European Urology, 2018, 73, e54-e55.	0.9	2
18	Ultrasound bladder wall thickness and detrusor overactivity: a multicentre test accuracy study. BJOG: an International Journal of Obstetrics and Gynaecology, 2017, 124, 1422-1429.	1.1	19

#	ARTICLE	IF	CITATIONS
19	Adverse events after first, single, mesh and non-mesh surgical procedures for stress urinary incontinence and pelvic organ prolapse in Scotland, 1997-2016: a population-based cohort study. <i>Lancet, The</i> , 2017, 389, 629-640.	6.3	115
20	Effect of procedure type and use of mesh on complications after surgery for pelvic organ prolapse: a retrospective cohort study. <i>Lancet, The</i> , 2015, 386, S56.	6.3	0
21	Bcl-2 may play a role in the progression of endometrial hyperplasia and early carcinogenesis, but not linked to further tumorigenesis. <i>Journal of Microscopy and Ultrastructure</i> , 2015, 3, 19.	0.1	7
22	Mixed urinary incontinence: international urogynecological association research and development committee opinion. <i>International Urogynecology Journal</i> , 2014, 25, 1303-1312.	0.7	39
23	Does traction on the cervix under anaesthesia tell us when to perform a concomitant hysterectomy? A 2-year follow-up of a prospective cohort study. <i>International Urogynecology Journal</i> , 2014, 25, 1213-1217.	0.7	4
24	Evaluation of three different surgical approaches in repairing paravaginal support defects: a comparative trial. <i>Archives of Gynecology and Obstetrics</i> , 2013, 288, 1341-1348.	0.8	16
25	Multicenter Prospective Randomized Study of Single-incision Mini-sling Vs Tension-free Vaginal Tape-obturator in Management of Female Stress Urinary Incontinence: A Minimum of 1-Year Follow-up. <i>Urology</i> , 2013, 82, 552-559.	0.5	40
26	Surgical Treatment of Recurrent Stress Urinary Incontinence in Women: A Systematic Review and Meta-analysis of Randomised Controlled Trials. <i>European Urology</i> , 2013, 64, 323-336.	0.9	45
27	Traction on the cervix in theatre before anterior repair: does it tell us when to perform a concomitant hysterectomy?. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2012, 160, 205-209.	0.5	16
28	Systematic review and meta-analysis of inside-out versus outside-in transobturator tapes in management of stress urinary incontinence in women. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2012, 162, 1-10.	0.5	26
29	A multicentre prospective randomised study of single-incision mini-sling (Adjust [®]) versus tension-free vaginal tape-obturator (TVT-O [®]) in the management of female stress urinary incontinence: pain profile and short-term outcomes. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2012, 165, 115-121.	0.5	49
30	Does pre-operative traction on the cervix approximate intra-operative uterine prolapse? Comment. <i>International Urogynecology Journal</i> , 2012, 23, 1811-1812.	0.7	2
31	Prospective multicentre study of adjustable single-incision mini-sling (Adjust [®]) in the management of stress urinary incontinence in women: 1-year follow-up study. <i>BJU International</i> , 2012, 109, 880-886.	1.3	61
32	Obesity and urinary incontinence in women: is the black box becoming grayer?. <i>International Urogynecology Journal</i> , 2011, 22, 257-258.	0.7	13
33	Unusual underlying cause of recurrent vulval abscess. <i>BMJ Case Reports</i> , 2010, 2010, bcr0820092142-bcr0820092142.	0.2	0
34	The relationship between bcl-2 oncogene expression and clinicopathological criteria in various stages of cervical neoplasia in Egyptian women. <i>European Journal of Gynaecological Oncology (discontinued)</i> , 2010, 31, 536-8.	0.3	1
35	Could the National Institute for Health and Clinical Excellence guidelines on urodynamics in urinary incontinence put some women at risk of a bad outcome from stress incontinence surgery?. <i>BJU International</i> , 2009, 103, 635-639.	1.3	60
36	Author Response to: The long-term effectiveness of antenatal pelvic floor muscle training: 8-year follow up of a randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2009, 116, 601-602.	1.1	0

#	ARTICLE	IF	CITATIONS
37	Efficacy of using electrosurgical bipolar vessel sealing during vaginal hysterectomy in patients with different degrees of operative difficulty: A randomised controlled trial. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2009, 147, 86-90.	0.5	39
38	The long-term effectiveness of antenatal pelvic floor muscle training: eight-year follow up of a randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2008, 115, 985-990.	1.1	48
39	Duloxetine for the treatment of stress urinary incontinence. <i>Expert Review of Obstetrics and Gynecology</i> , 2007, 2, 133-139.	0.4	2
40	Does antenatal pelvic floor muscle training affect the outcome of labour? A randomised controlled trial. <i>International Urogynecology Journal</i> , 2007, 19, 85-88.	0.7	28