

Oddmund Bakke

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9500238/oddmund-bakke-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

117
papers

6,532
citations

38
h-index

79
g-index

130
ext. papers

7,272
ext. citations

6.9
avg, IF

5.57
L-index

#	Paper	IF	Citations
117	De novo formation of early endosomes during Rab5-to-Rab7a transition. <i>Journal of Cell Science</i> , 2021 , 134,	5.3	4
116	Rab7b regulates dendritic cell migration by linking lysosomes to the actomyosin cytoskeleton. <i>Journal of Cell Science</i> , 2021 , 134,	5.3	3
115	Sjögren syndrome/scleroderma autoantigen 1 is a direct Tankyrase binding partner in cancer cells. <i>Communications Biology</i> , 2020 , 3, 123	6.7	3
114	Rab18 regulates focal adhesion dynamics by interacting with kinectin-1 at the endoplasmic reticulum. <i>Journal of Cell Biology</i> , 2020 , 219,	7.3	5
113	Invariant chain regulates endosomal fusion and maturation through an interaction with the SNARE Vti1b. <i>Journal of Cell Science</i> , 2020 , 133,	5.3	2
112	Micro-stepping extended focus reduces photobleaching and preserves structured illumination super-resolution features. <i>Journal of Cell Science</i> , 2020 , 133,	5.3	2
111	Receptor-Mediated Endocytosis of VEGF-A in Rat Liver Sinusoidal Endothelial Cells. <i>BioMed Research International</i> , 2019 , 2019, 5496197	3	5
110	Disentangling the immune response and host-pathogen interactions in <i>Francisella noatunensis</i> infected Atlantic cod. <i>Comparative Biochemistry and Physiology Part D: Genomics and Proteomics</i> , 2019 , 30, 333-346	2	6
109	Rab6 regulates cell migration and invasion by recruiting Cdc42 and modulating its activity. <i>Cellular and Molecular Life Sciences</i> , 2019 , 76, 2593-2614	10.3	12
108	Antifungal activity of well-defined chito-oligosaccharide preparations against medically relevant yeasts. <i>PLoS ONE</i> , 2019 , 14, e0210208	3.7	15
107	Antigen-delivery through invariant chain (CD74) boosts CD8 and CD4 T cell immunity. <i>Oncolimmunology</i> , 2019 , 8, 1558663	7.2	9
106	Rab7a modulates ER stress and ER morphology. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2018 , 1865, 781-793	4.9	9
105	Human c-SRC kinase (CSK) overexpression makes T cells dummy. <i>Cancer Immunology, Immunotherapy</i> , 2018 , 67, 525-536	7.4	6
104	TBC1D5 controls the GTPase cycle of Rab7b. <i>Journal of Cell Science</i> , 2018 , 131,	5.3	21
103	A protein kinase A-ezrin complex regulates connexin 43 gap junction communication in liver epithelial cells. <i>Cellular Signalling</i> , 2017 , 32, 1-11	4.9	13
102	Misdirection of endosomal trafficking mediated by herpes simplex virus-encoded glycoprotein B. <i>FASEB Journal</i> , 2017 , 31, 1650-1667	0.9	10
101	Overview of the membrane-associated RING-CH (MARCH) E3 ligase family. <i>New Biotechnology</i> , 2017 , 38, 7-15	6.4	30

100	Characterization of the role of RILP in cell migration. <i>European Journal of Histochemistry</i> , 2017 , 61, 2783-2811	2.1	6
99	Rab7b modulates autophagic flux by interacting with Atg4B. <i>EMBO Reports</i> , 2017 , 18, 1727-1739	6.5	22
98	Rab7a regulates cell migration through Rac1 and vimentin. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017 , 1864, 367-381	4.9	37
97	Endosomal binding kinetics of Eps15 and Hrs specifically regulate the degradation of RTKs. <i>Scientific Reports</i> , 2017 , 7, 17962	4.9	5
96	The multiple roles of Rab9 in the endolysosomal system. <i>Communicative and Integrative Biology</i> , 2016 , 9, e1204498	1.7	23
95	Spatiotemporal Resolution of Rab9 and CI-MPR Dynamics in the Endocytic Pathway. <i>Traffic</i> , 2016 , 17, 211-29	5.7	22
94	Bidirectional traffic between the Golgi and the endosomes - machineries and regulation. <i>Journal of Cell Science</i> , 2016 , 129, 3971-3982	5.3	57
93	Oligomerized, filamentous surface presentation of RANTES/CCL5 on vascular endothelial cells. <i>Scientific Reports</i> , 2015 , 5, 9261	4.9	17
92	Rab7b at the intersection of intracellular trafficking and cell migration. <i>Communicative and Integrative Biology</i> , 2015 , 8, e1023492	1.7	9
91	The human-specific invariant chain isoform Iip35 modulates Iip33 trafficking and function. <i>Immunology and Cell Biology</i> , 2014 , 92, 791-8	5	4
90	A novel interaction between Rab7b and actomyosin reveals a dual role in intracellular transport and cell migration. <i>Journal of Cell Science</i> , 2014 , 127, 4927-39	5.3	27
89	Emerging regulators of endosomal dynamics during mitosis. <i>Cell Cycle</i> , 2014 , 13, 349-50	4.7	
88	Invariant chain as a vehicle to load antigenic peptides on human MHC class I for cytotoxic T-cell activation. <i>European Journal of Immunology</i> , 2014 , 44, 774-84	6.1	16
87	B-cell tolerance to the B-cell receptor variable regions. <i>European Journal of Immunology</i> , 2013 , 43, 2577-87		4
86	BiP negatively affects ricin transport. <i>Toxins</i> , 2013 , 5, 969-82	4.9	9
85	Differential regulation of Rab GTPase expression in monocyte-derived dendritic cells upon lipopolysaccharide activation: a correlation to maturation-dependent functional properties. <i>PLoS ONE</i> , 2013 , 8, e73538	3.7	8
84	Dynamics of Rab7b-dependent transport of sorting receptors. <i>Traffic</i> , 2012 , 13, 1273-85	5.7	27
83	Rab GTPases are differentially regulated during DC maturation. <i>Molecular Immunology</i> , 2012 , 51, 8	4.3	

82	Differential regulation of MHC II and invariant chain expression during maturation of monocyte-derived dendritic cells. <i>Journal of Leukocyte Biology</i> , 2012 , 91, 729-37	6.5	9
81	Charcot-Marie-Tooth disease and intracellular traffic. <i>Progress in Neurobiology</i> , 2012 , 99, 191-225	10.9	48
80	The fusion of early endosomes induces molecular-motor-driven tubule formation and fission. <i>Journal of Cell Science</i> , 2012 , 125, 1910-9	5.3	40
79	Towards a systems understanding of MHC class I and MHC class II antigen presentation. <i>Nature Reviews Immunology</i> , 2011 , 11, 823-36	36.5	1090
78	Invariant chain increases the half-life of MHC II by delaying endosomal maturation. <i>Immunology and Cell Biology</i> , 2011 , 89, 619-29	5	21
77	SorLA regulates the activity of lipoprotein lipase by intracellular trafficking. <i>Journal of Cell Science</i> , 2011 , 124, 1095-105	5.3	58
76	Toll-like receptor 2 (P631H) mutant impairs membrane internalization and is a dominant negative allele. <i>Scandinavian Journal of Immunology</i> , 2010 , 71, 369-81	3.4	45
75	The protein phosphatase 1 regulator PNUTS is a new component of the DNA damage response. <i>EMBO Reports</i> , 2010 , 11, 868-75	6.5	52
74	Rab7b and receptors trafficking. <i>Communicative and Integrative Biology</i> , 2010 , 3, 401-4	1.7	24
73	Rab7b controls trafficking from endosomes to the TGN. <i>Journal of Cell Science</i> , 2010 , 123, 1480-91	5.3	87
72	Ubiquitination regulates MHC class II-peptide complex retention and degradation in dendritic cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 20465-70	11.5	81
71	Cadmium-induced inflammatory responses in cells relevant for lung toxicity: Expression and release of cytokines in fibroblasts, epithelial cells and macrophages. <i>Toxicology Letters</i> , 2010 , 193, 252-60	4.4	79
70	The Rab11a GTPase controls Toll-like receptor 4-induced activation of interferon regulatory factor-3 on phagosomes. <i>Immunity</i> , 2010 , 33, 583-96	32.3	151
69	Ultrastructural characterization of giant endosomes induced by GTPase-deficient Rab5. <i>Histochemistry and Cell Biology</i> , 2010 , 133, 41-55	2.4	80
68	MHC II and the endocytic pathway: regulation by invariant chain. <i>Scandinavian Journal of Immunology</i> , 2009 , 70, 184-93	3.4	59
67	Cell-cycle-dependent binding kinetics for the early endosomal tethering factor EEA1. <i>EMBO Reports</i> , 2008 , 9, 171-8	6.5	22
66	Major histocompatibility complex class II-peptide complexes internalize using a clathrin- and dynamin-independent endocytosis pathway. <i>Journal of Biological Chemistry</i> , 2008 , 283, 14717-27	5.4	99
65	Cellular trafficking of lipoteichoic acid and Toll-like receptor 2 in relation to signaling: role of CD14 and CD36. <i>Journal of Leukocyte Biology</i> , 2008 , 84, 280-91	6.5	114

64	Human PARM-1 is a novel mucin-like, androgen-regulated gene exhibiting proliferative effects in prostate cancer cells. <i>International Journal of Cancer</i> , 2008 , 122, 1229-35	7.5	19
63	Human NCU-G1 can function as a transcription factor and as a nuclear receptor co-activator. <i>BMC Molecular Biology</i> , 2007 , 8, 106	4.5	6
62	Sorting by the cytoplasmic domain of the amyloid precursor protein binding receptor SorLA. <i>Molecular and Cellular Biology</i> , 2007 , 27, 6842-51	4.8	152
61	Re-routing of the invariant chain to the direct sorting pathway by introduction of an AP3-binding motif from LIMP II. <i>European Journal of Cell Biology</i> , 2006 , 85, 457-67	6.1	15
60	Toll-like receptor 3 associates with c-Src tyrosine kinase on endosomes to initiate antiviral signaling. <i>EMBO Journal</i> , 2006 , 25, 3335-46	13	150
59	The adaptor protein AP-4 as a component of the clathrin coat machinery: a morphological study. <i>Biochemical Journal</i> , 2005 , 385, 503-10	3.8	28
58	The mouse CD1d cytoplasmic tail mediates CD1d trafficking and antigen presentation by adaptor protein 3-dependent and -independent mechanisms. <i>Journal of Immunology</i> , 2005 , 174, 3179-86	5.3	46
57	Characterization of a novel chemokine-containing storage granule in endothelial cells: evidence for preferential exocytosis mediated by protein kinase A and diacylglycerol. <i>Journal of Immunology</i> , 2005 , 175, 5358-69	5.3	55
56	c-Myb associates with PML in nuclear bodies in hematopoietic cells. <i>Experimental Cell Research</i> , 2004 , 297, 118-26	4.2	16
55	Rapid chemokine secretion from endothelial cells originates from 2 distinct compartments. <i>Blood</i> , 2004 , 104, 314-20	2.2	95
54	The cytoplasmic tail of invariant chain modulates antigen processing and presentation. <i>European Journal of Immunology</i> , 2003 , 33, 277-86	6.1	21
53	The adaptor protein AP-3 is required for CD1d-mediated antigen presentation of glycosphingolipids and development of Valpha14i NKT cells. <i>Journal of Experimental Medicine</i> , 2003 , 198, 1133-46	16.6	92
52	MHC class II loading of high or low affinity peptides directed by li/peptide fusion constructs: implications for T cell activation. <i>International Immunology</i> , 2003 , 15, 1291-9	4.9	14
51	Uncoating ATPase Hsc70 is recruited by invariant chain and controls the size of endocytic compartments. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 1515-20	11.5	37
50	Structural requirements for interactions between leucine-sorting signals and clathrin-associated adaptor protein complex AP3. <i>Journal of Biological Chemistry</i> , 2002 , 277, 47436-43	5.4	24
49	The cytoplasmic tail of invariant chain regulates endosome fusion and morphology. <i>Molecular Biology of the Cell</i> , 2002 , 13, 1846-56	3.5	36
48	Mechanism of interaction between leucine-based sorting signals from the invariant chain and clathrin-associated adaptor protein complexes AP1 and AP2. <i>Journal of Biological Chemistry</i> , 2002 , 277, 16484-8	5.4	31
47	Mitotic partitioning of endosomes and lysosomes. <i>Current Biology</i> , 2001 , 11, 644-51	6.3	76

46	The cytoplasmic tail of CD1d contains two overlapping basolateral sorting signals. <i>Journal of Biological Chemistry</i> , 2000 , 275, 8279-82	5.4	27
45	The MHC class II-associated chicken invariant chain shares functional properties with its mammalian homologs. <i>Experimental Cell Research</i> , 2000 , 259, 360-9	4.2	16
44	Overexpression of proteins containing tyrosine- or leucine-based sorting signals affects transferrin receptor trafficking. <i>Journal of Biological Chemistry</i> , 1999 , 274, 21139-48	5.4	10
43	The leucine-based sorting motifs in the cytoplasmic domain of the invariant chain are recognized by the clathrin adaptors AP1 and AP2 and their medium chains. <i>Journal of Biological Chemistry</i> , 1999 , 274, 36153-8	5.4	83
42	Intracellular traffic to compartments for MHC class II peptide loading: signals for endosomal and polarized sorting. <i>Immunological Reviews</i> , 1999 , 172, 171-87	11.3	35
41	Post-replicative base excision repair in replication foci. <i>EMBO Journal</i> , 1999 , 18, 3834-44	13	265
40	The leucine-based motif DDQxxLI is recognized both for internalization and basolateral sorting of invariant chain in MDCK cells. <i>European Journal of Cell Biology</i> , 1998 , 76, 25-32	6.1	24
39	Selection of phage displayed peptides from a random 10-mer library recognising a peptide target. <i>Immunotechnology: an International Journal of Immunological Engineering</i> , 1998 , 4, 21-8		12
38	Major histocompatibility complex class II-dependent antigen presentation by human intestinal endothelial cells. <i>Gastroenterology</i> , 1998 , 114, 649-56	13.3	34
37	A region from the medium chain adaptor subunit (μ) recognizes leucine- and tyrosine-based sorting signals. <i>Journal of Biological Chemistry</i> , 1998 , 273, 8638-45	5.4	56
36	Medium chains of adaptor complexes AP-1 and AP-2 recognize leucine-based sorting signals from the invariant chain. <i>Journal of Biological Chemistry</i> , 1998 , 273, 6005-8	5.4	89
35	Intracellular transport of molecules engaged in the presentation of exogenous antigens. <i>Current Topics in Microbiology and Immunology</i> , 1998 , 232, 179-215	3.3	15
34	Exon 6 is essential for invariant chain trimerization and induction of large endosomal structures. <i>Journal of Biological Chemistry</i> , 1997 , 272, 8281-7	5.4	32
33	MHC class II-associated invariant chain-induced enlarged endosomal structures: a morphological study. <i>Experimental Cell Research</i> , 1997 , 235, 79-92	4.2	34
32	A new triple-stranded alpha-helical bundle in solution: the assembling of the cytosolic tail of MHC-associated invariant chain. <i>Structure</i> , 1997 , 5, 1453-64	5.2	16
31	The various roles of invariant chain in the act of antigen Presentation 1996 , 15-41		1
30	Antigen presentation mediated by recycling of surface HLA-DR molecules. <i>Nature</i> , 1995 , 375, 603-6	50.4	238
29	Invariant chain induces a delayed transport from early to late endosomes. <i>Journal of Biological Chemistry</i> , 1995 , 270, 2741-6	5.4	36

28	Structure-activity relationship of the leucine-based sorting motifs in the cytosolic tail of the major histocompatibility complex-associated invariant chain. <i>Journal of Biological Chemistry</i> , 1995 , 270, 27165-71	5.4	43
27	Physiological functions of endosomal proteolysis. <i>Biochemical Journal</i> , 1995 , 307 (Pt 2), 313-26	3.8	80
26	Targeting major histocompatibility complex class II molecules to the cell surface by invariant chain allows antigen presentation upon recycling. <i>European Journal of Immunology</i> , 1994 , 24, 873-83	6.1	20
25	The invariant chain inhibits presentation of endogenous antigens by a human fibroblast cell line. <i>European Journal of Immunology</i> , 1994 , 24, 1632-9	6.1	38
24	The bio-logical role of invariant chain (Ii) in MHC class II antigen presentation. <i>Immunology Letters</i> , 1994 , 43, 47-55	4.1	5
23	Targeting of membrane proteins to endosomes and lysosomes. <i>Trends in Cell Biology</i> , 1994 , 4, 292-7	18.3	279
22	Nuclear and mitochondrial forms of human uracil-DNA glycosylase are encoded by the same gene. <i>Nucleic Acids Research</i> , 1993 , 21, 2579-84	20.1	120
21	Relationship between invariant chain expression and major histocompatibility complex class II transport into early and late endocytic compartments. <i>Journal of Experimental Medicine</i> , 1993 , 177, 583-96	16.6	138
20	Intracellular distribution of the MHC class II molecules and the associated invariant chain (Ii) in different cell lines. <i>International Immunology</i> , 1993 , 5, 903-17	4.9	51
19	Cell surface HLA-DR-invariant chain complexes are targeted to endosomes by rapid internalization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993 , 90, 8581-5	11.5	189
18	Intracellular transport and localization of major histocompatibility complex class II molecules and associated invariant chain. <i>Journal of Cell Biology</i> , 1991 , 115, 1213-23	7.3	137
17	MHC class II-associated invariant chain contains a sorting signal for endosomal compartments. <i>Cell</i> , 1990 , 63, 707-16	56.2	526
16	A flow cytometric and immunofluorescence microscopic study of tumor necrosis factor production and localization in human monocytes. <i>Cellular Immunology</i> , 1989 , 122, 405-15	4.4	37
15	Retinoic acid induces a specific membrane glycoprotein in human epithelial cell lines. <i>Experimental Cell Research</i> , 1989 , 180, 20-9	4.2	6
14	The Association of the Glucocorticoid Receptor with Mr 90,000 Heat Shock Protein and Tubulin 1989 , 41-53		
13	Intracellular localization of the glucocorticoid receptor: evidence for cytoplasmic and nuclear localization. <i>Endocrinology</i> , 1987 , 120, 1232-42	4.8	180
12	Effects of retinoic acid on cytokeratin and actin filaments in human NHIK 3025 cells. <i>Biochemical Society Transactions</i> , 1987 , 15, 858-859	5.1	1
11	Characterization and sequence-specific binding to mouse mammary tumor virus DNA of purified activated human glucocorticoid receptor. <i>Biochemistry</i> , 1987 , 26, 1697-704	3.2	16

10	Cell-mediated inhibition of proliferation and activation of alloreactive cytotoxic lymphocytes: maintenance of response potential of precursors and dissociation between proliferation and effector function of activated cytotoxic lymphocytes. <i>Cellular Immunology</i> , 1986 , 101, 105-21	4.4	3
9	Concentration-dependent effects of potassium dichromate on the cell cycle. <i>Cytometry</i> , 1984 , 5, 482-6		25
8	Steroid Structure Requirements, Cell Cycle Specificity and Protein Metabolism in Glucocorticoid Growth Regulation of Human NHIK 3025 Cells 1984 , 363-384		
7	Soluble cytostatic factor(s) released from human monocytes. II. Effects on target cell kinetics. <i>Scandinavian Journal of Immunology</i> , 1983 , 18, 13-20	3.4	15
6	Structure requirements for glucocorticoid growth of a human cell line (NHIK 3025). <i>The Journal of Steroid Biochemistry</i> , 1982 , 17, 489-93		4
5	The role of protein metabolism in glucocorticoid-induced prolongation of G1 phase in human NHIK 3025 cells. <i>Journal of Cellular Physiology</i> , 1982 , 113, 459-64	7	5
4	Effects of potassium dichromate on the cell cycle of an established human cell line (NHIK 3025). <i>Toxicology</i> , 1982 , 24, 281-92	4.4	15
3	Cell cycle-specific glucocorticoid growth regulation of a human cell line (NHIK 3025). <i>Journal of Cellular Physiology</i> , 1981 , 109, 489-96	7	29
2	Cell cycle characteristics of synchronized and asynchronous populations of human cells and effect of cooling of selected mitotic cells. <i>Cell Proliferation</i> , 1977 , 10, 511-22	7.9	20
1	A fast and accurate method for calculating Engelberg's synchronization index. <i>Cell Proliferation</i> , 1976 , 9, 389-93	7.9	3