

# Enrique Macpherson

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9502969/publications.pdf>

Version: 2024-02-01

150  
papers

6,796  
citations

76196

40  
h-index

69108

77  
g-index

153  
all docs

153  
docs citations

153  
times ranked

7235  
citing authors

#	ARTICLE	IF	CITATIONS
1	New species of deep-sea squat lobsters (Decapoda: Anomura: Galatheoidea) from Guadeloupe, French West Indies, unveiled through integrative taxonomy. <i>Journal of Crustacean Biology</i> , 2022, 42, .	0.3	4
2	Warming intensifies the interaction between the temperate seagrass <i>Posidonia oceanica</i> and its dominant fish herbivore <i>Sarpa salpa</i> . <i>Marine Environmental Research</i> , 2021, 165, 105237.	1.1	15
3	A new species of <i>Munida</i> Leach, 1820 (Crustacea: Decapoda: Anomura: Munididae) from seamounts of the Nazca-Desventuradas Marine Park. <i>PeerJ</i> , 2021, 9, e10531.	0.9	2
4	Temperature reduces fish dispersal as larvae grow faster to their settlement size. <i>Journal of Animal Ecology</i> , 2021, 90, 1419-1432.	1.3	24
5	Impact of individual early life traits in larval dispersal: A multispecies approach using backtracking models. <i>Progress in Oceanography</i> , 2021, 192, 102518.	1.5	20
6	Revision of the squat lobsters of the genus <i>Phylladorhynchus</i> Baba, 1969 (Crustacea, Decapoda,) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50</i>	0.2	5
7	Alain Crosnier's role in modern carcinology: exploration, international collaboration, and taxonomy. <i>Journal of Crustacean Biology</i> , 2021, 41, .	0.3	1
8	Individual-based population genomics reveal different drivers of adaptation in sympatric fish. <i>Scientific Reports</i> , 2020, 10, 12683.	1.6	13
9	New occurrences of squat lobsters of the genus <i>Eumunida</i> Smith, 1883 (Decapoda, Eumunididae) in New Caledonia, the Solomon Islands and Papua-New Guinea, with the description of a new species. <i>Zootaxa</i> , 2020, 4786, zootaxa.4786.4.2.	0.2	1
10	Deep-sea squat lobster biogeography (Munidopsidae: <i>Leiogalathea</i> ) unveils Tethyan vicariance and evolutionary patterns shared by shallow-water relatives. <i>Zoologica Scripta</i> , 2020, 49, 340-356.	0.7	9
11	Helping decision making for reliable and cost-effective RAD sequencing and genotyping analyses in non-model species. <i>Molecular Ecology Resources</i> , 2020, 20, 795-806.	2.2	12
12	A new species of squat lobster of the genus <i>Hendersonida</i> (Crustacea, Decapoda, Munididae) from Papua New Guinea. <i>ZooKeys</i> , 2020, 935, 25-35.	0.5	2
13	On some squat lobsters from India (Decapoda, Anomura, Munididae), with description of a new species of <i>Paramunida</i> Baba, 1988. <i>ZooKeys</i> , 2020, 965, 17-36.	0.5	6
14	High morphological similarity coupled with high genetic differentiation in new sympatric species of coral-reef squat lobsters (Crustacea: Decapoda: Galatheidae). <i>Zoological Journal of the Linnean Society</i> , 2019, 185, 984-1017.	1.0	13
15	Species delimitation and multi-locus species tree solve an old taxonomic problem for European squat lobsters of the genus <i>Munida</i> Leach, 1820. <i>Marine Biodiversity</i> , 2019, 49, 1751-1773.	0.3	16
16	Revision of the squat lobsters of the genus <i>Leiogalathea</i> Baba, 1969 (Crustacea, Decapoda,) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 142 T</i>	0.2	10
17	Evolution of squat lobsters (Crustacea, Galatheoidea): mitogenomic data suggest an early divergent Porcellanidae. <i>Hydrobiologia</i> , 2019, 833, 173-184.	1.0	5
18	How good is your marine protected area at curbing threats?. <i>Biological Conservation</i> , 2018, 221, 237-245.	1.9	69

#	ARTICLE	IF	CITATIONS
19	The richness of small pockets: Decapod species peak in small seagrass patches where fish predators are absent. <i>Marine Environmental Research</i> , 2018, 142, 1-6.	1.1	8
20	Three new species of squat lobsters of the genus <i>Munidopsis</i> Whiteaves, 1874, from Guadeloupe Island, Caribbean Sea (Crustacea, Decapoda, Munidopsidae). <i>Zootaxa</i> , 2018, 4422, 569.	0.2	8
21	Updating changes in the Iberian decapod crustacean fauna (excluding crabs) after 50 years. <i>Scientia Marina</i> , 2018, 82, 207.	0.3	7
22	Population genomics of an endemic Mediterranean fish: differentiation by fine scale dispersal and adaptation. <i>Scientific Reports</i> , 2017, 7, 43417.	1.6	83
23	Three new species of squat lobsters of the genus <i>Fennerogalatea</i> Baba, 1988 (Decapoda: Galatheidae) from the Pacific Ocean. <i>Zootaxa</i> , 2017, 4276, 46.	0.2	5
24	Phylogeny and evolution of shallow-water squat lobsters (Decapoda, Galatheoidea) from the Indo-Pacific. <i>Zoologica Scripta</i> , 2017, 46, 584-595.	0.7	11
25	A new species of squat lobster of the genus <i>Munida</i> (Galatheoidea, Munididae) from the Red Sea. <i>Crustaceana</i> , 2017, 90, 1005-1014.	0.1	1
26	<i>Uroptychus tuerkayi</i> sp. nov. (Anomura, Chirostylidae), a new squat lobster from the Atlantis-Great Meteor Seamount Chain in the eastern Atlantic. <i>Crustaceana</i> , 2017, 90, 807-817.	0.1	0
27	<i>Cancer strigosus</i> Linnaeus, 1760: neotype selection, its identification with <i>Cancer cancharus</i> Linnaeus, 1758, and reversal of precedence (Crustacea: Decapoda: Galatheidae). <i>Zootaxa</i> , 2017, 4323, 440.	0.2	0
28	Impact of life history traits on gene flow: A multispecies systematic review across oceanographic barriers in the Mediterranean Sea. <i>PLoS ONE</i> , 2017, 12, e0176419.	1.1	125
29	New sibling species and new occurrences of squat lobsters (Crustacea, Decapoda) from the western Indian Ocean. <i>European Journal of Taxonomy</i> , 2017, , .	0.6	2
30	Null alleles are ubiquitous at microsatellite loci in the Wedge Clam ( <i>Donax trunculus</i> ). <i>PeerJ</i> , 2017, 5, e3188.	0.9	35
31	Implications for management and conservation of the population genetic structure of the wedge clam <i>Donax trunculus</i> across two biogeographic boundaries. <i>Scientific Reports</i> , 2016, 6, 39152.	1.6	27
32	Temporal and spatial genetic differentiation in the crab <i>Liocarcinus depurator</i> across the Atlantic-Mediterranean transition. <i>Scientific Reports</i> , 2016, 6, 29892.	1.6	26
33	Some species of <i>Munidopsis</i> from the Gulf of Mexico, Florida Straits and Caribbean Sea (Decapoda:). <i>Tj ETQq1 1 0.784314 rgBT /Overlor</i>	0.2	6
34	The Enlargement of the Canal and Introduction of Non-Indigenous Species to the Mediterranean Sea. <i>Limnology and Oceanography Bulletin</i> , 2015, 24, 43-45.	0.2	38
35	Species of the genus <i>Galathea</i> Fabricius, 1793 (Crustacea, Decapoda, Galatheidae) from the Indian and Pacific Oceans, with descriptions of 92 new species. <i>Zootaxa</i> , 2015, 3913, 1-335.	0.2	11
36	“Double trouble”: the expansion of the Suez Canal and marine bioinvasions in the Mediterranean Sea. <i>Biological Invasions</i> , 2015, 17, 973-976.	1.2	170

#	ARTICLE	IF	CITATIONS
37	A new species of <i>Munidopsis</i> from a seamount of the Southwest Indian Ocean Ridge (Decapoda): <i>Tj ETQq1 1 0.784314 rgBT /Overloc</i>	0.2	11
38	<scp>SNP</scp> development from <scp>RNA</scp>â€seq data in a nonmodel fish: how many individuals are needed for accurate allele frequency prediction?. <i>Molecular Ecology Resources</i> , 2014, 14, 157-165.	2.2	38
39	Kinship analyses identify fish dispersal events on a temperate coastline. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2014, 281, 20140556.	1.2	39
40	Transcriptome analyses and differential gene expression in a non-model fish species with alternative mating tactics. <i>BMC Genomics</i> , 2014, 15, 167.	1.2	76
41	Distribution and biogeographic trends of decapod assemblages from Galicia Bank (NE Atlantic) at depths between 700 and 1800m, with connexions to regional water masses. <i>Deep-Sea Research Part II: Topical Studies in Oceanography</i> , 2014, 106, 165-178.	0.6	34
42	Large-Scale Assessment of Mediterranean Marine Protected Areas Effects on Fish Assemblages. <i>PLoS ONE</i> , 2014, 9, e91841.	1.1	146
43	A new species of <i>Paramunida</i> Baba, 1988 from the Central Pacific Ocean and a new genus to accommodate <i>P. granulata</i> (Henderson, 1885). <i>ZooKeys</i> , 2014, 425, 15-32.	0.5	4
44	Methane-Carbon Flow into the Benthic Food Web at Cold Seeps â€ A Case Study from the Costa Rica Subduction Zone. <i>PLoS ONE</i> , 2013, 8, e74894.	1.1	70
45	<strong>A new genus and some new species of the genus <em>Lauriea</em> Baba, 1971 (Crustacea, Decapoda, Galatheidae) from the Pacific and Indian Oceans, using molecular and morphological characters</strong>. <i>Zootaxa</i> , 2013, 3599, 136-160.	0.2	12
46	The Structure of Mediterranean Rocky Reef Ecosystems across Environmental and Human Gradients, and Conservation Implications. <i>PLoS ONE</i> , 2012, 7, e32742.	1.1	275
47	Genetic characterization of the endangered and endemic anchialine squat lobster <i>Munidopsis polymorpha</i> from Lanzarote (Canary Islands): management implications. <i>ICES Journal of Marine Science</i> , 2012, 69, 1030-1037.	1.2	10
48	The Magnitude of Global Marine Species Diversity. <i>Current Biology</i> , 2012, 22, 2189-2202.	1.8	797
49	New deep-sea squat lobsters of the genus <i>Galathea</i> Fabricius, 1793 (Decapoda, Galatheidae) from Vanuatu and New Caledonia. <i>Zoosystema</i> , 2012, 34, 409-427.	0.2	3
50	Phylogeographic patterns of decapod crustaceans at the Atlanticâ€Mediterranean transition. <i>Molecular Phylogenetics and Evolution</i> , 2012, 62, 664-672.	1.2	59
51	Permanent Genetic Resources added to Molecular Ecology Resources Database 1 April 2012 â€ 31 May 2012. <i>Molecular Ecology Resources</i> , 2012, 12, 972-974.	2.2	18
52	DEEP UNDER THE SEA: UNRAVELING THE EVOLUTIONARY HISTORY OF THE DEEP-SEA SQUAT LOBSTER <i>PARAMUNIDA</i> (DECAPODA, MUNIDIDAE). <i>Evolution; International Journal of Organic Evolution</i> , 2012, 66, 1878-1896.	1.1	28
53	A new squat lobster (Crustacea: Decapoda: Anomura: Chirostylidae) from off NW Spain. <i>Zootaxa</i> , 2012, 3224, 49.	0.2	5
54	The squat lobsters of the genus <i>Sadayoshia</i> Baba, 1969 (Crustacea: Decapoda: Anomura: Munididae): new records including six new species from the Pacific Ocean. <i>Zootaxa</i> , 2012, 3589, 30.	0.2	5

#	ARTICLE	IF	CITATIONS
55	Barcoding type specimens helps to identify synonyms and an unnamed new species in Eumunida Smith, 1883 (Decapoda : Eumunididae). <i>Invertebrate Systematics</i> , 2011, 25, 322.	0.5	48
56	Matching genetics with oceanography: directional gene flow in a Mediterranean fish species. <i>Molecular Ecology</i> , 2011, 20, 5167-5181.	2.0	121
57	Effect of oceanographic barriers and overfishing on the population genetic structure of the European spiny lobster (<i>Palinurus elephas</i>). <i>Biological Journal of the Linnean Society</i> , 2011, 104, 407-418.	0.7	30
58	Allogalathea (Decapoda: Galatheidae): a monospecific genus of squat lobster?. <i>Zoological Journal of the Linnean Society</i> , 2011, 162, 245-270.	1.0	23
59	Genetic connectivity patterns in an endangered species: The dusky grouper ( <i>Epinephelus marginatus</i> ). <i>Journal of Experimental Marine Biology and Ecology</i> , 2011, 401, 126-133.	0.7	40
60	A new squat lobster of the genus <i>Munidopsis</i> (Crustacea: Decapoda: Munidopsidae) from the Mediterranean Sea. <i>Scientia Marina</i> , 2011, 75, 525-532.	0.3	6
61	Squat lobster assemblages on seamounts differ from some, but not all, deep-sea habitats of comparable depth. <i>Marine Ecology</i> , 2010, 31, 63-83.	0.4	37
62	Shallow-water squat lobsters (Crustacea, Decapoda, Galatheidae) from Mayotte (Comoros Island), La Réunion and Madagascar, with the description of a new genus and two new species. <i>Zootaxa</i> , 2010, 2612, 57.	0.2	13
63	Taxonomic revision of the genus <i>Paramunida</i> Baba, 1988 (Crustacea: Decapoda: Galatheidae): a morphological and molecular approach. <i>Zootaxa</i> , 2010, 2712, .	0.2	17
64	A new classification of the Galatheaidea (Crustacea: Decapoda: Anomura). <i>Zootaxa</i> , 2010, 2676, 57.	0.2	101
65	Occurrence of lithodid crabs (Decapoda, Lithodidae) on the Pacific coast of Costa Rica, Central America. <i>Crustaceana</i> , 2010, 83, 143-151.	0.1	7
66	Biogeography of the deep-sea galatheid squat lobsters of the Pacific Ocean. <i>Deep-Sea Research Part I: Oceanographic Research Papers</i> , 2010, 57, 228-238.	0.6	38
67	Genetic Diversity Levels in Fishery-Exploited Spiny Lobsters of the Genus <i>Palinurus</i> (Decapoda:) <a href="#">Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 1</a>	0.3	8
68	Rapid radiation in spiny lobsters ( <i>Palinurus</i> spp) as revealed by classic and ABC methods using mtDNA and microsatellite data. <i>BMC Evolutionary Biology</i> , 2009, 9, 263.	3.2	31
69	Development and characterization of microsatellite markers for the endangered anchialine squat lobster <i>Munidopsis polymorpha</i> . <i>Conservation Genetics</i> , 2009, 10, 673-676.	0.8	4
70	Morphological and molecular description of new species of squat lobster (Crustacea: Decapoda:) <a href="#">Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 1</a> <i>Linnean Society</i> , 2009, 156, 465-493.	1.0	38
71	Phylogenetic relationships between spiny, slipper and coral lobsters (Crustacea, Decapoda, Achelata). <i>Molecular Phylogenetics and Evolution</i> , 2009, 50, 152-162.	1.2	65
72	Patterns of genetic differentiation between two co-occurring demersal species: the red mullet ( <i>Mullus barbatus</i> ) and the striped red mullet ( <i>Mullus surmuletus</i> ). <i>Canadian Journal of Fisheries and Aquatic Sciences</i> , 2009, 66, 1478-1490.	0.7	27

#	ARTICLE	IF	CITATIONS
73	New species of squat lobsters of the genera <i>Munida</i> and <i>Raymunida</i> (Crustacea, Decapoda, Galatheididae) from Vanuatu and New Caledonia. <i>Zoosystema</i> , 2009, 31, 431-451.	0.2	8
74	The influence of oceanographic fronts and early-life-history traits on connectivity among littoral fish species. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 1473-1478.	3.3	263
75	Utility of pairwise mtDNA genetic distances for predicting cross-species microsatellite amplification and polymorphism success in fishes. <i>Conservation Genetics</i> , 2008, 9, 181-190.	0.8	29
76	Phylogeography of the European spiny lobster ( <i>Palinurus elephas</i> ): Influence of current oceanographical features and historical processes. <i>Molecular Phylogenetics and Evolution</i> , 2008, 48, 708-717.	1.2	82
77	A New Genus of Squat Lobster (Decapoda: Anomura: Galatheididae) from the South West Pacific and Indian Ocean Inferred from Morphological and Molecular Evidence. <i>Journal of Crustacean Biology</i> , 2008, 28, 68-75.	0.3	22
78	Some lithodid crabs (Crustacea: Decapoda: Lithodidae) from Taiwan and adjacent waters, with the description of one new species from Guam. <i>Zootaxa</i> , 2008, 1924, 43-52.	0.2	5
79	Catalogue of squat lobsters of the world (Crustacea: Decapoda: Anomura—families Chirostylidae.) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 467</i>	0.2	145
80	Molecular Phylogeny of the East Pacific Squat Lobsters of the Genus <i>Munidopsis</i> (Decapoda:) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 467</i>	0.3	40
81	Cross-amplification of 10 new isolated polymorphic microsatellite loci for red mullet ( <i>Mullus</i> ) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10</i>	1.7	6
82	Isolation and characterization of nine polymorphic microsatellite markers in the two-banded sea bream ( <i>Diplodus vulgaris</i> ) and cross-species amplification in the white sea bream ( <i>Diplodus sargus</i> ) and the saddled bream ( <i>Oblada melanura</i> ). <i>Molecular Ecology Notes</i> , 2007, 7, 661-663.	1.7	14
83	High self-recruitment levels in a Mediterranean littoral fish population revealed by microsatellite markers. <i>Marine Biology</i> , 2007, 151, 719-727.	0.7	45
84	Polymorphic microsatellite loci for the cardinal fish ( <i>Apogon imberbis</i> ). <i>Conservation Genetics</i> , 2007, 8, 1251-1253.	0.8	2
85	Isolation of eight microsatellites loci from the saddled bream, <i>Oblada melanura</i> and cross-species amplification in two sea bream species of the genus <i>Diplodus</i> . <i>Conservation Genetics</i> , 2007, 8, 1255-1257.	0.8	6
86	Species of the genus <i>Munidopsis</i> Whiteaves, 1784 from the Indian and Pacific Oceans and reestablishment of the genus <i>Galacantha</i> A. Milne-Edwards, 1880 (Crustacea, Decapoda, Galatheididae). <i>Zootaxa</i> , 2007, 1417, 1-135.	0.2	38
87	A review of the <i>Tripterygion tripteronotus</i> (Risso, 1810) complex, with a description of a new species from the Mediterranean Sea (Teleostei: Tripterygiidae). <i>Scientia Marina</i> , 2007, 71, 75-86.	0.3	11
88	Some new records of shallow-water galatheid crustaceans (Anomura: Galatheididae) from the Dampier Archipelago, Western Australia. <i>Records of the Western Australian Museum, Supplement</i> , 2007, 73, 289.	0.5	5
89	Pere Rubi's Guardiola (1949-2007). <i>Scientia Marina</i> , 2007, 71, 825-826.	0.3	0
90	Effect of brine discharge from a desalination plant on macrobenthic communities in the NW Mediterranean. <i>Marine Environmental Research</i> , 2006, 62, 1-14.	1.1	64

#	ARTICLE	IF	CITATIONS
91	Characterization of 12 microsatellite markers in <i>Serranus cabrilla</i> (Pisces: Serranidae). <i>Molecular Ecology Notes</i> , 2006, 6, 204-206.	1.7	4
92	Isolation and characterization of polymorphic microsatellite markers for peacock wrasse ( <i>Symphodus tinca</i> ). <i>Molecular Ecology Notes</i> , 2006, 6, 747-749.	1.7	2
93	Population structure within and between subspecies of the Mediterranean triplefin fish <i>Tripterygion delaisi</i> revealed by highly polymorphic microsatellite loci. <i>Molecular Ecology</i> , 2006, 15, 3527-3539.	2.0	59
94	Seamount endemism questioned by the geographic distribution and population genetic structure of marine invertebrates. <i>Marine Biology</i> , 2006, 149, 1463-1475.	0.7	162
95	First stage zoeal descriptions of five Galatheaidea species from Western Pacific (Crustacea: Decapoda: Tj ETQq1 10,784314, rgBT /Ove	0.2	15
96	Relationship between pelagic larval duration and geographic distribution of Mediterranean littoral fishes. <i>Marine Ecology - Progress Series</i> , 2006, 327, 257-265.	0.9	120
97	Rapid radiation and cryptic speciation in mediterranean triplefin blennies (Pisces: Tripterygiidae) combining multiple genes. <i>Molecular Phylogenetics and Evolution</i> , 2005, 37, 751-761.	1.2	41
98	Ecological impact of coastal defence structures on sediment and mobile fauna: Evaluating and forecasting consequences of unavoidable modifications of native habitats. <i>Coastal Engineering</i> , 2005, 52, 1027-1051.	1.7	180
99	Environmental influences on temporal patterns of settlement in two littoral labrid fishes in the Mediterranean Sea. <i>Estuarine, Coastal and Shelf Science</i> , 2005, 63, 479-487.	0.9	24
100	Species of the genus <i>Munidopsis</i> (Crustacea, Decapoda, Galatheaidae) from the deep Atlantic Ocean, including cold-seep and hydrothermal vent areas. <i>Zootaxa</i> , 2005, 1095, 1-60.	0.2	65
101	Use of morphological and molecular data to identify three new sibling species of the genus <i>Munida</i> Leach, 1820 (Crustacea, Decapoda, Galatheaidae) from New Caledonia. <i>Journal of Natural History</i> , 2005, 39, 819-834.	0.2	29
102	Effect of pelagic larval growth and size-at-hatching on post-settlement survivorship in two temperate labrid fish of the genus <i>Symphodus</i> . <i>Marine Ecology - Progress Series</i> , 2005, 285, 205-211.	0.9	50
103	Population structure and reproduction of three sympatric species of hermit crabs in the north-western Mediterranean. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2004, 84, 371-376.	0.4	19
104	Isolation and characterization of microsatellite loci in <i>Tripterygion delaisi</i> . <i>Molecular Ecology Notes</i> , 2004, 4, 438-439.	1.7	10
105	A new species and new records of lithodid crabs (Crustacea: Decapoda: Lithodidae) from the Crozet and Kerguelen Islands area (Subantarctica). <i>Polar Biology</i> , 2004, 27, 418-422.	0.5	17
106	Rapid radiation and cryptic speciation in squat lobsters of the genus <i>Munida</i> (Crustacea, Decapoda) and related genera in the South West Pacific: molecular and morphological evidence. <i>Molecular Phylogenetics and Evolution</i> , 2004, 33, 259-279.	1.2	82
107	Species range size distributions for some marine taxa in the Atlantic Ocean. Effect of latitude and depth. <i>Biological Journal of the Linnean Society</i> , 2003, 80, 437-455.	0.7	54
108	Some lithodid crabs (Crustacea, Decapoda, Lithodidae) from the Salomon Islands (SW Pacific Ocean) with the description of a new species. <i>Scientia Marina</i> , 2003, 67, 413-418.	0.3	8

#	ARTICLE	IF	CITATIONS
109	On the genus <i>Munida</i> Leach, 1820 (Decapoda, Galatheididae) from the western and southern Indian Ocean, with the description of four new species. <i>Crustaceana</i> , 2002, 75, 465-484.	0.1	12
110	Large-scale species richness gradients in the Atlantic Ocean. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2002, 269, 1715-1720.	1.2	166
111	Biomass Size Spectra in Littoral Fishes in Protected and Unprotected Areas in the NW Mediterranean. <i>Estuarine, Coastal and Shelf Science</i> , 2002, 55, 777-788.	0.9	63
112	Planktonic larval duration and settlement marks on the otoliths of Mediterranean littoral fishes. <i>Marine Biology</i> , 2001, 138, 1115-1120.	0.7	127
113	Direct estimation of natural mortality rates for littoral marine fishes using populational data from a marine reserve. <i>Marine Biology</i> , 2000, 137, 1067-1076.	0.7	29
114	RAYMUNIDA, NEW GENUS (DECAPODA: ANOMURA: GALATHEIDAE) FROM THE INDIAN AND PACIFIC OCEANS. <i>Journal of Crustacean Biology</i> , 2000, 20, 253-258.	0.3	16
115	The mine tailing accident in Aznalcollar. <i>Science of the Total Environment</i> , 1999, 242, 3-11.	3.9	378
116	Spatio-temporal variability in growth of juvenile sparid fishes from the Mediterranean littoral zone. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 1999, 79, 137-143.	0.4	50
117	Temporal and spatial variability of settlement success and recruitment level in three blennoid fishes in the northwestern Mediterranean. <i>Marine Ecology - Progress Series</i> , 1999, 182, 269-282.	0.9	24
118	Ontogenetic shifts in habitat use and aggregation in juvenile sparid fishes. <i>Journal of Experimental Marine Biology and Ecology</i> , 1998, 220, 127-150.	0.7	145
119	Spatial and temporal patterns of settlement among sparid fishes of the genus <i>Diplodus</i> in the northwestern Mediterranean. <i>Marine Ecology - Progress Series</i> , 1998, 168, 45-56.	0.9	90
120	Mortality of juvenile fishes of the genus <i>Diplodus</i> in protected and unprotected areas in the western Mediterranean Sea. <i>Marine Ecology - Progress Series</i> , 1997, 160, 135-147.	0.9	76
121	Biomass spectra in benthic fish assemblages in the Benguela System. <i>Marine Ecology - Progress Series</i> , 1996, 138, 27-32.	0.9	35
122	Substrate use and temporal pattern of recruitment in juvenile fishes of the Mediterranean littoral. <i>Marine Biology</i> , 1995, 124, 35-42.	0.7	195
123	<i>Gastroptychus Formosus</i> (Filhol, 1884) (Decapoda, Anomura, Chirostylidae): Taxonomic History and First Record From the Western Atlantic. <i>Crustaceana</i> , 1995, 68, 484-488.	0.1	3
124	<i>Gastroptychus Formosus</i> (Filhol, 1884) (Decapoda, Anomura, Chirostylidae): Taxonomic History and First Record From the Western Atlantic. <i>Crustaceana</i> , 1995, 68, 484-488.	0.1	2
125	Some observations on the pelagic decapod <i>Pasiphaea semispinosa</i> Holthuis 1951 in the Benguela upwelling system. <i>African Journal of Marine Science</i> , 1994, 14, 59-67.	0.6	5
126	Effect of prey densities on cannibalism in Cape hake ( <i>Merluccius capensis</i> ) off Namibia. <i>Marine Biology</i> , 1994, 119, 145-149.	0.7	25



#	ARTICLE	IF	CITATIONS
127	Patterns in species richness, size, and latitudinal range of East Atlantic fishes. <i>Ecography</i> , 1994, 17, 242-248.	2.1	92
128	<i>Paralomis Phrixa</i> (Decapoda, Anomura, Lithodidae), a New Species From Northern Peru, and a Key To the Eastern Pacific Species of the Genus. <i>Crustaceana</i> , 1992, 63, 313-317.	0.1	4
129	Epibiosis and Rhizocephalan Infestation Patterns in Relation to the Reproductive Biology of <i>Lithodes ferox</i> (Filhol, 1885) (Anomura: Lithodidae). <i>Journal of Crustacean Biology</i> , 1992, 12, 561.	0.3	28
130	Trends in the demersal fish community off Namibia from 1983 to 1990. <i>African Journal of Marine Science</i> , 1992, 12, 635-649.	0.6	18
131	Distribution Patterns and Migration of <i>Lithodes ferox</i> (Filhol) (Anomura: Lithodidae) off Namibia. <i>Journal of Crustacean Biology</i> , 1991, 11, 261.	0.3	10
132	Diurnal variation in the feeding activity and catch rate of cape hake ( <i>Merluccius capensis</i> and <i>M.</i> ) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 5</i>	0.9	23
133	Biogeography and Community Structure of the Decapod Crustacean Fauna Off Namibia (Southeast) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 5</i>	0.3	37
134	Relationship between measurements of hake biomass and sea surface temperature off southern Namibia. <i>African Journal of Marine Science</i> , 1991, 10, 213-217.	0.6	14
135	Influence of Benguela upwelling on the structure of demersal fish populations off Namibia. <i>Marine Biology</i> , 1990, 104, 175-182.	0.7	36
136	Food selection by a sit-and-wait predator, the monkfish, <i>Lophius upsicephalus</i> , off Namibia (South) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 5</i>	0.4	11
137	The validation of <i>Palaemon peringueyi</i> (Stebbing, 1915) from Southern African waters and its relationship with <i>Palaemon pacificus</i> (Stimpson, 1860) (Decapoda, Palaemonidae). <i>Journal of Natural History</i> , 1990, 24, 627-633.	0.2	7
138	Les espèces atlantiques du genre <i>Eumunida</i> Smith, 1883 (Crustacea: Decapoda: Chirostylidae). <i>Journal of Natural History</i> , 1990, 24, 647-666.	0.2	14
139	Influence of environmental conditions on the distribution of <i>Pterygosquilla armata capensis</i> (Crustacea: Stomatopoda) off Namibia. <i>African Journal of Marine Science</i> , 1990, 9, 169-175.	0.6	3
140	The Identity of <i>Xaiva Pulchella</i> Macleay, 1838 (Decapoda, Portunidae). <i>Crustaceana</i> , 1989, 57, 107-110.	0.1	1
141	Gastric emptying in <i>Scyllorhinus canicula</i> (L.): a comparison of surface-dependent and non-surface dependent models. <i>Journal of Fish Biology</i> , 1989, 35, 37-48.	0.7	20
142	Distribution of <i>Bathynectes piperitus</i> (Brachyura: Portunidae) in the Benguela Upwelling Region and Its Relationship with Some Environmental Parameters. <i>Journal of Crustacean Biology</i> , 1989, 9, 373.	0.3	2
143	Three new species of <i>Paralomis</i> (Crustacea, Decapoda, Anomura, Lithodidae) from the Pacific and Antarctic Oceans. <i>Zoologica Scripta</i> , 1988, 17, 69-75.	0.7	22
144	Feeding of <i>Merluccius capensis</i> and <i>M. paradoxus</i> off Namibia. <i>African Journal of Marine Science</i> , 1988, 6, 227-243.	0.6	35

#	ARTICLE	IF	CITATIONS
145	Trophic relationships in the demersal fish community off Namibia. African Journal of Marine Science, 1987, 5, 585-596.	0.6	62
146	CVPA, an expanded VPA with cannibalism. Application to a hake population. Fisheries Research, 1985, 3, 61-79.	0.9	18
147	Feeding pattern of the kingklip ( <i>Genypterus capensis</i> ) and its effect on the hake ( <i>Merluccius capensis</i> ) resource off the coast of Namibia. Marine Biology, 1983, 78, 105-112.	0.7	12
148	A New Species of <i>Paralomis</i> (Decapoda Anomura) From the Southeastern Atlantic. Crustaceana, 1982, 43, 142-146.	0.1	4
149	Ecological overlap between macrourids in the western mediterranean sea. Marine Biology, 1979, 53, 149-159.	0.7	75
150	On the Occurrence of <i>Richardina Fredericii</i> Lo Bianco, 1903 (Decapoda, Stenopodidae) in Spanish Waters. Crustaceana, 1978, 35, 107-109.	0.1	2