Agnieszka SÅ,owik

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/950491/publications.pdf

Version: 2024-02-01

111 2,288 19
papers citations h-index

113 113 5104 all docs docs citations times ranked citing authors

40

g-index

#	Article	IF	CITATIONS
1	Multi-ethnic genome-wide association study for atrial fibrillation. Nature Genetics, 2018, 50, 1225-1233.	21.4	552
2	Meta-analysis of Genome-wide Association Studies Identifies 1q22 as a Susceptibility Locus for Intracerebral Hemorrhage. American Journal of Human Genetics, 2014, 94, 511-521.	6.2	235
3	Genome-wide association study of intracranial aneurysms identifies 17 risk loci and genetic overlap with clinical risk factors. Nature Genetics, 2020, 52, 1303-1313.	21.4	163
4	Genome-wide association study of cerebral small vessel disease reveals established and novel loci. Brain, 2019, 142, 3176-3189.	7.6	76
5	Big Data Approaches to Phenotyping Acute Ischemic Stroke Using Automated Lesion Segmentation of Multi-Center Magnetic Resonance Imaging Data. Stroke, 2019, 50, 1734-1741.	2.0	52
6	Outcome after acute ischemic stroke is linked to sex-specific lesion patterns. Nature Communications, 2021, 12, 3289.	12.8	50
7	White matter hyperintensity quantification in large-scale clinical acute ischemic stroke cohorts – The MRI-GENIE study. Neurolmage: Clinical, 2019, 23, 101884.	2.7	48
8	Reduced ex vivo release of pro-inflammatory cytokines and elevated plasma interleukin-6 are inflammatory signatures of post-stroke delirium. Journal of Neuroinflammation, 2018, 15, 111.	7.2	43
9	Association of Apolipoprotein E With Intracerebral Hemorrhage Risk by Race/Ethnicity. JAMA Neurology, 2019, 76, 480.	9.0	43
10	Early Neurological Change After Ischemic Stroke Is Associated With 90-Day Outcome. Stroke, 2021, 52, 132-141.	2.0	36
11	Genetically Elevated <scp>LDL</scp> Associates with Lower Risk of Intracerebral Hemorrhage. Annals of Neurology, 2020, 88, 56-66.	5.3	35
12	Genetic variants in CETP increase risk of intracerebral hemorrhage. Annals of Neurology, 2016, 80, 730-740.	5.3	33
13	Reduction in Cerebrospinal Fluid Volume as an Early Quantitative Biomarker of Cerebral Edema After Ischemic Stroke. Stroke, 2020, 51, 462-467.	2.0	33
14	Plasma endotoxin activity rises during ischemic stroke and is associated with worse short-term outcome. Journal of Neuroimmunology, 2016, 297, 76-80.	2.3	30
15	FTIR, Raman and AFM characterization of the clinically valid biochemical parameters of the thrombi in acute ischemic stroke. Scientific Reports, 2019, 9, 15475.	3.3	27
16	PRospective Observational POLIsh Study on post-stroke delirium (PROPOLIS): methodology of hospital-based cohort study on delirium prevalence, predictors and diagnostic tools. BMC Neurology, 2015, 15, 94.	1.8	26
17	Repetitive transcranial magnetic stimulation for the treatment of cognitive impairment in frontotemporal dementia: an open-label pilot study. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 749-755.	2.2	26
18	Co-medication and potential drug interactions among patients with epilepsy. Seizure: the Journal of the British Epilepsy Association, 2019, 66, 47-52.	2.0	25

#	Article	IF	CITATIONS
19	Clinical course and outcome of SARS-CoV-2 infection in multiple sclerosis patients treated with disease-modifying therapies — the Polish experience. Neurologia I Neurochirurgia Polska, 2021, 55, 212-222.	1.2	24
20	Decompensated Heart Failure Is a Strong Independent Predictor of Functional Outcome After Ischemic Stroke. Journal of Cardiac Failure, 2015, 21, 642-646.	1.7	21
21	Gender-Related Differences in Diagnostic Evaluation and Outcome of Ischemic Stroke in Poland. Stroke, 2009, 40, 980-982.	2.0	20
22	Why do some women with epilepsy use valproic acid despite current guidelines? A single-center cohort study. Epilepsy and Behavior, 2019, 98, 1-5.	1.7	20
23	The association between plasma endotoxin, endotoxin pathway proteins and outcome after ischemic stroke. Atherosclerosis, 2018, 269, 138-143.	0.8	19
24	Association between cardiovascular disease, cardiovascular drug therapy, and in-hospital outcomes in patients with COVID-19: data from a large single-center registry in Poland. Kardiologia Polska, 2021, 79, 773-780.	0.6	19
25	Brain Volume: An Important Determinant of Functional Outcome After Acute Ischemic Stroke. Mayo Clinic Proceedings, 2020, 95, 955-965.	3.0	18
26	COVID-19 among patients with epilepsy: Risk factors and course of the disease. Epilepsy and Behavior, 2021, 120, 107996.	1.7	18
27	Systemic response to rupture of intracranial aneurysms involves expression of specific gene isoforms. Journal of Translational Medicine, 2019, 17, 141.	4.4	17
28	Overlap in the Genetic Architecture of Stroke Risk, Early Neurological Changes, and Cardiovascular Risk Factors. Stroke, 2019, 50, 1339-1345.	2.0	17
29	Inflammatory Responses Induced by the Rupture of Intracranial Aneurysms Are Modulated by miRNAs. Molecular Neurobiology, 2020, 57, 988-996.	4.0	16
30	Quantitative Serial CT Imaging-Derived Features Improve Prediction of Malignant Cerebral Edema after Ischemic Stroke. Neurocritical Care, 2020, 33, 785-792.	2.4	16
31	Menstrual disorders and their determinants among women with epilepsy. Neuropsychiatric Disease and Treatment, 2018, Volume 14, 2657-2664.	2.2	15
32	Fasting Hyperglycemia and Long-term Outcome in Patients with Acute Ischemic Stroke Treated with Mechanical Thrombectomy. Journal of Stroke and Cerebrovascular Diseases, 2020, 29, 104774.	1.6	15
33	C-reactive protein and post-stroke depressive symptoms. Scientific Reports, 2020, 10, 1431.	3.3	15
34	The FGA Thr312Ala polymorphism and risk of intracerebral haemorrhage in Polish and Greek populations. Neurologia I Neurochirurgia Polska, 2014, 48, 105-110.	1,2	14
35	Safety of switching from brand-name to generic levetiracetam in patients with epilepsy. Drug Design, Development and Therapy, 2017, Volume 11, 2287-2291.	4.3	14
36	Early apathetic, but not depressive, symptoms are associated with poor outcome after stroke. European Journal of Neurology, 2021, 28, 1949-1957.	3.3	14

#	Article	IF	CITATIONS
37	Hemispheric CSF volume ratio quantifies progression and severity of cerebral edema after acute hemispheric stroke. Journal of Cerebral Blood Flow and Metabolism, 2021, 41, 2907-2915.	4.3	14
38	Transcranial Magnetic Stimulation as a Diagnostic and Therapeutic Tool in Various Types of Dementia. Journal of Clinical Medicine, 2021, 10, 2875.	2.4	14
39	Catalase activity in blood fractions of patients with sporadic ALS. Pharmacological Reports, 2014, 66, 704-707.	3.3	13
40	Can Prediction of Functional Outcome after Stroke Be Improved by Adding Fibrinogen to Prognostic Model?. Journal of Stroke and Cerebrovascular Diseases, 2016, 25, 2752-2755.	1.6	13
41	Prognostic Significance of Stroke-Associated Infection and other Readily Available Parameters in Acute Ischemic Stroke Treated by Intravenous Thrombolysis. Journal of Stroke and Cerebrovascular Diseases, 2021, 30, 105525.	1.6	13
42	The Need for New Biomarkers to Assist with Stroke Prevention and Prediction of Post-Stroke Therapy Based on Plasma-Derived Extracellular Vesicles. Biomedicines, 2021, 9, 1226.	3.2	13
43	Genetic variants influencing elevated myeloperoxidase levels increase risk of stroke. Brain, 2017, 140, 2663-2672.	7.6	12
44	Psychiatric comedication in patients with epilepsy. Epilepsy and Behavior, 2018, 83, 207-211.	1.7	12
45	Clinical Relevance of Changes in Peripheral Blood Cells After Intracranial Aneurysm Rupture. Journal of Stroke and Cerebrovascular Diseases, 2020, 29, 105293.	1.6	12
46	MRI Radiomic Signature of White Matter Hyperintensities Is Associated With Clinical Phenotypes. Frontiers in Neuroscience, 2021, 15, 691244.	2.8	12
47	Excessive White Matter Hyperintensity Increases Susceptibility to Poor Functional Outcomes After Acute Ischemic Stroke. Frontiers in Neurology, 2021, 12, 700616.	2.4	11
48	Reduced release of TNF \hat{l} ± and IP-10 after ex vivo blood stimulation with endotoxin is associated with poor outcome after stroke. Cytokine, 2018, 102, 51-54.	3.2	10
49	Implementation of the new ILAE classification of epilepsies into clinical practice $\hat{a} \in \text{``}$ A cohort study. Epilepsy and Behavior, 2019, 96, 28-32.	1.7	10
50	Single nucleotide variations in <i>ZBTB46</i> are associated with post-thrombolytic parenchymal haematoma. Brain, 2021, 144, 2416-2426.	7.6	10
51	Multiple sclerosis incidence and prevalence in Poland: Data from administrative health claims. Multiple Sclerosis and Related Disorders, 2021, 55, 103162.	2.0	10
52	The AGTR1 gene A1166C polymorphism as a risk factor and outcome predictor of primary intracerebral and aneurysmal subarachnoid hemorrhages. Neurologia I Neurochirurgia Polska, 2014, 48, 242-247.	1.2	9
53	Brand-to-generic oxcarbazepine switch $\hat{a}\in$ A prospective observational study. Epilepsy Research, 2019, 151, 75-77.	1.6	9
54	Toll-like receptor 4-mediated cytokine synthesis and post-stroke depressive symptoms. Translational Psychiatry, 2021, 11, 246.	4.8	9

#	Article	IF	CITATIONS
55	Accelerating Prediction of Malignant Cerebral Edema After Ischemic Stroke with Automated Image Analysis and Explainable Neural Networks. Neurocritical Care, 2022, 36, 471-482.	2.4	9
56	Anthracycline-induced cardiotoxicity prevention with angiotensin-converting enzyme inhibitor ramipril in women with low-risk breast cancer: results of a prospective randomized study. Kardiologia Polska, 2020, 78, 131-137.	0.6	9
57	Symptoms after COVID-19 Infection in Individuals with Multiple Sclerosis in Poland. Journal of Clinical Medicine, 2021, 10, 5225.	2.4	9
58	Dynamic cerebral autoregulation is compromised in ischaemic stroke of undetermined aetiology only in the non-affected hemisphere. Neurologia I Neurochirurgia Polska, 2014, 48, 91-97.	1.2	8
59	Genetics of the thrombomodulin-endothelial cell protein C receptor system and the risk of early-onset ischemic stroke. PLoS ONE, 2018, 13, e0206554.	2.5	8
60	Treatment of high- and intermediate-risk pulmonary embolism using the AngioJet percutaneous mechanical thrombectomy system in patients with contraindications for thrombolytic treatment – a pilot study. Wideochirurgia I Inne Techniki Maloinwazyjne, 2018, 13, 233-242.	0.7	8
61	Use of complementary and alternative medicine among adults with epilepsy in a university epilepsy clinic in Poland. Epilepsy and Behavior, 2019, 98, 40-44.	1.7	8
62	Association of early and later depressive symptoms with functional outcome after ischemic stroke. Journal of Neural Transmission, 2021, 128, 679-686.	2.8	8
63	Sex-specific lesion pattern of functional outcomes after stroke. Brain Communications, 2022, 4, fcac020.	3.3	8
64	Delirium and subsyndromal delirium are associated with the long-term risk of death after ischaemic stroke. Aging Clinical and Experimental Research, 2022, 34, 1459-1462.	2.9	8
65	Improvement of survival in Polish stroke patients is related to reduced stroke severity and better control of risk factors: the Krakow Stroke Database. Archives of Medical Science, 2016, 3, 552-555.	0.9	7
66	Common allergies do not influence the prevalence of cutaneous hypersensitivity reactions to antiepileptic drugs. Epilepsy Research, 2017, 135, 9-13.	1.6	7
67	The specific ex vivo released cytokine profile is associated with ischemic stroke outcome and improves its prediction. Journal of Neuroinflammation, 2020, 17, 7.	7.2	7
68	Mean platelet volume and its genetic variants relate to stroke severity and 1-year mortality. Neurology, 2020, 95, e1153-e1162.	1.1	7
69	Safety of Vaccines against SARS-CoV-2 among Polish Patients with Multiple Sclerosis Treated with Disease-Modifying Therapies. Vaccines, 2022, 10, 763.	4.4	7
70	RP11-362K2.2:RP11-767I20.1 Genetic Variation Is Associated with Post-Reperfusion Therapy Parenchymal Hematoma. A GWAS Meta-Analysis. Journal of Clinical Medicine, 2021, 10, 3137.	2.4	6
71	<p>Status epilepticus in patients with genetic (idiopathic) generalized epilepsy</p> . Neuropsychiatric Disease and Treatment, 2019, Volume 15, 1585-1592.	2.2	5
72	Diffusion-Weighted Imaging, MR Angiography, and Baseline Data in a Systematic Multicenter Analysis of 3,301 MRI Scans of Ischemic Stroke Patients—Neuroradiological Review Within the MRI-GENIE Study. Frontiers in Neurology, 2020, 11, 577.	2.4	5

#	Article	IF	CITATIONS
73	Magnetisation transfer imaging revealed microstructural changes related to apathy symptoms after ischaemic stroke. International Journal of Geriatric Psychiatry, 2021, 36, 1264-1273.	2.7	5
74	C-Reactive Protein and White Blood Cell Count in Non-Infective Acute Ischemic Stroke Patients Treated with Intravenous Thrombolysis. Journal of Clinical Medicine, 2021, 10, 1610.	2.4	5
75	Gender-Related Differences in Prodromal Multiple Sclerosis Characteristics: A 7-Year Observation Study. Journal of Clinical Medicine, 2021, 10, 3821.	2.4	5
76	Association between C677T polymorphism of MTHFR gene and risk of amyotrophic lateral sclerosis: Polish population study and a meta-analysis. Neurologia I Neurochirurgia Polska, 2017, 51, 135-139.	1.2	4
77	Real-World Comparison of Human and Software Image Assessment in Acute Ischemic Stroke Patients' Qualification for Reperfusion Treatment. Journal of Clinical Medicine, 2020, 9, 3383.	2.4	4
78	Glucocorticoid Resistance is Associated with Poor Functional Outcome After Stroke. Cellular and Molecular Neurobiology, 2020, 40, 1321-1326.	3.3	4
79	Elevated plasma levels of galectin-3 binding protein are associated with post-stroke delirium – A pilot study. Journal of Neuroimmunology, 2021, 356, 577579.	2.3	4
80	Recommendations of the Polish Medical Society of Radiology and the Polish Society of Neurology for a protocol concerning routinely used magnetic resonance imaging in patients with multiple sclerosis. Neurologia I Neurochirurgia Polska, 2020, 54, 410-415.	1.2	4
81	Post-stroke infection in acute ischemic stroke patients treated with mechanical thrombectomy does not affect long-term outcome. Postepy W Kardiologii Interwencyjnej, 2020, 16, 452-459.	0.2	4
82	Predictors of remission in patients with epilepsy. Neurologia I Neurochirurgia Polska, 2020, 54, 434-439.	1.2	4
83	Acute-onset chorea and confusional state in 77-year-old COVID-19 patient: a case report. Neurologia I Neurochirurgia Polska, 2022, 56, 106-110.	1.2	4
84	Body mass index and aromatase inhibitors: a step forward in individualizing therapy for breast cancer patients?. Expert Review of Anticancer Therapy, 2016, 16, 759-766.	2.4	3
85	Genetic testing and the phenotype of Polish patients with Unverricht–Lundborg disease (EPM1) — A cohort study. Epilepsy and Behavior, 2020, 112, 107439.	1.7	3
86	Olfactory Dysfunction in Patients With Relapsing-Remitting Multiple Sclerosis Treated With Disease-Modifying Therapies. Ear, Nose and Throat Journal, 2020, , 014556132097377.	0.8	3
87	Clinical utility of brain computed tomography in prediction of post-stroke delirium. Journal of Neural Transmission, 2021, 128, 207-213.	2.8	3
88	Lacosamide and myoclonic seizures: what is the risk of aggravation?. Neurologia I Neurochirurgia Polska, 2021, 55, 107-109.	1.2	3
89	Lipopolysaccharide binding protein and sCD14 as risk markers of stroke-associated pneumonia. Journal of Neuroimmunology, 2021, 354, 577532.	2.3	3
90	Subtypes of delirium after ischaemic strokeâ€"predisposing factors and outcomes: a prospective observational study (PROPOLIS). European Journal of Neurology, 2022, 29, 478-485.	3.3	3

#	Article	IF	Citations
91	Somatic comorbidity in Polish patients with epilepsy. Polish Archives of Internal Medicine, 2019, 129, 303-307.	0.4	3
92	Obstetric and neonatal outcomes in women with epilepsy in Poland — a two-centre study. Neurologia I Neurochirurgia Polska, 2020, 54, 62-65.	1.2	3
93	Seasonal Variations in the Concentration of Particulate Matter in the Air of Cracow Affect the Magnitude of CD4+ T Cell Subsets Cytokine Production in Patients with Inflammatory and Autoimmune Disorders. Atmosphere, 2022, 13, 529.	2.3	3
94	Two novel mutations of the SETX gene and ataxia with oculomotor apraxia type 2. Clinical Neurology and Neurosurgery, 2015, 128, 44-46.	1.4	2
95	Endovascular management of carotid and vertebral artery dissections with new generation double-mesh stent and protection systems – single-center early and midterm results. Postepy W Kardiologii Interwencyjnej, 2019, 15, 321-327.	0.2	2
96	Outcome estimation based on multimodal computed tomography examination in acute ischaemic stroke patients treated with mechanical thrombectomy. Wideochirurgia I Inne Techniki Maloinwazyjne, 2019, 14, 560-566.	0.7	2
97	Ex vivo synthesized cytokines as a biomarker of stroke-associated pneumonia. Clinica Chimica Acta, 2020, 510, 260-263.	1.1	2
98	Opposite regulation of piRNAs, rRNAs and miRNAs in the blood after subarachnoid hemorrhage. Journal of Molecular Medicine, 2020, 98, 887-896.	3.9	2
99	Fasting Normoglycemia after Intravenous Thrombolysis Predicts Favorable Long-Term Outcome in Non-Diabetic Patients with Acute Ischemic Stroke. Journal of Clinical Medicine, 2021, 10, 3005.	2.4	2
100	Additional Value of [18F]FDG PET/CT in Detection of Suspected Malignancy in Patients with Paraneoplastic Neurological Syndromes Having Negative Results of Conventional Radiological Imaging. Journal of Clinical Medicine, 2022, 11, 1537.	2.4	2
101	The use of anticholinergic medication is associated with an increased risk of stroke-associated pneumonia. Aging Clinical and Experimental Research, 2022, 34, 1935-1938.	2.9	2
102	Left ventricular geometry and white matter lesions in ischemic stroke patients. Blood Pressure, 2016, 25, 149-154.	1.5	1
103	4C Mortality Score correlates with in-hospital functional outcome after COVID-19-associated ischaemic stroke. Neurologia I Neurochirurgia Polska, 2021, 55, 295-299.	1.2	1
104	Statement by a Working Group conceived by the Polish National Consultants in Cardiology and Neurology addressing the use of implantable cardiac monitors in patients after ischaemic embolic stroke of undetermined source. Neurologia I Neurochirurgia Polska, 2019, 53, 181-189.	1.2	1
105	Prognostic significance of age in patients with acute ischaemic stroke treated with intravenous thrombolysis. Neurologia I Neurochirurgia Polska, 2022, , .	1.2	1
106	Mechanical thrombectomy in COVID-19-associated ischaemic stroke: patient characteristics and outcomes in a single-centre study. Neurologia I Neurochirurgia Polska, 2022, , .	1.2	1
107	Paraoxonase 2 C311S single nucleotide polymorphism is associated with type C lesions in coronary atherosclerosis. Clinical Biochemistry, 2022, 105-106, 64-69.	1.9	1
108	Reply to "Depression among patients with epilepsy — A diagnostic approach― Epilepsy and Behavior, 2018, 88, 397.	1.7	0

#	Article	IF	CITATIONS
109	Use of dietary and herbal supplements in adult patients with epilepsy. Epilepsy Research, 2019, 156, 106168.	1.6	0
110	The prognostic significance of large vessel occlusion in stroke patients treated by intravenous thrombolysis. Polish Journal of Radiology, 2021, 86, 344-352.	0.9	0
111	Recommendations of the Polish Medical Society of Radiology and the Polish Society of Neurology for a protocol concerning routinely used magnetic resonance imaging in patients with multiple sclerosis. Polish Journal of Radiology, 2020, 85, 272-276.	0.9	0