

Ernesto Akio Taketomi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/9505376/ernesto-akio-taketomi-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

16

papers

260

citations

11

h-index

16

g-index

21

ext. papers

283

ext. citations

3.2

avg, IF

1.97

L-index

#	Paper	IF	Citations
16	IgE, IgG1, and IgG4 Reactivity to Glycosylated Extract in Allergic Patients. <i>BioMed Research International</i> , 2019, 2019, 9840890	3	1
15	Allergen-Specific Immunotherapy Follow-Up by Measuring Allergen-Specific IgG as an Objective Parameter 2017,		1
14	Allergen Microarray Indicates Pooidae Sensitization in Brazilian Grass Pollen Allergic Patients. <i>PLoS ONE</i> , 2015, 10, e0128402	3.7	4
13	Reply: To PMID 24299565. <i>Pediatric Allergy and Immunology</i> , 2014, 25, 417-8		4.2
12	Modulation of mucosal/systemic antibody response after sublingual immunotherapy in mite-allergic children. <i>Pediatric Allergy and Immunology</i> , 2013, 24, 752-61	4.2	15
11	Allergen-specific IgG antibodies purified from mite-allergic patients sera block the IgE recognition of Dermatophagoides pteronyssinus antigens: an in vitro study. <i>Clinical and Developmental Immunology</i> , 2013, 2013, 657424		16
10	Serum and salivary IgE, IgA, and IgG4 antibodies to Dermatophagoides pteronyssinus and its major allergens, Der p1 and Der p2, in allergic and nonallergic children. <i>Clinical and Developmental Immunology</i> , 2011, 2011, 302739		28
9	Antibody and cytokine responses to house dust mite allergens and Toxoplasma gondii antigens in atopic and non-atopic Brazilian subjects. <i>Clinical Immunology</i> , 2010, 136, 148-56	9	18
8	IgG1, IgG4, and IgE antibody responses in human strongyloidiasis by ELISA using <i>Strongyloides ratti</i> saline extract as heterologous antigen. <i>Parasitology Research</i> , 2007, 101, 1209-14	2.4	43
7	Doen�a al『rgica pol『ica: polens al『rgenos e seus principais al『genos. <i>Revista Brasileira De Otorrinolaringologia</i> , 2006, 72, 562-567		10
6	Strongyloides ratti antigenic components recognized by IgE antibodies in immunoblotting as an additional tool for improving the immunodiagnosis in human strongyloidiasis. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2004, 99, 89-93	2.6	18
5	Mite and cat allergen exposure in Brazilian public transport vehicles. <i>Annals of Allergy, Asthma and Immunology</i> , 2004, 93, 179-84	3.2	13
4	Heterologous antigen extract in ELISA for the detection of human IgE anti-Strongyloides stercoralis. <i>Revista Do Instituto De Medicina Tropical De Sao Paulo</i> , 2003, 45, 265-8	2.2	13
3	A sensitive reverse ELISA for the measurement of specific IgE to Der p 2, a major Dermatophagoides pteronyssinus allergen. <i>Annals of Allergy, Asthma and Immunology</i> , 2001, 86, 545-50	3.2	12
2	Dermatophagoides farinae (Der f 1) and Dermatophagoides pteronyssinus (Der p 1) allergen exposure among subjects living in Uberl�ndia, Brazil. <i>International Archives of Allergy and Immunology</i> , 2000, 122, 257-63	3.7	22
1	Recombinant allergens for immunotherapy: a Der p 2 variant with reduced IgE reactivity retains T-cell epitopes. <i>Journal of Allergy and Clinical Immunology</i> , 1998, 101, 423-5	11.5	33