

Shigeto Sudo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9506990/publications.pdf>

Version: 2024-02-01

41
papers

1,598
citations

331259

21
h-index

301761

39
g-index

42
all docs

42
docs citations

42
times ranked

1891
citing authors

#	ARTICLE	IF	CITATIONS
1	Influence of Azolla incorporation and/or dual cropping on CH ₄ and N ₂ O emission from a paddy field. <i>Soil Science and Plant Nutrition</i> , 2022, 68, 246-255.	0.8	2
2	Effect of Paddy-Upland Rotation System on the Net Greenhouse Gas Balance as the Sum of Methane and Nitrous Oxide Emissions and Soil Carbon Storage: A Case in Western Japan. <i>Agriculture (Switzerland)</i> , 2021, 11, 52.	1.4	7
3	Effects of controlled release N fertilizers and reduced application rate on nitrous oxide emissions from soybean fields converted from rice paddies. <i>Soil Science and Plant Nutrition</i> , 2021, 67, 716-726.	0.8	4
4	Assessment of crop residue and palm shell biochar incorporation on greenhouse gas emissions during the fallow and crop growing seasons of broccoli (<i>Brassica oleracea</i> var. <i>italica</i>). <i>Soil and Tillage Research</i> , 2020, 196, 104435.	2.6	21
5	Azolla incorporation and dual cropping influences CH ₄ and N ₂ O emissions from flooded paddy ecosystems. <i>Soil Science and Plant Nutrition</i> , 2020, 66, 152-162.	0.8	12
6	Field Validation of the DNDC-Rice Model for Methane and Nitrous Oxide Emissions from Double-Cropping Paddy Rice under Different Irrigation Practices in Tamil Nadu, India. <i>Agriculture (Switzerland)</i> , 2020, 10, 355.	1.4	9
7	Influence of pruning waste biochar and oyster shell on N ₂ O and CO ₂ emissions from Japanese pear orchard soil. <i>Heliyon</i> , 2018, 4, e00568.	1.4	12
8	Forage rice varieties Fukuhibiki and Tachisuzuka emit larger CH ₄ than edible rice Haenuki. <i>Soil Science and Plant Nutrition</i> , 2018, 64, 77-83.	0.8	23
9	Azolla cover significantly decreased CH ₄ but not N ₂ O emissions from flooding rice paddy to atmosphere. <i>Soil Science and Plant Nutrition</i> , 2018, 64, 68-76.	0.8	20
10	Methane and nitrous oxide emissions from conventional and modified rice cultivation systems in South India. <i>Agriculture, Ecosystems and Environment</i> , 2018, 252, 148-158.	2.5	88
11	Aerated Irrigation and Pruning Residue Biochar on N ₂ O Emission, Yield and Ion Uptake of Komatsuna. <i>Horticulturae</i> , 2018, 4, 33.	1.2	1
12	Mitigation Potential and Yield-Scaled Global Warming Potential of Early-Season Drainage from a Rice Paddy in Tamil Nadu, India. <i>Agronomy</i> , 2018, 8, 202.	1.3	19
13	Returning Tea Pruning Residue and Its Biochar Had a Contrasting Effect on Soil N ₂ O and CO ₂ Emissions from Tea Plantation Soil. <i>Atmosphere</i> , 2018, 9, 109.	1.0	11
14	Effect of dolomite and biochar addition on N ₂ O and CO ₂ emissions from acidic tea field soil. <i>PLoS ONE</i> , 2018, 13, e0192235.	1.1	46
15	Greenhouse gas emissions, soil carbon sequestration and crop yields in a rain-fed rice field with crop rotation management. <i>Agriculture, Ecosystems and Environment</i> , 2017, 237, 109-120.	2.5	52
16	Development of a method for estimating total CH ₄ emission from rice paddies in Japan using the DNDC-Rice model. <i>Science of the Total Environment</i> , 2016, 547, 429-440.	3.9	36
17	Multi-site monitoring for N ₂ O emission factors of synthetic fertilizer in various soils with different redoximorphic features across Japan. <i>Nutrient Cycling in Agroecosystems</i> , 2015, 103, 87-99.	1.1	4
18	Differences in the Spatial Variability Among CO ₂ , CH ₄ , and N ₂ O Gas Fluxes from an Urban Forest Soil in Japan. <i>Ambio</i> , 2015, 44, 55-66.	2.8	12

#	ARTICLE	IF	CITATIONS
19	Potential of prolonged midseason drainage for reducing methane emission from rice paddies in Japan: a long-term simulation using the DNDC-Rice model. <i>Biology and Fertility of Soils</i> , 2014, 50, 879-889.	2.3	39
20	Trace gas and particle emissions from open burning of three cereal crop residues: Increase in residue moistness enhances emissions of carbon monoxide, methane, and particulate organic carbon. <i>Atmospheric Environment</i> , 2014, 95, 36-44.	1.9	74
21	Nitrification, ammonia-oxidizing communities, and N ₂ O and CH ₄ fluxes in an imperfectly drained agricultural field fertilized with coated urea with and without dicyandiamide. <i>Biology and Fertility of Soils</i> , 2013, 49, 213-223.	2.3	83
22	Evaluation of uncertainties in N ₂ O and NO fluxes from agricultural soil using a hierarchical Bayesian model. <i>Journal of Geophysical Research</i> , 2012, 117, .	3.3	14
23	Characterization and production and consumption processes of N ₂ O emitted from temperate agricultural soils determined via isotopomer ratio analysis. <i>Global Biogeochemical Cycles</i> , 2011, 25, n/a-n/a.	1.9	123
24	Combined emission of CH ₄ and N ₂ O from a paddy field was reduced by preceding upland crop cultivation. <i>Soil Science and Plant Nutrition</i> , 2011, 57, 167-178.	0.8	44
25	Mitigation of methane emissions from paddy fields by prolonging midseason drainage. <i>Agriculture, Ecosystems and Environment</i> , 2011, 141, 359-372.	2.5	122
26	N ₂ O and NO emissions from an Andisol field as influenced by pelleted poultry manure. <i>Soil Biology and Biochemistry</i> , 2009, 41, 521-529.	4.2	107
27	Automated sampling system for long-term monitoring of nitrous oxide and methane fluxes from soils. <i>Soil Science and Plant Nutrition</i> , 2009, 55, 435-440.	0.8	17
28	Effect of land use change from paddy rice cultivation to upland crop cultivation on soil carbon budget of a cropland in Japan. <i>Agriculture, Ecosystems and Environment</i> , 2008, 125, 9-20.	2.5	116
29	An Empirical Model of Soil Chemical Properties that Regulate Methane Production in Japanese Rice Paddy Soils. <i>Journal of Environmental Quality</i> , 2007, 36, 1920-1925.	1.0	37
30	Time-lagged induction of N ₂ O emission and its trade-off with NO emission from a nitrogen fertilized Andisol. <i>Soil Science and Plant Nutrition</i> , 2007, 53, 362-372.	0.8	7
31	Hydrogen and carbon isotopic measurements of methane from agricultural combustion: Implications for isotopic signatures of global biomass burning sources. <i>Journal of Geophysical Research</i> , 2006, 111, .	3.3	9
32	Temporal and Spatial Variations in N ₂ O Emissions from a Chinese Cabbage Field as a Function of Type of Fertilizer and Application. <i>Nutrient Cycling in Agroecosystems</i> , 2006, 74, 147-155.	1.1	22
33	Continuous, Automated Nitrous Oxide Measurements from Paddy Soils Converted to Upland Crops. <i>Soil Science Society of America Journal</i> , 2005, 69, 1977-1986.	1.2	41
34	Development of a System for Simultaneous and Continuous Measurement of Carbon Dioxide, Methane and Nitrous Oxide Fluxes from Croplands Based on the Automated Closed Chamber Method. <i>Soil Science and Plant Nutrition</i> , 2005, 51, 557-564.	0.8	27
35	The variation of greenhouse gas emissions from soils of various land-use/cover types in Jambi province, Indonesia. <i>Nutrient Cycling in Agroecosystems</i> , 2005, 71, 17-32.	1.1	74
36	Spatial patterns of greenhouse gas emission in a tropical rainforest in Indonesia. <i>Nutrient Cycling in Agroecosystems</i> , 2005, 71, 55-62.	1.1	46

#	ARTICLE	IF	CITATIONS
37	Methane and nitrous oxide emissions from a paddy field with Japanese conventional water management and fertilizer application. <i>Global Biogeochemical Cycles</i> , 2004, 18, n/a-n/a.	1.9	119
38	Annual and El Niño-Southern Oscillation variations in observations of in situ stratospheric ozone over Peninsular Malaysia. <i>Journal of Geophysical Research</i> , 2002, 107, ACH 12-1.	3.3	3
39	Title is missing!. <i>Nutrient Cycling in Agroecosystems</i> , 2002, 63, 231-238.	1.1	35
40	Reactivity study of alloy clusters made of aluminum and some transition metals with hydrogen. <i>Chemical Physics Letters</i> , 1989, 164, 427-432.	1.2	59
41	Effects of a new compost-chemical fertilizer mixture on CO ₂ , and N ₂ O production and plant growth in a Chernozem and an Andosol. <i>Soil Science and Plant Nutrition</i> , 0, , 1-8.	0.8	1