

Maria Swennergren Hansen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9514430/publications.pdf>

Version: 2024-02-01

19
papers

173
citations

1307594

7
h-index

1125743

13
g-index

21
all docs

21
docs citations

21
times ranked

133
citing authors

#	ARTICLE	IF	CITATIONS
1	Estimation of Patient Acceptable Symptom State (PASS) and Treatment Failure (TF) Threshold Values for the Achilles Tendon Total Rupture Score (ATRS) at 6 Months, 1 Year, and 2 Years After Acute Achilles Tendon Rupture. <i>Journal of Foot and Ankle Surgery</i> , 2022, 61, 503-507.	1.0	5
2	The Copenhagen Achilles Length Measurement performed at time of acute Achilles tendon rupture is correlated to tendon elongation after one year.. <i>Journal of Foot and Ankle Surgery</i> , 2022, , .	1.0	3
3	The Achilles Tendon Length Measure and the Achilles Tendon Resting Angle show acceptable construct validity using the Copenhagen Achilles Length Measure as gold standard. <i>Foot and Ankle Surgery</i> , 2021, 27, 655-659.	1.7	6
4	Diabetes and treatment with orally administrated corticosteroids negatively affect treatment outcome at follow-up after acute Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 1584-1592.	4.2	5
5	The heel-rise work test overestimates the performed work with 21â€“25% after an Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 1604-1611.	4.2	5
6	No clinically relevant difference between operative and non-operative treatment in tendon elongation measured with the Achilles tendon resting angle (ATRA) 1Åyear after acute Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 1617-1626.	4.2	6
7	Neither heel-rise Height (HRH) nor Achilles tendon resting angle (ATRA) show strong correlations to patient limitations and return to previous activities one year after acute Achilles tendon rupture. <i>Foot and Ankle Surgery</i> , 2021, , .	1.7	3
8	Efficacy of early controlled motion of the ankle compared with immobilisation in non-operative treatment of patients with an acute Achilles tendon rupture: an assessor-blinded, randomised controlled trial. <i>British Journal of Sports Medicine</i> , 2020, 54, 719-724.	6.7	37
9	Reliability of the Copenhagen Achilles length measure (CALM) on patients with an Achilles tendon rupture. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 281-290.	4.2	11
10	Outcome after acute Achilles tendon rupture is not negatively affected by female sex and age over 65Åyears. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2020, 28, 3994-4002.	4.2	8
11	Individualized treatment for acute Achilles tendon rupture based on the Copenhagen Achilles Rupture Treatment Algorithm (CARTA): a study protocol for a multicenter randomized controlled trial. <i>Trials</i> , 2020, 21, 399.	1.6	8
12	Performance of the Achilles Tendon Total Rupture Score Over Time in a Large National Database: Development of an Instruction Manual for Accurate Use. <i>American Journal of Sports Medicine</i> , 2020, 48, 1423-1429.	4.2	10
13	12â€“...Efficacy of early controlled motion of the ankle in non-operative treatment of acute achilles tendon rupture. An assessor-blinded RCT. , 2019, , .		2
14	14â€“...Individual treatment selection for acute achilles tendon rupture based on the copenhagen achilles length measurements (CALM). , 2019, , .		0
15	20â€“...The achilles tendon total rupture score should be used with caution the first 6 months after injury. , 2019, , .		0
16	21â€“...Validation of achilles tendon length measure and achilles tendon resting angle in relation to copenhagen achilles length measure. , 2019, , .		0
17	Development and reliability of the Achilles Tendon Length Measure and comparison with the Achilles Tendon Resting Angle on patients with an Achilles tendon rupture. <i>Foot and Ankle Surgery</i> , 2017, 23, 275-280.	1.7	22
18	Efficacy of early controlled motion of the ankle compared with no motion after non-operative treatment of an acute Achilles tendon rupture: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 564.	1.6	10

#	ARTICLE	IF	CITATIONS
19	Achilles tendon Total Rupture Score at 3 months can predict patients' ability to return to sport 1 year after injury. Knee Surgery, Sports Traumatology, Arthroscopy, 2016, 24, 1365-1371.	4.2	31