

Nicole Dukers

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9525428/nicole-dukers-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

142
papers

3,035
citations

30
h-index

50
g-index

149
ext. papers

3,473
ext. citations

5.2
avg, IF

5.12
L-index

#	Paper	IF	Citations
142	Homosexual men change to risky sex when perceiving less threat of HIV/AIDS since availability of highly active antiretroviral therapy: a longitudinal study. <i>Aids</i> , 2004 , 18, 303-9	3.5	188
141	Sexual risk behaviour relates to the virological and immunological improvements during highly active antiretroviral therapy in HIV-1 infection. <i>Aids</i> , 2001 , 15, 369-78	3.5	179
140	High prevalence of syphilis and other sexually transmitted diseases among sex workers in China: potential for fast spread of HIV. <i>Aids</i> , 2001 , 15, 753-9	3.5	150
139	Seroconversion for human herpesvirus 8 during HIV infection is highly predictive of Kaposi's sarcoma. <i>Aids</i> , 1998 , 12, 2481-8	3.5	143
138	Increase in sexually transmitted infections among homosexual men in Amsterdam in relation to HAART. <i>Sexually Transmitted Infections</i> , 2001 , 77, 184-6	2.8	131
137	High performance and acceptability of self-collected rectal swabs for diagnosis of Chlamydia trachomatis and Neisseria gonorrhoeae in men who have sex with men and women. <i>Sexually Transmitted Diseases</i> , 2009 , 36, 493-7	2.4	101
136	HIV incidence on the increase among homosexual men attending an Amsterdam sexually transmitted disease clinic: using a novel approach for detecting recent infections. <i>Aids</i> , 2002 , 16, F19-24	3.5	89
135	Human herpesvirus 8 infections in the Amsterdam Cohort Studies (1984-1997): analysis of seroconversions to ORF65 and ORF73. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 4838-43	11.5	69
134	Perceived viral load, but not actual HIV-1-RNA load, is associated with sexual risk behaviour among HIV-infected homosexual men. <i>Aids</i> , 2004 , 18, 1943-9	3.5	66
133	What is needed to guide testing for anorectal and pharyngeal Chlamydia trachomatis and Neisseria gonorrhoeae in women and men? Evidence and opinion. <i>BMC Infectious Diseases</i> , 2015 , 15, 533	4	61
132	High co-occurrence of anorectal chlamydia with urogenital chlamydia in women visiting an STI clinic revealed by routine universal testing in an observational study; a recommendation towards a better anorectal chlamydia control in women. <i>BMC Infectious Diseases</i> , 2014 , 14, 274	4	56
131	Lack of Epstein-Barr virus- and HIV-specific CD27- CD8+ T cells is associated with progression to viral disease in HIV-infection. <i>Aids</i> , 2002 , 16, 2001-11	3.5	55
130	Standard symptom- and sexual history-based testing misses anorectal Chlamydia trachomatis and neisseria gonorrhoeae infections in swingers and men who have sex with men. <i>Sexually Transmitted Diseases</i> , 2013 , 40, 285-9	2.4	54
129	Evaluation of the anatomical site distribution of chlamydia and gonorrhoea in men who have sex with men and in high-risk women by routine testing: cross-sectional study revealing missed opportunities for treatment strategies. <i>Sexually Transmitted Infections</i> , 2014 , 90, 58-60	2.8	49
128	The impact of experiencing lipodystrophy on the sexual behaviour and well-being among HIV-infected homosexual men. <i>Aids</i> , 2001 , 15, 812-3	3.5	48
127	The value of testing multiple anatomic sites for gonorrhoea and chlamydia in sexually transmitted infection centres in the Netherlands, 2006-2010. <i>International Journal of STD and AIDS</i> , 2012 , 23, 626-31	1.4	47
126	Q fever: single-point source outbreak with high attack rates and massive numbers of undetected infections across an entire region. <i>Clinical Infectious Diseases</i> , 2012 , 55, 1591-9	11.6	46

125	High prevalence of Epstein-Barr virus type 2 among homosexual men is caused by sexual transmission. <i>Journal of Infectious Diseases</i> , 2000 , 181, 2045-9	7	45
124	High non-responsiveness of males and the elderly to standard hepatitis B vaccination among a large cohort of healthy employees. <i>Journal of Clinical Virology</i> , 2013 , 58, 262-4	14.5	44
123	Waterborne gastroenteritis outbreak at a scouting camp caused by two norovirus genogroups: GI and GII. <i>Journal of Clinical Virology</i> , 2010 , 47, 268-72	14.5	42
122	High Proportion of Anorectal Chlamydia trachomatis and Neisseria gonorrhoeae After Routine Universal Urogenital and Anorectal Screening in Women Visiting the Sexually Transmitted Infection Clinic. <i>Clinical Infectious Diseases</i> , 2017 , 64, 1705-1710	11.6	39
121	Chlamydia trachomatis test-of-cure cannot be based on a single highly sensitive laboratory test taken at least 3 weeks after treatment. <i>PLoS ONE</i> , 2012 , 7, e34108	3.7	39
120	High and persistent excretion of hepatitis A virus in immunocompetent patients. <i>Journal of Medical Virology</i> , 2006 , 78, 1398-405	19.7	39
119	HIV incidence and HIV testing behavior in men who have sex with men: using three incidence sources, The Netherlands, 1984-2005. <i>Aids</i> , 2007 , 21, 491-9	3.5	38
118	Relation between infection and pelvic inflammatory disease, ectopic pregnancy and tubal factor infertility in a Dutch cohort of women previously tested for chlamydia in a chlamydia screening trial. <i>Sexually Transmitted Infections</i> , 2019 , 95, 300-306	2.8	38
117	Detection of anorectal and cervicovaginal Chlamydia trachomatis infections following azithromycin treatment: prospective cohort study with multiple time-sequential measures of rRNA, DNA, quantitative load and symptoms. <i>PLoS ONE</i> , 2013 , 8, e81236	3.7	35
116	The Spatial Distribution of Hepatitis C Virus Infections and Associated Determinants--An Application of a Geographically Weighted Poisson Regression for Evidence-Based Screening Interventions in Hotspots. <i>PLoS ONE</i> , 2015 , 10, e0135656	3.7	34
115	Treatment Effectiveness of Azithromycin and Doxycycline in Uncomplicated Rectal and Vaginal Chlamydia trachomatis Infections in Women: A Multicenter Observational Study (FemCure). <i>Clinical Infectious Diseases</i> , 2019 , 69, 1946-1954	11.6	33
114	Socially isolated individuals are more prone to have newly diagnosed and prevalent type 2 diabetes mellitus - the Maastricht study. <i>BMC Public Health</i> , 2017 , 17, 955	4.1	31
113	What explains anorectal chlamydia infection in women? Implications of a mathematical model for test and treatment strategies. <i>Sexually Transmitted Infections</i> , 2017 , 93, 270-275	2.8	30
112	Molecular epidemiology of Neisseria gonorrhoeae in Amsterdam, The Netherlands, shows distinct heterosexual and homosexual networks. <i>Journal of Clinical Microbiology</i> , 2006 , 44, 2689-97	9.7	30
111	Prevalence of and Factors Associated with Rectal-Only Chlamydia and Gonorrhoea in Women and in Men Who Have Sex with Men. <i>PLoS ONE</i> , 2015 , 10, e0140297	3.7	30
110	Seroprevalence and determinants of eight high-risk human papillomavirus types in homosexual men, heterosexual men, and women: a population-based study in Amsterdam. <i>Sexually Transmitted Diseases</i> , 2010 , 37, 672-80	2.4	28
109	Correlates of resistance to HIV-1 infection in homosexual men with high-risk sexual behaviour. <i>Aids</i> , 2004 , 18, 1117-26	3.5	28
108	Chemsex among men who have sex with men living outside major cities and associations with sexually transmitted infections: A cross-sectional study in the Netherlands. <i>PLoS ONE</i> , 2019 , 14, e02167327	3.7	27

107	Low CD4 T cell counts before HIV-1 seroconversion do not affect disease progression in Ethiopian factory workers. <i>Journal of Infectious Diseases</i> , 2005 , 192, 739-48	7	27
106	Body mass index course in asymptomatic HIV-infected homosexual men and the predictive value of a decrease of body mass index for progression to AIDS. <i>Journal of Acquired Immune Deficiency Syndromes</i> , 1998 , 19, 254-9		27
105	Older and swinging; need to identify hidden and emerging risk groups at STI clinics. <i>Sexually Transmitted Infections</i> , 2010 , 86, 315-7	2.8	26
104	Chlamydia trachomatis and the Risk of Pelvic Inflammatory Disease, Ectopic Pregnancy, and Female Infertility: A Retrospective Cohort Study Among Primary Care Patients. <i>Clinical Infectious Diseases</i> , 2019 , 69, 1517-1525	11.6	25
103	Trends in antibiotic prescribing in adults in Dutch general practice. <i>PLoS ONE</i> , 2012 , 7, e51860	3.7	25
102	Simple markers for initiating antiretroviral therapy among HIV-infected Ethiopians. <i>Aids</i> , 2003 , 17, 815-93.5		25
101	Viability-PCR Shows That NAAT Detects a High Proportion of DNA from Non-Viable Chlamydia trachomatis. <i>PLoS ONE</i> , 2016 , 11, e0165920	3.7	25
100	Revaccination with Fendrix [®] or HBVaxPro [®] results in better response rates than does revaccination with three doses of Engerix-B [®] in previous non-responders. <i>Vaccine</i> , 2012 , 30, 6734-7	4.1	23
99	High burden of STI and HIV in male sex workers working as internet escorts for men in an observational study: a hidden key population compared with female sex workers and other men who have sex with men. <i>BMC Infectious Diseases</i> , 2015 , 15, 291	4	22
98	Effectiveness of an opting-out strategy for HIV testing: evaluation of 4 years of standard HIV testing in a STI clinic. <i>Sexually Transmitted Infections</i> , 2009 , 85, 226-30	2.8	20
97	Chlamydia trachomatis load in population-based screening and STI-clinics: implications for screening policy. <i>PLoS ONE</i> , 2015 , 10, e0121433	3.7	19
96	Social Network Characteristics Are Associated With Type 2 Diabetes Complications: The Maastricht Study. <i>Diabetes Care</i> , 2018 , 41, 1654-1662	14.6	18
95	Re-screening Chlamydia trachomatis positive subjects: a comparison of practices between an STI clinic, general practitioners and gynaecologists. <i>Sexually Transmitted Infections</i> , 2013 , 89, 25-7	2.8	18
94	Who tests whom? A comprehensive overview of Chlamydia trachomatis test practices in a Dutch region among different STI care providers for urogenital, anorectal and oropharyngeal sites in young people: a cross-sectional study. <i>Sexually Transmitted Infections</i> , 2016 , 92, 211-7	2.8	18
93	Screening for HIV, hepatitis B and syphilis on dried blood spots: A promising method to better reach hidden high-risk populations with self-collected sampling. <i>PLoS ONE</i> , 2017 , 12, e0186722	3.7	17
92	Drug use, sexual risk behaviour and sexually transmitted infections among swingers: a cross-sectional study in The Netherlands. <i>Sexually Transmitted Infections</i> , 2015 , 91, 31-6	2.8	16
91	Oral and topical antibiotic prescriptions for children in general practice. <i>Archives of Disease in Childhood</i> , 2013 , 98, 228-31	2.2	16
90	Influence of temperature, medium, and storage duration on Chlamydia trachomatis DNA detection by PCR. <i>Journal of Clinical Microbiology</i> , 2013 , 51, 990-2	9.7	16

89	Anorectal Chlamydia trachomatis Load Is Similar in Men Who Have Sex with Men and Women Reporting Anal Sex. <i>PLoS ONE</i> , 2015 , 10, e0134991	3.7	15
88	A comprehensive overview of urogenital, anorectal and oropharyngeal Neisseria gonorrhoeae testing and diagnoses among different STI care providers: a cross-sectional study. <i>BMC Infectious Diseases</i> , 2017 , 17, 290	4	13
87	Design of the FemCure study: prospective multicentre study on the transmission of genital and extra-genital Chlamydia trachomatis infections in women receiving routine care. <i>BMC Infectious Diseases</i> , 2016 , 16, 381	4	13
86	Strong decline in herpes simplex virus antibodies over time among young homosexual men is associated with changing sexual behavior. <i>American Journal of Epidemiology</i> , 2000 , 152, 666-73	3.8	13
85	Using intervention mapping for the development of a targeted secure web-based outreach strategy named SafeFriend, for Chlamydia trachomatis testing in young people at risk. <i>BMC Public Health</i> , 2013 , 13, 996	4.1	12
84	Rise in seroprevalence of herpes simplex virus type 1 among highly sexual active homosexual men and an increasing association between herpes simplex virus type 2 and HIV over time (1984-2003). <i>European Journal of Epidemiology</i> , 2007 , 22, 937-44	12.1	12
83	Review of Chlamydia trachomatis viability methods: assessing the clinical diagnostic impact of NAAT positive results. <i>Expert Review of Molecular Diagnostics</i> , 2018 , 18, 739-747	3.8	11
82	Coxiella burnetii Infection Is Lower in Children than in Adults After Community Exposure: Overlooked Cause of Infrequent Q Fever Reporting in the Young. <i>Pediatric Infectious Disease Journal</i> , 2015 , 34, 1283-8	3.4	11
81	Low positivity rate after systematic screening for Trichomonas vaginalis in three patient cohorts from general practitioners, STI clinic and a national population-based chlamydia screening study. <i>Sexually Transmitted Infections</i> , 2013 , 89, 532-4	2.8	11
80	Characterization of an HIV-1 group M variant that is distinct from the known subtypes. <i>AIDS Research and Human Retroviruses</i> , 2007 , 23, 466-70	1.6	11
79	Low HIV-testing rates among younger high-risk homosexual men in Amsterdam. <i>Sexually Transmitted Infections</i> , 2007 , 83, 387-91	2.8	11
78	Spontaneous clearance of urogenital, anorectal and oropharyngeal and in women, MSM and heterosexual men visiting the STI clinic: a prospective cohort study. <i>Sexually Transmitted Infections</i> , 2019 , 95, 505-510	2.8	10
77	Chlamydia trachomatis testing among young people: what is the role of stigma?. <i>BMC Public Health</i> , 2015 , 15, 651	4.1	10
76	Detecting Hepatitis B and C by Combined Public Health and Primary Care Birth Cohort Testing. <i>Annals of Family Medicine</i> , 2018 , 16, 21-27	2.9	10
75	Barriers to and facilitators of partner notification for chlamydia trachomatis among health care professionals. <i>BMC Health Services Research</i> , 2014 , 14, 647	2.9	10
74	Nasal carriage of Streptococcus pneumoniae serotypes and Staphylococcus aureus in Streptococcus pneumoniae-vaccinated and non-vaccinated young children. <i>Epidemiology and Infection</i> , 2013 , 141, 631-8	4.3	10
73	Two decades of hepatitis B infections among drug users in Amsterdam: are they still a high-risk group?. <i>Journal of Medical Virology</i> , 2009 , 81, 1163-9	19.7	10
72	Acceptance of Home-Based Chlamydia Genital and Anorectal Testing Using Short Message Service (SMS) in Previously Tested Young People and Their Social and Sexual Networks. <i>PLoS ONE</i> , 2015 , 10, e0133575	3.7	10

71	Incidence of repeat testing and diagnoses of and in swingers, homosexual and heterosexual men and women at two large Dutch STI clinics, 2006-2013. <i>Sexually Transmitted Infections</i> , 2017 , 93, 383-389	2.8	9
70	Antibiotic use before chlamydia and gonorrhea genital and extragenital screening in the sexually transmitted infection clinical setting. <i>Antimicrobial Agents and Chemotherapy</i> , 2015 , 59, 121-8	5.9	9
69	Chlamydia trachomatis bacterial load, estimated by Cq values, in urogenital samples from men and women visiting the general practice, hospital or STI clinic. <i>PLoS ONE</i> , 2019 , 14, e0215606	3.7	8
68	Is the current pertussis incidence only the results of testing? A spatial and space-time analysis of pertussis surveillance data using cluster detection methods and geographically weighted regression modelling. <i>PLoS ONE</i> , 2017 , 12, e0172383	3.7	8
67	Hepatitis E prevalence in a sexual high-risk population compared to the general population. <i>PLoS ONE</i> , 2018 , 13, e0191798	3.7	8
66	Predictors of Chlamydia Trachomatis testing: perceived norms, susceptibility, changes in partner status, and underestimation of own risk. <i>BMC Public Health</i> , 2016 , 16, 55	4.1	8
65	Assessment of rectal viable load in women by viability-PCR. <i>Sexually Transmitted Infections</i> , 2020 , 96, 85-88	2.8	8
64	Test of cure, retesting and extragenital testing practices for Chlamydia trachomatis and Neisseria gonorrhoeae among general practitioners in different socioeconomic status areas: A retrospective cohort study, 2011-2016. <i>PLoS ONE</i> , 2018 , 13, e0194351	3.7	8
63	Where to go to in chlamydia control? From infection control towards infectious disease control. <i>Sexually Transmitted Infections</i> , 2021 , 97, 501-506	2.8	8
62	Genital and anal bacterial load in concurrently infected women: a cross-sectional study. <i>Sexually Transmitted Infections</i> , 2019 , 95, 317-321	2.8	7
61	Spontaneous clearance of accounting for bacterial viability in vaginally or rectally infected women (FemCure). <i>Sexually Transmitted Infections</i> , 2020 , 96, 541-548	2.8	7
60	Evaluation of one-sample testing of self-obtained vaginal swabs and first catch urine samples separately and in combination for the detection of Chlamydia trachomatis by two amplified DNA assays in women visiting a sexually transmitted disease clinic. <i>Sexually Transmitted Diseases</i> , 2011 , 36, 500-505	2.4	7
59	What Is the Optimal Testing Strategy for Oropharyngeal Neisseria gonorrhoeae in Men Who Have Sex With Men? Comparing Selective Testing Versus Routine Universal Testing From Dutch Sexually Transmitted Infection Clinic Data (2008-2017). <i>Clinical Infectious Diseases</i> , 2020 , 71, 944-951	11.6	7
58	Men and Women Have an Equal Oropharyngeal and Anorectal Chlamydia trachomatis Bacterial Load: A Comparison of 3 Anatomic Sites. <i>Journal of Infectious Diseases</i> , 2021 , 223, 1582-1589	7	7
57	Q fever: Evidence of a massive yet undetected cross-border outbreak, with ongoing risk of extra mortality, in a Dutch-German border region. <i>Transboundary and Emerging Diseases</i> , 2020 , 67, 1660-1670	4.2	6
56	Social networks in relation to self-reported symptomatic infections in individuals aged 40-75 - the Maastricht study. <i>BMC Infectious Diseases</i> , 2018 , 18, 300	4	6
55	Improving sexual health for HIV patients by providing a combination of integrated public health and hospital care services; a one-group pre- and post test intervention comparison. <i>BMC Public Health</i> , 2012 , 12, 1118	4.1	6
54	Identification of hidden key hepatitis C populations: an evaluation of screening practices using mixed epidemiological methods. <i>PLoS ONE</i> , 2012 , 7, e51194	3.7	6

53	Retrieval and re-evaluation of previously diagnosed chronic hepatitis C infections lost to medical follow-up in the Netherlands. <i>European Journal of Gastroenterology and Hepatology</i> , 2020 , 32, 851-856	2.2	6
52	Standardisation is necessary in urogenital and extragenital bacterial load determination by quantitative PCR: a review of literature and retrospective study. <i>Sexually Transmitted Infections</i> , 2019 , 95, 562-568	2.8	5
51	Do swingers self-identify as swingers when attending STI services for testing? A cross-sectional study. <i>Sexually Transmitted Infections</i> , 2018 , 94, 559-561	2.8	5
50	A Web-Based Respondent Driven Sampling Pilot Targeting Young People at Risk for Chlamydia Trachomatis in Social and Sexual Networks with Testing: A Use Evaluation. <i>International Journal of Environmental Research and Public Health</i> , 2015 , 12, 9889-906	4.6	5
49	The added value of chlamydia screening between 2008-2010 in reaching young people in addition to chlamydia testing in regular care; an observational study. <i>BMC Infectious Diseases</i> , 2014 , 14, 612	4	5
48	Prevalence of Non-Volitional Sex Types and Associated Factors: A National Sample of Young People. <i>PLoS ONE</i> , 2015 , 10, e0132847	3.7	5
47	Men and Women Have Similar Neisseria gonorrhoeae Bacterial Loads: a Comparison of Three Anatomical Sites. <i>Journal of Clinical Microbiology</i> , 2020 , 58,	9.7	5
46	Routine universal testing versus selective or incidental testing for oropharyngeal Neisseria gonorrhoeae in women in the Netherlands: a retrospective cohort study. <i>Lancet Infectious Diseases</i> , 2021 , 21, 858-867	25.5	5
45	Association between social network characteristics and prevalent and incident depression: The Maastricht Study. <i>Journal of Affective Disorders</i> , 2021 , 293, 338-346	6.6	5
44	Viable Bacterial Load Is Key to Azithromycin Treatment Failure in Rectally Chlamydia trachomatis Infected Women (FemCure). <i>Journal of Infectious Diseases</i> , 2019 , 220, 1389-1390	7	4
43	Participation, retention, and associated factors of women in a prospective multicenter study on Chlamydia trachomatis infections (FemCure). <i>PLoS ONE</i> , 2020 , 15, e0230413	3.7	4
42	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study. <i>PLoS ONE</i> , 2020 , 15, e0235467	3.7	4
41	Mucosal shedding of human herpesvirus 8. <i>New England Journal of Medicine</i> , 2001 , 344, 690-1; author reply 691-2	59.2	4
40	Social networks and type 2 diabetes: a narrative review. <i>Diabetologia</i> , 2021 , 64, 1905-1916	10.3	4
39	Comparison of urine samples and penile swabs for detection of human papillomavirus in HIV-negative Dutch men. <i>Sexually Transmitted Infections</i> , 2016 , 92, 467-9	2.8	4
38	Oropharyngeal in women; spontaneous clearance and cure after treatment (FemCure). <i>Sexually Transmitted Infections</i> , 2021 , 97, 147-151	2.8	4
37	The Characteristics of Patients Frequently Tested and Repeatedly Infected with. <i>International Journal of Environmental Research and Public Health</i> , 2020 , 17,	4.6	3
36	Antibody Testing in Estimating Past Exposure to C in the Netherlands Chlamydia Cohort Study. <i>Microorganisms</i> , 2019 , 7,	4.9	3

35	Improving STD testing behavior among high-risk young adults by offering STD testing at a vocational school. <i>BMC Public Health</i> , 2011 , 11, 750	4.1	3
34	Pregnancies and Time to Pregnancy in Women With and Without a Previous Chlamydia trachomatis Infection. <i>Sexually Transmitted Diseases</i> , 2020 , 47, 739-747	2.4	3
33	The characteristics of patients frequently tested and repeatedly infected with Chlamydia trachomatis in Southwest Limburg, the Netherlands. <i>BMC Public Health</i> , 2020 , 20, 1239	4.1	3
32	Barriers and facilitators to infection prevention and control in Dutch residential care facilities for people with intellectual and developmental disabilities: A theory-informed qualitative study. <i>PLoS ONE</i> , 2021 , 16, e0258701	3.7	2
31	Sexually Transmitted Infection and Chemsex Also Highly Prevalent Among Male Sex Workers Outside Amsterdam, The Netherlands. <i>Sexually Transmitted Diseases</i> , 2020 , 47, e15	2.4	2
30	Social networks and infectious diseases prevention behavior: A cross-sectional study in people aged 40 years and older. <i>PLoS ONE</i> , 2021 , 16, e0251862	3.7	2
29	Determinants associated with viable genital or rectal bacterial load (FemCure). <i>Sexually Transmitted Infections</i> , 2021 ,	2.8	2
28	Pharyngeal Chlamydia trachomatis in men who have sex with men (MSM) in the Netherlands: a large retrospective cohort study. <i>Clinical Infectious Diseases</i> , 2021 ,	11.6	2
27	Pilot implementation of a home-care programme with chlamydia, gonorrhoea, hepatitis B, and syphilis self-sampling in HIV-positive men who have sex with men. <i>BMC Infectious Diseases</i> , 2020 , 20, 925	4	1
26	Incident urogenital and anorectal in women: the role of sexual exposure and autoinoculation: a multicentre observational study (FemCure).. <i>Sexually Transmitted Infections</i> , 2022 ,	2.8	1
25	Online respondent-driven detection for enhanced contact tracing of close-contact infectious diseases: benefits and barriers for public health practice		1
24	Men and Women Repeatedly Infected With Chlamydia trachomatis Have a Lower Urogenital Bacterial Load. <i>Sexually Transmitted Diseases</i> , 2020 , 47, e51-e53	2.4	1
23	Sex Abroad Among Men Who Have Sex With Men and Its Association With Chemsex, Sexual Risk Behavior, and Sexually Transmitted Diseases: A Cross-Sectional Study in the Netherlands. <i>Sexually Transmitted Diseases</i> , 2020 , 47, e29-e32	2.4	1
22	Use of doxycycline and other antibiotics to prevent STIs among men who have sex with men visiting sexual health clinics in the Netherlands. <i>Sexually Transmitted Infections</i> , 2020 , 96, 550-551	2.8	1
21	The Availability and Feasibility of Chemsex Services Within Sexually Transmitted Disease Clinics in the Netherlands: A Cross-Sectional Survey Among Sexually Transmitted Disease Nurses. <i>Sexually Transmitted Diseases</i> , 2020 , 47, 639-644	2.4	1
20	Prevalence of drug use during sex among swingers and perceived benefits and risks - a cross-sectional internet survey in the Netherlands. <i>Sexually Transmitted Infections</i> , 2020 , 96, 40-46	2.8	1
19	Can Previous Associations of Single Nucleotide Polymorphisms in the , , , and Genes in the Susceptibility to and Severity of Infections Be Confirmed?. <i>Pathogens</i> , 2021 , 10,	4.5	1
18	Routine universal testing versus selective or incidental testing for oropharyngeal Chlamydia trachomatis in women in the Netherlands: a retrospective cohort study.. <i>Lancet Infectious Diseases</i> , The , 2021 ,	25.5	1

17	HIV testing behaviour and associated factors in men who have sex with men by level of urbanisation: a cross-sectional study in the Netherlands.. <i>BMJ Open</i> , 2022 , 12, e049175	3	○
16	Online respondent-driven detection for enhanced contact tracing of close-contact infectious diseases: benefits and barriers for public health practice. <i>BMC Infectious Diseases</i> , 2021 , 21, 358	4	○
15	How aware are swingers about their swing sex partnersRisk behaviours, and sexually transmitted infection status?. <i>BMC Infectious Diseases</i> , 2021 , 21, 172	4	○
14	Controversies and evidence on Chlamydia testing and treatment in asymptomatic women and men who have sex with men: a narrative review.. <i>BMC Infectious Diseases</i> , 2022 , 22, 255	4	○
13	Barriers and facilitators to infection prevention and control in Dutch psychiatric institutions: a theory-informed qualitative study.. <i>BMC Infectious Diseases</i> , 2022 , 22, 243	4	○
12	Exposure factors associated with SARS-CoV-2 seroprevalence during the first eight months of the COVID-19 pandemic in the Netherlands: A cross-sectional study.. <i>PLoS ONE</i> , 2022 , 17, e0268057	3.7	○
11	Pertussis diagnostic practices of general practitioners in the Netherlands: A survey study. <i>European Journal of General Practice</i> , 2019 , 25, 214-219	2.8	
10	Limits of chlamydial diagnostic tests in epidemiological studies. <i>Journal of Infectious Diseases</i> , 2014 , 209, 2017-8	7	
9	Signs and symptoms do not predict, but may help rule out acute Q fever in favour of other respiratory tract infections, and reduce antibiotics overuse in primary care. <i>BMC Infectious Diseases</i> , 2020 , 20, 690	4	
8	Feasibility of anorectal chlamydia testing in women: a cross-sectional survey among general practitioners. <i>Family Practice</i> , 2021 , 38, 724-730	1.9	
7	Culture-Independent Genotyping Revealed 3 Strain Clusters in a Potential Neisseria gonorrhoeae Outbreak in Young Heterosexuals (. <i>Sexually Transmitted Diseases</i> , 2021 , 48, 536-541	2.4	
6	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		
5	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		
4	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		
3	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		
2	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		
1	Attitude and beliefs about the social environment associated with chemsex among MSM visiting STI clinics in the Netherlands: An observational study 2020 , 15, e0235467		