Vaclav Horejsi

List of Publications by Citations

Source: https://exaly.com/author-pdf/9538810/vaclav-horejsi-publications-by-citations.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

138 8,212 49 87 g-index

151 8,782 7.6 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
138	GPI-anchored cell-surface molecules complexed to protein tyrosine kinases. <i>Science</i> , 1991 , 254, 1016-9	33.3	780
137	Phosphoprotein associated with glycosphingolipid-enriched microdomains (PAG), a novel ubiquitously expressed transmembrane adaptor protein, binds the protein tyrosine kinase csk and is involved in regulation of T cell activation. <i>Journal of Experimental Medicine</i> , 2000 , 191, 1591-604	16.6	401
136	Urokinase plasminogen activator receptor, beta 2-integrins, and Src-kinases within a single receptor complex of human monocytes. <i>Journal of Experimental Medicine</i> , 1995 , 181, 1381-90	16.6	341
135	Biochemical characterization of a soluble form of the 53-kDa monocyte surface antigen. <i>European Journal of Immunology</i> , 1986 , 16, 1583-9	6.1	214
134	Tetraspan microdomains distinct from lipid rafts enrich select peptide-MHC class II complexes. <i>Nature Immunology</i> , 2002 , 3, 61-8	19.1	192
133	Disulfide bond-mediated dimerization of HLA-G on the cell surface. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 16180-5	11.5	185
132	Non-T cell activation linker (NTAL): a transmembrane adaptor protein involved in immunoreceptor signaling. <i>Journal of Experimental Medicine</i> , 2002 , 196, 1617-26	16.6	176
131	Transmembrane adaptor proteins: organizers of immunoreceptor signalling. <i>Nature Reviews Immunology</i> , 2004 , 4, 603-16	36.5	166
130	Phosphorylation-dependent regulation of T-cell activation by PAG/Cbp, a lipid raft-associated transmembrane adaptor. <i>Molecular and Cellular Biology</i> , 2003 , 23, 2017-28	4.8	166
129	M6P/IGFII-receptor complexes urokinase receptor and plasminogen for activation of transforming growth factor-beta1. <i>European Journal of Immunology</i> , 1999 , 29, 1004-13	6.1	144
128	Characterization of monoclonal antibodies recognizing HLA-G or HLA-E: new tools to analyze the expression of nonclassical HLA class I molecules. <i>Human Immunology</i> , 2003 , 64, 315-26	2.3	134
127	LAT displacement from lipid rafts as a molecular mechanism for the inhibition of T cell signaling by polyunsaturated fatty acids. <i>Journal of Biological Chemistry</i> , 2002 , 277, 28418-23	5.4	134
126	Structural relationship between the soluble and membrane-bound forms of human monocyte surface glycoprotein CD14. <i>Molecular Immunology</i> , 1989 , 26, 657-62	4.3	133
125	Release from tonic inhibition of T cell activation through transient displacement of C-terminal Src kinase (Csk) from lipid rafts. <i>Journal of Biological Chemistry</i> , 2001 , 276, 29313-8	5.4	128
124	Complexes of HLA-G protein on the cell surface are important for leukocyte Ig-like receptor-1 function. <i>Journal of Immunology</i> , 2003 , 171, 1343-51	5.3	124
123	Characterization of a broadly expressed human leucocyte surface antigen MEM-43 anchored in membrane through phosphatidylinositol. <i>Molecular Immunology</i> , 1989 , 26, 153-61	4.3	124
122	Expression and release of soluble HLA-E is an immunoregulatory feature of endothelial cell activation. <i>Blood</i> , 2007 , 109, 2806-14	2.2	122

121	Exogenous glycosyl phosphatidylinositol-anchored CD59 associates with kinases in membrane clusters on U937 cells and becomes Ca(2+)-signaling competent. <i>Journal of Cell Biology</i> , 1995 , 131, 669-	7 ⁷ 7 ³	120
120	The roles of membrane microdomains (rafts) in T cell activation. <i>Immunological Reviews</i> , 2003 , 191, 148-	64 .3	116
119	T cell activation-associated epitopes of CD147 in regulation of the T cell response, and their definition by antibody affinity and antigen density. <i>International Immunology</i> , 1999 , 11, 777-86	4.9	114
118	Novel structurally distinct family of leucocyte surface glycoproteins including CD9, CD37, CD53 and CD63. <i>FEBS Letters</i> , 1991 , 288, 1-4	3.8	113
117	Negative regulation of mast cell signaling and function by the adaptor LAB/NTAL. <i>Journal of Experimental Medicine</i> , 2004 , 200, 1001-13	16.6	111
116	The functional interactions between CD98, beta1-integrins, and CD147 in the induction of U937 homotypic aggregation. <i>Blood</i> , 2001 , 98, 374-82	2.2	111
115	Colocalization of the tetraspanins, CO-029 and CD151, with integrins in human pancreatic adenocarcinoma: impact on cell motility. <i>Clinical Cancer Research</i> , 2005 , 11, 2840-52	12.9	106
114	NTAL phosphorylation is a pivotal link between the signaling cascades leading to human mast cell degranulation following Kit activation and Fc epsilon RI aggregation. <i>Blood</i> , 2004 , 104, 207-14	2.2	102
113	Noncovalent associations of T lymphocyte surface proteins. <i>European Journal of Immunology</i> , 1996 , 26, 2335-43	6.1	97
112	LIME: a new membrane Raft-associated adaptor protein involved in CD4 and CD8 coreceptor signaling. <i>Journal of Experimental Medicine</i> , 2003 , 198, 1453-62	16.6	94
111	Differential role of glycolipid-enriched membrane domains in glycoprotein VI- and integrin-mediated phospholipase Cgamma2 regulation in platelets. <i>Biochemical Journal</i> , 2002 , 364, 755-	.≩\$ ⁸	93
110	CD molecules 2005: human cell differentiation molecules. <i>Blood</i> , 2005 , 106, 3123-6	2.2	92
109	The CD85J/leukocyte inhibitory receptor-1 distinguishes between conformed and beta 2-microglobulin-free HLA-G molecules. <i>Journal of Immunology</i> , 2005 , 175, 4866-74	5.3	92
108	Cross-linking of CD59 and of other glycosyl phosphatidylinositol-anchored molecules on neutrophils triggers cell activation via tyrosine kinase. <i>European Journal of Immunology</i> , 1993 , 23, 2841-	50 ¹	90
107	Large, detergent-resistant complexes containing murine antigens Thy-1 and Ly-6 and protein tyrosine kinase p56lck. <i>European Journal of Immunology</i> , 1993 , 23, 825-31	6.1	89
106	CD59 molecule: a second ligand for CD2 in T cell adhesion. <i>European Journal of Immunology</i> , 1992 , 22, 2943-7	6.1	83
105	CD Nomenclature 2015: Human Leukocyte Differentiation Antigen Workshops as a Driving Force in Immunology. <i>Journal of Immunology</i> , 2015 , 195, 4555-63	5.3	80
104	LST1/A is a myeloid leukocyte-specific transmembrane adaptor protein recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to the plasma membrane <i>Journal of Biological Chemistry</i> , 2013 ,	5.4	78

103	Activation of human monocytes and granulocytes by monoclonal antibodies to glycosylphosphatidylinositol-anchored antigens. <i>European Journal of Immunology</i> , 1993 , 23, 2782-91	6.1	78
102	Grb2 and the non-T cell activation linker NTAL constitute a Ca(2+)-regulating signal circuit in B lymphocytes. <i>Immunity</i> , 2004 , 21, 681-91	32.3	73
101	GPI-microdomains (membrane rafts) and signaling of the multi-chain interleukin-2 receptor in human lymphoma/leukemia T cell lines. <i>FEBS Journal</i> , 2002 , 269, 1199-208		72
100	The Src family kinase Hck regulates mast cell activation by suppressing an inhibitory Src family kinase Lyn. <i>Blood</i> , 2007 , 110, 2511-9	2.2	64
99	Release and intercellular transfer of cell surface CD81 via microparticles. <i>Journal of Immunology</i> , 2002 , 169, 5531-7	5.3	63
98	Some theoretical aspects of affinity electrophoresis. <i>Journal of Chromatography A</i> , 1979 , 178, 1-13	4.5	61
97	Review: Affinity electrophoresis. <i>Analytical Biochemistry</i> , 1981 , 112, 1-8	3.1	57
96	Combined spatial and enzymatic regulation of Csk by cAMP and protein kinase a inhibits T cell receptor signaling. <i>Journal of Biological Chemistry</i> , 2003 , 278, 17597-600	5.4	56
95	Kit- and Fc epsilonRI-induced differential phosphorylation of the transmembrane adaptor molecule NTAL/LAB/LAT2 allows flexibility in its scaffolding function in mast cells. <i>Cellular Signalling</i> , 2008 , 20, 195-205	4.9	55
94	CD4 segregates into specific detergent-resistant T-cell membrane microdomains. <i>Tissue Antigens</i> , 1999 , 53, 33-40		55
93	The human leucocyte surface antigen CD53 is a protein structurally similar to the CD37 and MRC OX-44 antigens. <i>Immunogenetics</i> , 1990 , 32, 281-5	3.2	54
92	Human monocyte activation induced by an anti-CD14 monoclonal antibody. <i>Immunology Letters</i> , 1988 , 19, 321-7	4.1	54
91	Adapters in lymphocyte signaling. <i>Journal of Clinical Investigation</i> , 2002 , 109, 301-309	15.9	54
90	A novel negative regulatory function of the phosphoprotein associated with glycosphingolipid-enriched microdomains: blocking Ras activation. <i>Blood</i> , 2007 , 110, 596-615	2.2	49
89	Constitutive exclusion of Csk from Hck-positive membrane microdomains permits Src kinase-dependent proliferation of Theileria-transformed B lymphocytes. <i>Blood</i> , 2003 , 101, 1874-81	2.2	49
88	TGF-beta-induced apoptosis in endothelial cells mediated by M6P/IGFII-R and mini-plasminogen. <i>Journal of Cell Science</i> , 2005 , 118, 4577-86	5.3	48
87	The N terminus of mannose 6-phosphate/insulin-like growth factor 2 receptor in regulation of fibrinolysis and cell migration. <i>Journal of Biological Chemistry</i> , 2002 , 277, 40575-82	5.4	48
86	Non-T cell activation linker (NTAL) negatively regulates TREM-1/DAP12-induced inflammatory cytokine production in myeloid cells. <i>Journal of Immunology</i> , 2007 , 178, 1991-9	5.3	47

85	Lipid rafts and their roles in T-cell activation. <i>Microbes and Infection</i> , 2005 , 7, 310-6	9.3	47
84	Novel molecular mechanisms of dendritic cell-induced T cell activation. <i>International Immunology</i> , 2000 , 12, 1051-61	ļ.9	46
83	Monoclonal antibodies specific for the empty conformation of HLA-DR1 reveal aspects of the conformational change associated with peptide binding. <i>Journal of Biological Chemistry</i> , 2004 , 279, 1656	i: / 70	44
82	SCIMP, a transmembrane adaptor protein involved in major histocompatibility complex class II signaling. <i>Molecular and Cellular Biology</i> , 2011 , 31, 4550-62	μ.8	41
81	Qualitative and quantitative applications of affinity electrophoresis for the study of protein I gand interactions: A review. <i>Biomedical Applications</i> , 1986 , 376, 49-67		41
80	Single and combined deletions of the NTAL/LAB and LAT adaptors minimally affect B-cell development and function. <i>Molecular and Cellular Biology</i> , 2005 , 25, 4455-65	₄ .8	40
79	Membrane microdomains in immunoreceptor signaling. FEBS Letters, 2014 , 588, 2392-7	3.8	39
78	CD43-mediated signals induce DNA binding activity of AP-1, NF-AT, and NFkappa B transcription factors in human T lymphocytes. <i>Journal of Biological Chemistry</i> , 2000 , 275, 31460-8	5.4	39
77	MicroRNA editing facilitates immune elimination of HCMV infected cells. <i>PLoS Pathogens</i> , 2014 , 10, e100	3 0 63	35
76	Btk is a positive regulator in the TREM-1/DAP12 signaling pathway. <i>Blood</i> , 2011 , 118, 936-45	2.2	35
75	Transmembrane adaptor molecules: a new category of lymphoid-cell markers. <i>Blood</i> , 2006 , 107, 213-21 2	2.2	35
74	Transmembrane adaptor proteins in membrane microdomains: important regulators of immunoreceptor signaling. <i>Immunology Letters</i> , 2004 , 92, 43-9	ļ.1	34
73	Nitrocellulose membrane as an antigen or antibody carrier for screening hybridoma cultures. <i>Journal of Immunological Methods</i> , 1983 , 62, 325-9	2.5	34
72	Antibody array analysis with label-based detection and resolution of protein size. <i>Molecular and Cellular Proteomics</i> , 2009 , 8, 245-57	7.6	32
71	Regulation of CD43-induced U937 homotypic aggregation. <i>Experimental Cell Research</i> , 2003 , 290, 155-674	ļ. 2	32
70	Conformational variation of surface class II MHC proteins during myeloid dendritic cell differentiation accompanies structural changes in lysosomal MIIC. <i>Journal of Immunology</i> , 2005 , 175, 4935-47	5.3	32
69	A new type of membrane raft-like microdomains and their possible involvement in TCR signaling. Journal of Immunology, 2010 , 184, 3689-96	5.3	31
68	Structural study of the O-linked sugar chains of human leukocyte tyrosine phosphatase CD45. <i>FEBS Journal</i> , 1998 , 251, 288-94		31

67	HLA-G levels in serum and plasma. <i>Tissue Antigens</i> , 2006 , 67, 111-6		31
66	Palmitoylated transmembrane adaptor proteins in leukocyte signaling. <i>Cellular Signalling</i> , 2014 , 26, 895	5-29.092	30
65	The Csk-binding protein PAG regulates PDGF-induced Src mitogenic signaling via GM1. <i>Journal of Cell Biology</i> , 2008 , 182, 603-14	7.3	29
64	HLA-E: strong association with beta2-microglobulin and surface expression in the absence of HLA class I signal sequence-derived peptides. <i>Journal of Immunology</i> , 2008 , 181, 5442-50	5.3	29
63	Fab antibody fragment-functionalized liposomes for specific targeting of antigen-positive cells. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2018 , 14, 123-130	6	28
62	Special organization of the HLA-G protein on the cell surface. <i>Human Immunology</i> , 2003 , 64, 1011-6	2.3	28
61	A novel anti-CD18 mAb recognizes an activation-related epitope and induces a high-affinity conformation in leukocyte integrins. <i>Immunobiology</i> , 2001 , 203, 687-98	3.4	28
60	Melanoma cells constitutively release an anchor-positive soluble form of protectin (sCD59) that retains functional activities in homologous complement-mediated cytotoxicity. <i>Journal of Clinical Investigation</i> , 1997 , 100, 1248-55	15.9	28
59	Regulation of Src family kinases involved in T cell receptor signaling by protein-tyrosine phosphatase CD148. <i>Journal of Biological Chemistry</i> , 2011 , 286, 22101-12	5.4	27
58	Amino acids at the N- and C-termini of human glutamate carboxypeptidase II are required for enzymatic activity and proper folding. <i>FEBS Journal</i> , 2004 , 271, 2782-90		27
57	Molecular mechanisms involved in CD43-mediated apoptosis of TF-1 cells. Roles of transcription Daxx expression, and adhesion molecules. <i>Journal of Biological Chemistry</i> , 2002 , 277, 7955-61	5.4	26
56	An alternative way of CD4 and CD8 association with protein kinases of the Src family. <i>Immunogenetics</i> , 1995 , 41, 110-6	3.2	25
55	LFA-1-mediated leukocyte adhesion regulated by interaction of CD43 with LFA-1 and CD147. <i>Molecular Immunology</i> , 2008 , 45, 1703-11	4.3	24
54	The nature of the subset of MHC class II molecules carrying the CDw78 epitopes. <i>International Immunology</i> , 1999 , 11, 491-8	4.9	24
53	Association of the GPI-anchored leucocyte surface glycoproteins with ganglioside GM3. <i>Biochemical and Biophysical Research Communications</i> , 1994 , 203, 1069-75	3.4	23
52	Association of human NK cell surface receptors NKR-P1 and CD94 with Src-family protein kinases. <i>Immunogenetics</i> , 1997 , 46, 231-6	3.2	22
51	The epitope recognized by pan-HLA class I-reactive monoclonal antibody W6/32 and its relationship to unusual stability of the HLA-B27/beta2-microglobulin complex. <i>Immunogenetics</i> , 2001 , 53, 440-6	3.2	22
50	Monoclonal antibodies against human alpha-fetoprotein. Exploitation of an unusual calcium-dependent interaction with the antigen for analytical and preparative purposes. <i>Journal of Immunological Methods</i> , 1988 , 111, 67-73	2.5	22

(2001-2013)

49	Nonredundant roles of Src-family kinases and Syk in the initiation of B-cell antigen receptor signaling. <i>Journal of Immunology</i> , 2013 , 190, 1807-18	5.3	21
48	The lipopolysaccharide co-receptor CD14 is present and functional in seminal plasma and expressed on spermatozoa. <i>Immunology</i> , 2001 , 104, 317-23	7.8	21
47	Dysregulation of Src family kinases in mast cells from epilepsy-resistant ASK versus epilepsy-prone EL mice. <i>Journal of Immunology</i> , 2007 , 178, 455-62	5.3	20
46	Adapters in lymphocyte signaling. <i>Journal of Clinical Investigation</i> , 2002 , 109, 301-9	15.9	20
45	Transient activation of the c-Jun N-terminal kinase (JNK) activity by ligation of the tetraspan CD53 antigen in different cell types. <i>FEBS Journal</i> , 2002 , 269, 1012-21		19
44	Interaction of late apoptotic and necrotic cells with vitronectin. <i>PLoS ONE</i> , 2011 , 6, e19243	3.7	17
43	LST1/A is a myeloid leukocyte-specific transmembrane adaptor protein recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to the plasma membrane. <i>Journal of Biological Chemistry</i> , 2012 , 287, 228	a∮2 ¹ -21	17
42	EVI2B is a C/EBPItarget gene required for granulocytic differentiation and functionality of hematopoietic progenitors. <i>Cell Death and Differentiation</i> , 2017 , 24, 705-716	12.7	16
41	Pre-sorting endosomal transport of the GPI-anchored protein, CD59, is regulated by EHD1. <i>Traffic</i> , 2011 , 12, 102-20	5.7	16
40	The human leucocyte antigen CD48 (MEM-102) is closely related to the activation marker Blast-1. <i>Immunogenetics</i> , 1991 , 33, 108-12	3.2	16
39	Lentil lectin effectively induces allotransplantation tolerance in mice. <i>Nature</i> , 1980 , 284, 273-5	50.4	16
38	Deletion of the LIME adaptor protein minimally affects T and B cell development and function. <i>European Journal of Immunology</i> , 2007 , 37, 3259-69	6.1	15
37	Human leukocytes contain a large pool of free forms of CD18. <i>Biochemical and Biophysical Research Communications</i> , 2000 , 275, 295-9	3.4	15
36	The effects of membrane compartmentalization of csk on TCR signaling. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2011 , 1813, 367-76	4.9	13
35	Affinity electrophoresis. <i>Trends in Biochemical Sciences</i> , 1979 , 4, N6-N7	10.3	13
34	Affinity electrophoresis: new simple and general methods of preparation of affinity gels. <i>Analytical Biochemistry</i> , 1982 , 125, 358-69	3.1	13
33	CDw78a determinant on a major histocompatibility complex class II subpopulation that can be induced to associate with the cytoskeleton. <i>European Journal of Immunology</i> , 1997 , 27, 3206-13	6.1	12
32	Production of HIV-1 by resting memory T lymphocytes. <i>Aids</i> , 2001 , 15, 1931-40	3.5	12

31	Genomic structure of the human CD53 gene. <i>Immunogenetics</i> , 1993 , 38, 272-9	3.2	12
30	The genes for CD37, CD53, and R2, all members of a novel gene family, are located on different chromosomes. <i>Immunogenetics</i> , 1993 , 37, 461-5	3.2	12
29	Human leucocyte surface glycoprotein CDw44 and lymphocyte homing receptor are identical molecules. <i>Immunogenetics</i> , 1989 , 29, 402-4	3.2	12
28	PRR7 is a transmembrane adaptor protein expressed in activated T cells involved in regulation of T cell receptor signaling and apoptosis. <i>Journal of Biological Chemistry</i> , 2011 , 286, 19617-29	5.4	10
27	Characterization of seven new monoclonal antibodies against human DR, DR + DP and DQ1 + DQ3 antigens. <i>Tissue Antigens</i> , 1986 , 28, 288-97		10
26	Expression pattern of adaptor protein PAG: correlation between secondary lymphatic follicle and histogenetically related malignant lymphomas. <i>Immunology Letters</i> , 2005 , 100, 94-7	4.1	10
25	The use of styrene-maleic acid copolymer (SMA) for studies on T cell membrane rafts. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2019 , 1861, 130-141	3.8	9
24	Murine hybridoma monoclonal antibodies against insulin: cross-reactivity with insulins of three species and blocking of insulin binding to its receptor. <i>Immunology Letters</i> , 1984 , 8, 279-83	4.1	9
23	Equilibrium in the protein-immobilized-ligand-soluble-ligand system: estimation of dissociation constants of protein-soluble-ligand complexes from binding-inhibition data. <i>Molecular Immunology</i> , 1985 , 22, 125-33	4.3	9
22	LATan important raft-associated transmembrane adaptor protein. Delivered on 6 July 2009 at the 34th FEBS Congress in Prague, Czech Republic. <i>FEBS Journal</i> , 2010 , 277, 4383-97	5.7	7
21	UniCAR T cell immunotherapy enables efficient elimination of radioresistant cancer cells. <i>OncoImmunology</i> , 2020 , 9, 1743036	7.2	6
20	CD Antigens 2001. <i>Modern Pathology</i> , 2002 , 15, 71-6	9.8	6
19	CDw149 antibodies recognize a clustered subset of CD47 molecules associated with cytoplasmic signaling molecules. <i>Tissue Antigens</i> , 2000 , 56, 258-67		6
18	The CBF.78 monoclonal antibody to human sialophorin has distinct properties giving new insights into the CD43 marker and its activation pathway. <i>Tissue Antigens</i> , 1999 , 54, 1-15		6
17	Simple polyacrylamide gel electrophoresis in continuous carbonate buffer system suitable for the analysis of ascitic fluids of hybridoma bearing mice. <i>Journal of Immunological Methods</i> , 1986 , 86, 103-5	2.5	6
16	The transmembrane region is responsible for targeting of adaptor protein LAX into "heavy rafts". <i>PLoS ONE</i> , 2012 , 7, e36330	3.7	4
15	Association of leukocyte surface receptors with protein kinases. <i>International Archives of Allergy and Immunology</i> , 1996 , 110, 1-6	3.7	4
14	Exogenous CD59 incorporated into U937 cells through its glycosyl phosphatidylinositol anchor becomes associated with signalling molecules in a time dependent manner. <i>Biochemical Society Transactions</i> , 1995 , 23, 269S	5.1	4

LIST OF PUBLICATIONS

1	13	Dendritic cells sensitize TCRs through self-MHC-mediated Src family kinase activation. <i>Journal of Immunology</i> , 2007 , 178, 2262-71	5.3	3	
1	2	The 5th EFIS Tatra Immunology Conference on RMolecular determinants of T cell immunityRheld in the High Tatra Mountains, Slovakia, September 7-11, 2002. <i>Immunology Letters</i> , 2003 , 86, 1-6	4.1	3	
1	1	The HLDA8 blind panel: findings and conclusions. <i>Journal of Immunological Methods</i> , 2005 , 305, 75-83	2.5	3	
1	Ο.	Cross-reactivity between tubulin and denatured human serum albumin demonstrated by monoclonal antibody TU-01. <i>Immunology Letters</i> , 1984 , 8, 285-8	4.1	3	
9)	The mannose 6-phosphate/insulin-like growth factor 2 receptor mediates plasminogen-induced efferocytosis. <i>Journal of Leukocyte Biology</i> , 2019 , 105, 519-530	6.5	2	
8	3	Czech bibliometric system fosters mediocre research. <i>Nature</i> , 2009 , 460, 1079	50.4	1	
7	7	A clustered subset of MHC class II molecules. <i>Trends in Immunology</i> , 1998 , 19, 486		1	
6	5	A novel family of leucocyte surface antigens. <i>Trends in Immunology</i> , 1991 , 12, 287		1	
5	;	Czech science. <i>Nature</i> , 1992 , 359, 99-99	50.4	1	
4	ł	Development of Novel Anti-CD10 Target Modules for Redirection of Universal CAR T Cells Against CD10-Positive Malignancies. <i>Blood</i> , 2019 , 134, 5612-5612	2.2	1	
3	}	Tumour devascularisation as a potential immunotherapeutic strategy. <i>Oncolmmunology</i> , 2019 , 8, e1526	56 / 1. <u>4</u>		
2	2	A novel monoclonal reagent recognizing native and denatured Vbeta5.3-related chains of human T cell receptor. <i>Immunology Letters</i> , 2003 , 88, 105-8	4.1		
1		Non-lineage antigens: section report. <i>Cellular Immunology</i> , 2005 , 236, 42-7	4.4		