

Camilla Pramfalk

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9540296/publications.pdf>

Version: 2024-02-01

27
papers

738
citations

471061

17
h-index

552369

26
g-index

28
all docs

28
docs citations

28
times ranked

1525
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>Soat2</i> ties cholesterol metabolism to β -oxidation and glucose tolerance in male mice. <i>Journal of Internal Medicine</i> , 2022, 292, 296-307.	2.7	6
2	Abstract 10013: PRDs Are Multifunctional Oral Inhibitors of PCSK9 and ACAT2. <i>Circulation</i> , 2021, 144, .	1.6	0
3	Modifying nutritional substrates induces macrovesicular lipid droplet accumulation and metabolic alterations in a cellular model of hepatic steatosis. <i>Physiological Reports</i> , 2020, 8, e14482.	0.7	7
4	Generation of new hepatocyte-like in vitro models better resembling human lipid metabolism. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020, 1865, 158659.	1.2	2
5	Hepatic de novo lipogenesis is suppressed and fat oxidation is increased by omega-3 fatty acids at the expense of glucose metabolism. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e000871.	1.2	46
6	Effects on hepatic lipid metabolism in human hepatoma cells following overexpression of TGF β 2 induced factor homeobox 1 or 2. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019, 1864, 756-762.	1.2	3
7	Fasting hepatic de novo lipogenesis is not reliably assessed using circulating fatty acid markers. <i>American Journal of Clinical Nutrition</i> , 2019, 109, 260-268.	2.2	21
8	Ezetimibe in Combination With Simvastatin Reduces Remnant Cholesterol Without Affecting Biliary Lipid Concentrations in Gallstone Patients. <i>Journal of the American Heart Association</i> , 2018, 7, e009876.	1.6	24
9	Overexpression of transforming growth factor β 2 induced factor homeobox 1 represses NPC1L1 and lowers markers of intestinal cholesterol absorption. <i>Atherosclerosis</i> , 2018, 275, 246-255.	0.4	4
10	In vitro cellular models of human hepatic fatty acid metabolism: differences between Huh7 and HepG2 cell lines in human and fetal bovine culturing serum. <i>Physiological Reports</i> , 2017, 5, e13532.	0.7	48
11	Fasting Plasma Insulin Concentrations Are Associated With Changes in Hepatic Fatty Acid Synthesis and Partitioning Prior to Changes in Liver Fat Content in Healthy Adults. <i>Diabetes</i> , 2016, 65, 1858-1867.	0.3	37
12	Diabetes Mellitus Is Associated With Reduced High-Density Lipoprotein Sphingosine-1-Phosphate Content and Impaired High-Density Lipoprotein Cardiac Cell Protection. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 817-824.	1.1	61
13	Culturing of HepG2 cells with human serum improve their functionality and suitability in studies of lipid metabolism. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 51-59.	1.2	11
14	Sex-Specific Differences in Hepatic Fat Oxidation and Synthesis May Explain the Higher Propensity for NAFLD in Men. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 4425-4433.	1.8	108
15	From whole body to cellular models of hepatic triglyceride metabolism: man has got to know his limitations. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2015, 308, E1-E20.	1.8	30
16	Role of TG-interacting factor (Tgif) in lipid metabolism. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2015, 1851, 9-12.	1.2	7
17	TG-interacting factor 1 acts as a transcriptional repressor of sterol O-acyltransferase 2. <i>Journal of Lipid Research</i> , 2014, 55, 709-717.	2.0	11
18	The storage stability and concentration of acetoacetate differs between blood fractions. <i>Clinica Chimica Acta</i> , 2014, 433, 278-283.	0.5	18

#	ARTICLE	IF	CITATIONS
19	Estrogen Signalling and the Metabolic Syndrome: Targeting the Hepatic Estrogen Receptor Alpha Action. PLoS ONE, 2013, 8, e57458.	1.1	46
20	Cholesteryl esters and ACAT. European Journal of Lipid Science and Technology, 2012, 114, 624-633.	1.0	21
21	Role of thyroid receptor β 2 in lipid metabolism. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2011, 1812, 929-937.	1.8	48
22	Hepatic Niemann-Pick C1-like 1. Current Opinion in Lipidology, 2011, 22, 225-230.	1.2	25
23	HNF1 α and SREBP2 are important regulators of NPC1L1 in human liver. Journal of Lipid Research, 2010, 51, 1354-1362.	2.0	32
24	HNF1 α and SREBP2 are important regulators of NPC1L1 in human liver. Journal of Lipid Research, 2010, 51, 1354-1362.	2.0	46
25	Thyroid hormones and thyroid hormone receptors: effects of thyromimetics on reverse cholesterol transport. World Journal of Gastroenterology, 2010, 16, 5958-64.	1.4	26
26	Cholesterol regulates ACAT2 gene expression and enzyme activity in human hepatoma cells. Biochemical and Biophysical Research Communications, 2007, 364, 402-409.	1.0	22
27	Control of ACAT2 liver expression by HNF1. Journal of Lipid Research, 2005, 46, 1868-1876.	2.0	28