

# Christine M O connor

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/9547689/christine-m-oconnor-publications-by-year.pdf>

**Version:** 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

28  
papers

624  
citations

13  
h-index

24  
g-index

33  
ext. papers

750  
ext. citations

5.9  
avg, IF

4.62  
L-index

#	Paper	IF	Citations
28	Modulation of host cell signaling during cytomegalovirus latency and reactivation. <i>Virology Journal</i> , <b>2021</b> , 18, 207	6.1	1
27	CMV-encoded GPCR pUL33 activates CREB and facilitates its recruitment to the MIE locus for efficient viral reactivation. <i>Journal of Cell Science</i> , <b>2021</b> , 134,	5.3	6
26	Methods for Studying the Function of Cytomegalovirus GPCRs. <i>Methods in Molecular Biology</i> , <b>2021</b> , 2244, 159-197	1.4	
25	Innate Immune Responses to Herpesvirus Infection. <i>Cells</i> , <b>2021</b> , 10,	7.9	2
24	Regulation of the MIE Locus During HCMV Latency and Reactivation. <i>Pathogens</i> , <b>2020</b> , 9,	4.5	9
23	The Requirement for US28 During Cytomegalovirus Latency Is Independent of US27 and US29 Gene Expression. <i>Frontiers in Cellular and Infection Microbiology</i> , <b>2020</b> , 10, 186	5.9	6
22	Identification of a novel signaling complex containing host chemokine receptor CXCR4, Interleukin-10 receptor, and human cytomegalovirus US27. <i>Virology</i> , <b>2020</b> , 548, 49-58	3.6	1
21	Activator protein-1 transactivation of the major immediate early locus is a determinant of cytomegalovirus reactivation from latency. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2020</b> , 117, 20860-20867	11.5	10
20	Interferon-Responsive Genes Are Targeted during the Establishment of Human Cytomegalovirus Latency. <i>MBio</i> , <b>2019</b> , 10,	7.8	22
19	Human cytomegalovirus G protein-coupled receptor US28 promotes latency by attenuating c-fos. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2019</b> , 116, 1755-1764	11.5	36
18	Human Cytomegalovirus UL111A and US27 Gene Products Enhance the CXCL12/CXCR4 Signaling Axis via Distinct Mechanisms. <i>Journal of Virology</i> , <b>2018</b> , 92,	6.6	13
17	Selective 4-Thiouracil Labeling of RNA Transcripts within Latently Infected Cells after Infection with Human Cytomegalovirus Expressing Functional Uracil Phosphoribosyltransferase. <i>Journal of Virology</i> , <b>2018</b> , 92,	6.6	10
16	US28: HCMV's Swiss Army Knife. <i>Viruses</i> , <b>2018</b> , 10,	6.2	37
15	The Natural Flavonoid Compound Deguelin Inhibits HCMV Lytic Replication within Fibroblasts. <i>Viruses</i> , <b>2018</b> , 10,	6.2	13
14	The Human Cytomegalovirus US27 Gene Product Constitutively Activates Antioxidant Response Element-Mediated Transcription through G $\beta$ Phosphoinositide 3-Kinase, and Nuclear Respiratory Factor 1. <i>Journal of Virology</i> , <b>2018</b> , 92,	6.6	9
13	Inhibition of the FACT Complex Reduces Transcription from the Human Cytomegalovirus Major Immediate Early Promoter in Models of Lytic and Latent Replication. <i>Journal of Virology</i> , <b>2016</b> , 90, 4249-4253	6.6	19
12	Human Cytomegalovirus US28 Is Important for Latent Infection of Hematopoietic Progenitor Cells. <i>Journal of Virology</i> , <b>2015</b> , 90, 2959-70	6.6	74

11	Four levels of hierarchical organization, including noncovalent chainmail, brace the mature tumor herpesvirus capsid against pressurization. <i>Structure</i> , <b>2014</b> , 22, 1385-98	5.2	14
10	Quantitative proteomic discovery of dynamic epigenome changes that control human cytomegalovirus (HCMV) infection. <i>Molecular and Cellular Proteomics</i> , <b>2014</b> , 13, 2399-410	7.6	25
9	Host microRNA regulation of human cytomegalovirus immediate early protein translation promotes viral latency. <i>Journal of Virology</i> , <b>2014</b> , 88, 5524-32	6.6	71
8	Methods for studying the function of cytomegalovirus GPCRs. <i>Methods in Molecular Biology</i> , <b>2014</b> , 1119, 133-64	1.4	4
7	US28 is a potent activator of phospholipase C during HCMV infection of clinically relevant target cells. <i>PLoS ONE</i> , <b>2012</b> , 7, e50524	3.7	37
6	A myeloid progenitor cell line capable of supporting human cytomegalovirus latency and reactivation, resulting in infectious progeny. <i>Journal of Virology</i> , <b>2012</b> , 86, 9854-65	6.6	91
5	Human cytomegalovirus pUL78 G protein-coupled receptor homologue is required for timely cell entry in epithelial cells but not fibroblasts. <i>Journal of Virology</i> , <b>2012</b> , 86, 11425-33	6.6	40
4	Human cytomegalovirus pUS27 G protein-coupled receptor homologue is required for efficient spread by the extracellular route but not for direct cell-to-cell spread. <i>Journal of Virology</i> , <b>2011</b> , 85, 3700-7	6.6	70
3	The Requirement For US28 During Cytomegalovirus Latency Is Independent Of US27 And US29 Gene Expression		1
2	Activator protein-1 transactivation of the major immediate early locus is a determinant of cytomegalovirus reactivation from latency		1
1	The human cytomegalovirus G-protein coupled receptor US28 promotes latency by attenuating c-fos		1