## Christopher A Natale

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9550625/publications.pdf

Version: 2024-02-01

1040056 1474206 11 582 9 9 citations g-index h-index papers 17 17 17 1299 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	MLL1 is essential for the senescence-associated secretory phenotype. Genes and Development, 2016, 30, 321-336.	5.9	121
2	Activation of G protein-coupled estrogen receptor signaling inhibits melanoma and improves response to immune checkpoint blockade. ELife, $2018, 7, \ldots$	6.0	98
3	Sex steroids regulate skin pigmentation through nonclassical membrane-bound receptors. ELife, 2016, 5, .	6.0	89
4	$\langle i \rangle$ CDKN2B $\langle i \rangle$ Loss Promotes Progression from Benign Melanocytic Nevus to Melanoma. Cancer Discovery, 2015, 5, 1072-1085.	9.4	78
5	Systematic Epigenomic Analysis Reveals Chromatin States Associated with Melanoma Progression. Cell Reports, 2017, 19, 875-889.	6.4	78
6	Pharmacologic Activation of the G Protein–Coupled Estrogen Receptor Inhibits Pancreatic Ductal Adenocarcinoma. Cellular and Molecular Gastroenterology and Hepatology, 2020, 10, 868-880.e1.	4.5	35
7	Four Previously Identified Petunia inflata S-Locus F-Box Genes Are Involved in Pollen Specificity in Self-Incompatibility. Molecular Plant, 2014, 7, 567-569.	8.3	33
8	The integrin $\hat{l}\pm v$ -TGF $\hat{l}^2$ signaling axis is necessary for epidermal proliferation during cutaneous wound healing. Cell Cycle, 2016, 15, 2077-2086.	2.6	29
9	ZIP9 Is a Druggable Determinant of Sex Differences in Melanoma. Cancer Research, 2021, 81, 5991-6003.	0.9	14
10	Abstract 1239: CDKN2B loss promotes progression from benign melanocytic nevus to melanoma. , 2015, , .		3
11	Non-Classical Estrogen Signaling Inhibits Melanoma and Improves Response to PD-1 Blockade. SSRN Electronic Journal, 0, , .	0.4	1