

# Miwako Yamasaki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9562896/publications.pdf>

Version: 2024-02-01

78  
papers

4,393  
citations

126907

33  
h-index

114465

63  
g-index

89  
all docs

89  
docs citations

89  
times ranked

6201  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | L-DOPA-Induced Neurogenesis in the Hippocampus Is Mediated Through GPR143, a Distinct Mechanism of Dopamine. <i>Stem Cells</i> , 2022, 40, 215-226.   | 3.2 | 5         |
| 2  | SIPA1L1/SPAR1 Interacts with the Neurabin Family of Proteins and is Involved in GPCR Signaling. <i>Journal of Neuroscience</i> , 2022, 42, 2448-2473.   | 3.6 | 5         |
| 3  | Activation of Extrasynaptic Kainate Receptors Drives Hilar Mossy Cell Activity. <i>Journal of Neuroscience</i> , 2022, 42, 2872-2884.   | 3.6 | 8         |
| 4  | Histochemical Characterization of the Dorsal Raphe-Periaqueductal Grey Dopamine Transporter Neurons Projecting to the Extended Amygdala. <i>ENeuro</i> , 2022, 9, ENEURO.0121-22.2022.  | 1.9 | 4         |
| 5  | Single-scan volumetric imaging throughout thick tissue specimens by one-touch installable light-needle creating device. <i>Scientific Reports</i> , 2022, 12, .   | 3.3 | 4         |
| 6  | Compartmentalized Inputâ€œOutput Organization of Lugaro Cells in the Cerebellar Cortex. <i>Neuroscience</i> , 2021, 462, 89-105.  | 2.3 | 15        |
| 7  | Nectinâ€² is localized at cholinergic neuron dendrites and regulates synapse formation in the medial habenula. <i>Journal of Comparative Neurology</i> , 2021, 529, 450-477.  | 1.6 | 4         |
| 8  | Kv11 (related gene) voltage-dependent K <sup>+</sup> channels promote resonance and oscillation of subthreshold membrane potentials. <i>Journal of Physiology</i> , 2021, 599, 547-569.   | 2.9 | 7         |
| 9  | mGluR1 signaling in cerebellar Purkinje cells: Subcellular organization and involvement in cerebellar function and disease. <i>Neuropharmacology</i> , 2021, 194, 108629.   | 4.1 | 16        |
| 10 | Fluorescent In Situ Hybridization for Sensitive and Specific Labeling. <i>Neuromethods</i> , 2021, , 145-160.   | 0.3 | 0         |
| 11 | Spike firing attenuation of serotonin neurons in learned helplessness rats is reversed by ketamine. <i>Brain Communications</i> , 2021, 3, fcab285.   | 3.3 | 2         |
| 12 | Expression mapping, quantification, and complex formation of GluD1 and GluD2 glutamate receptors in adult mouse brain. <i>Journal of Comparative Neurology</i> , 2020, 528, 1003-1027.  | 1.6 | 33        |
| 13 | TMEM163 Regulates ATP-Gated P2X Receptor and Behavior. <i>Cell Reports</i> , 2020, 31, 107704.  | 6.4 | 19        |
| 14 | Development of an L-type Ca <sup>2+</sup> channel-dependent Ca <sup>2+</sup> transient during the radial migration of cortical excitatory neurons. <i>Neuroscience Research</i> , 2020, 169, 17-26.   | 1.9 | 8         |
| 15 | mGluR1 in cerebellar Purkinje cells is essential for the formation but not expression of associative eyeblink memory. <i>Scientific Reports</i> , 2019, 9, 7353.  | 3.3 | 10        |
| 16 | Localization of phospholipase C Î²3 in the major salivary glands of adult mice. <i>Acta Histochemica</i> , 2019, 121, 484-490.  | 1.8 | 6         |
| 17 | Heterogeneous localization of muscarinic cholinceptor M1 in the salivary ducts of adult mice. <i>Archives of Oral Biology</i> , 2019, 100, 14-22.   | 1.8 | 3         |
| 18 | Localization of nectinâ€² at the boundary between the adjacent somata of the clustered cholinergic neurons and its regulatory role in the subcellular localization of the voltage-gated A-type K <sup>+</sup> channel Kv4.2 in the medial habenula. <i>Journal of Comparative Neurology</i> , 2018, 526, 1527-1549. | 1.6 | 4         |

| #  | ARTICLE   | IF   | CITATIONS |
|----|---|------|-----------|
| 19 | Silent Learning. <i>Current Biology</i> , 2018, 28, 3508-3515.e5.   | 3.9  | 35        |
| 20 | Olig2-Lineage Astrocytes: A Distinct Subtype of Astrocytes That Differs from GFAP Astrocytes. <i>Frontiers in Neuroanatomy</i> , 2018, 12, 8.   | 1.7  | 71        |
| 21 | Localization of photoperiod responsive circadian oscillators in the mouse suprachiasmatic nucleus. <i>Scientific Reports</i> , 2017, 7, 8210.   | 3.3  | 31        |
| 22 | Glutamate transporter GLAST controls synaptic wrapping by Bergmann glia and ensures proper wiring of Purkinje cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 7438-7443.                           | 7.1  | 54        |
| 23 | Locus Coeruleus and Dopamine-Dependent Memory Consolidation. <i>Neural Plasticity</i> , 2017, 2017, 1-15.   | 2.2  | 50        |
| 24 | TARP $\hat{\imath}^3$ -2 and $\hat{\imath}^3$ -8 Differentially Control AMPAR Density Across Schaffer Collateral/Commissural Synapses in the Hippocampal CA1 Area. <i>Journal of Neuroscience</i> , 2016, 36, 4296-4312.                                      | 3.6  | 36        |
| 25 | Transsynaptic Modulation of Kainate Receptor Functions by C1q-like Proteins. <i>Neuron</i> , 2016, 90, 752-767.   | 8.1  | 150       |
| 26 | Locus coeruleus and dopaminergic consolidation of everyday memory. <i>Nature</i> , 2016, 537, 357-362.  | 27.8 | 561       |
| 27 | Distinct Subunit Domains Govern Synaptic Stability and Specificity of the Kainate Receptor. <i>Cell Reports</i> , 2016, 16, 531-544.  | 6.4  | 33        |
| 28 | Ionic Basis for Membrane Potential Resonance in Neurons of the Inferior Olive. <i>Cell Reports</i> , 2016, 16, 994-1004.  | 6.4  | 32        |
| 29 | The active zone protein $\langle scp \rangle$ CAST $\langle /scp \rangle$ regulates synaptic vesicle recycling and quantal size in the mouse hippocampus. <i>European Journal of Neuroscience</i> , 2016, 44, 2272-2284.                                      | 2.6  | 17        |
| 30 | Molecular and anatomical evidence for the input pathway- and target cell type-dependent regulation of glutamatergic synapses. <i>Anatomical Science International</i> , 2016, 91, 8-21.   | 1.0  | 4         |
| 31 | Developmental Switch in Spike Timing-Dependent Plasticity and Cannabinoid-Dependent Reorganization of the Thalamocortical Projection in the Barrel Cortex. <i>Journal of Neuroscience</i> , 2016, 36, 7039-7054.  | 3.6  | 18        |
| 32 | Territories of heterologous inputs onto Purkinje cell dendrites are segregated by mGluR1-dependent parallel fiber synapse elimination. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 2282-2287.         | 7.1  | 66        |
| 33 | Fluorescent In Situ Hybridization for Sensitive and Specific Labeling. <i>Neuromethods</i> , 2016, , 127-142.   | 0.3  | 2         |
| 34 | QRFP-Deficient Mice Are Hypophagic, Lean, Hypoactive and Exhibit Increased Anxiety-Like Behavior. <i>PLoS ONE</i> , 2016, 11, e0164716.   | 2.5  | 28        |
| 35 | VGlut3-Expressing CCK-Positive Basket Cells Construct Invaginating Synapses Enriched with Endocannabinoid Signaling Proteins in Particular Cortical and Cortex-Like Amygdaloid Regions of Mouse Brains. <i>Journal of Neuroscience</i> , 2015, 35, 4215-4228. | 3.6  | 33        |
| 36 | Involvement of diacylglycerol kinase $\hat{\imath}^2$ in the spine formation at distal dendrites of striatal medium spiny neurons. <i>Brain Research</i> , 2015, 1594, 36-45.   | 2.2  | 11        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | The glutamate receptor $\text{GluN}2$ subunit regulates synaptic trafficking of AMPA receptors in the neonatal mouse brain. <i>European Journal of Neuroscience</i> , 2014, 40, 3136-3146.   | 2.6  | 14        |
| 38 | Neuron type- and input pathway-dependent expression of Slc4a10 in adult mouse brains. <i>European Journal of Neuroscience</i> , 2014, 40, 2797-2810.   | 2.6  | 4         |
| 39 | Enriched Expression of GluD1 in Higher Brain Regions and Its Involvement in Parallel Fiber Interneuron Synapse Formation in the Cerebellum. <i>Journal of Neuroscience</i> , 2014, 34, 7412-7424.  | 3.6  | 89        |
| 40 | Opposing Role of NMDA Receptor GluN2B and GluN2D in Somatosensory Development and Maturation. <i>Journal of Neuroscience</i> , 2014, 34, 11534-11548.  | 3.6  | 49        |
| 41 | Global Scaling Down of Excitatory Postsynaptic Responses in Cerebellar Purkinje Cells Impairs Developmental Synapse Elimination. <i>Cell Reports</i> , 2014, 8, 1119-1129.   | 6.4  | 19        |
| 42 | Protocadherin 17 Regulates Presynaptic Assembly in Topographic Corticobasal Ganglia Circuits. <i>Neuron</i> , 2013, 78, 839-854.   | 8.1  | 67        |
| 43 | Homeostatic Control of Synaptic Transmission by Distinct Glutamate Receptors. <i>Neuron</i> , 2013, 78, 687-699.   | 8.1  | 31        |
| 44 | Type 2 $\text{K}^+$ cotransporter is preferentially recruited to climbing fiber synapses during development and the stellate cell targeting dendritic zone at adulthood in cerebellar Purkinje cells. <i>European Journal of Neuroscience</i> , 2013, 37, 532-543. | 2.6  | 8         |
| 45 | Autoantibodies to Epilepsy-Related LGI1 in Limbic Encephalitis Neutralize LGI1-ADAM22 Interaction and Reduce Synaptic AMPA Receptors. <i>Journal of Neuroscience</i> , 2013, 33, 18161-18174.  | 3.6  | 288       |
| 46 | Three Types of Neurochemical Projection from the Bed Nucleus of the Stria Terminalis to the Ventral Tegmental Area in Adult Mice. <i>Journal of Neuroscience</i> , 2012, 32, 18035-18046.  | 3.6  | 158       |
| 47 | $\text{Ca}^{v}2.1$ in Cerebellar Purkinje Cells Regulates Competitive Excitatory Synaptic Wiring, Cell Survival, and Cerebellar Biochemical Compartmentalization. <i>Journal of Neuroscience</i> , 2012, 32, 1311-1328.  | 3.6  | 74        |
| 48 | Lack of Molecular-Anatomical Evidence for GABAergic Influence on Axon Initial Segment of Cerebellar Purkinje Cells by the Pinceau Formation. <i>Journal of Neuroscience</i> , 2012, 32, 9438-9448.   | 3.6  | 47        |
| 49 | Distinct Neurochemical and Functional Properties of GAD67-Containing 5-HT Neurons in the Rat Dorsal Raphe Nucleus. <i>Journal of Neuroscience</i> , 2012, 32, 14415-14426.   | 3.6  | 47        |
| 50 | Rewiring of Afferent Fibers in the Somatosensory Thalamus of Mice Caused by Peripheral Sensory Nerve Transection. <i>Journal of Neuroscience</i> , 2012, 32, 6917-6930.  | 3.6  | 25        |
| 51 | Glutamate receptor $\gamma 2$ is essential for input pathway-dependent regulation of synaptic AMPAR contents in cerebellar Purkinje cells. <i>Neuroscience Research</i> , 2011, 71, e93.   | 1.9  | 0         |
| 52 | Invaginating inhibitory synapse with particularly rich endocannabinoid signaling machinery in the basal nucleus of the amygdala. <i>Neuroscience Research</i> , 2011, 71, e93-e94.   | 1.9  | 0         |
| 53 | Cellular expression and subcellular localization of secretogranin II in the mouse hippocampus and cerebellum. <i>European Journal of Neuroscience</i> , 2011, 33, 82-94.   | 2.6  | 14        |
| 54 | Distinct functions of kainate receptors in the brain are determined by the auxiliary subunit Neto1. <i>Nature Neuroscience</i> , 2011, 14, 866-873.  | 14.8 | 111       |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Molecular and Morphological Configuration for 2-Arachidonoylglycerol-Mediated Retrograde Signaling at Mossy Cell-Granule Cell Synapses in the Dentate Gyrus. <i>Journal of Neuroscience</i> , 2011, 31, 7700-7714.   | 3.6 | 75        |
| 56 | Glutamate Receptor $\hat{2}$ Is Essential for Input Pathway-Dependent Regulation of Synaptic AMPAR Contents in Cerebellar Purkinje Cells. <i>Journal of Neuroscience</i> , 2011, 31, 3362-3374.  | 3.6 | 79        |
| 57 | Unique inhibitory synapse with particularly rich endocannabinoid signaling machinery on pyramidal neurons in basal amygdaloid nucleus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 3059-3064.        | 7.1 | 100       |
| 58 | Developmental Switching of Perisomatic Innervation from Climbing Fibers to Basket Cell Fibers in Cerebellar Purkinje Cells. <i>Journal of Neuroscience</i> , 2011, 31, 16916-16927.  | 3.6 | 52        |
| 59 | Orexin Neurons Receive Glycinergic Innervations. <i>PLoS ONE</i> , 2011, 6, e25076.  | 2.5 | 26        |
| 60 | TARPs $\hat{2}$ and $\hat{7}$ are essential for AMPA receptor expression in the cerebellum. <i>European Journal of Neuroscience</i> , 2010, 31, 2204-2220.   | 2.6 | 76        |
| 61 | Cytochemical and cytological properties of perineuronal oligodendrocytes in the mouse cortex. <i>European Journal of Neuroscience</i> , 2010, 32, 1326-1336.   | 2.6 | 81        |
| 62 | Inositol 1,4,5-trisphosphate signaling maintains the activity of glutamate uptake in Bergmann glia. <i>European Journal of Neuroscience</i> , 2010, 32, 1668-1677.   | 2.6 | 19        |
| 63 | Ablation of Glutamate Receptor GluR $\hat{2}$ in Adult Purkinje Cells Causes Multiple Innervation of Climbing Fibers by Inducing Aberrant Invasion to Parallel Fiber Innervation Territory. <i>Journal of Neuroscience</i> , 2010, 30, 15196-15209.          | 3.6 | 55        |
| 64 | Preferential Localization of Muscarinic M <sub>1</sub> Receptor on Dendritic Shaft and Spine of Cortical Pyramidal Cells and Its Anatomical Evidence for Volume Transmission. <i>Journal of Neuroscience</i> , 2010, 30, 4408-4418.                          | 3.6 | 187       |
| 65 | Imaging extrasynaptic glutamate dynamics in the brain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 6526-6531.  | 7.1 | 156       |
| 66 | Ablation of glutamate receptor GluR2 in adult Purkinje cells causes multiple innervation of climbing fibers by ectopic innervation of transverse collaterals. <i>Neuroscience Research</i> , 2010, 68, e86.  | 1.9 | 1         |
| 67 | Lentiviral vector-mediated rescue of motor behavior in spontaneously occurring hereditary ataxic mice. <i>Neurobiology of Disease</i> , 2009, 35, 457-465.   | 4.4 | 13        |
| 68 | Rescue of abnormal phenotypes in $\hat{2}$ glutamate receptor-deficient mice by the extracellular N-terminal and intracellular C-terminal domains of the $\hat{2}$ glutamate receptor. <i>European Journal of Neuroscience</i> , 2009, 30, 355-365.          | 2.6 | 21        |
| 69 | NMDA Receptor GluN2B (GluR $\hat{2}$ /NR2B) Subunit Is Crucial for Channel Function, Postsynaptic Macromolecular Organization, and Actin Cytoskeleton at Hippocampal CA3 Synapses. <i>Journal of Neuroscience</i> , 2009, 29, 10869-10882.                   | 3.6 | 138       |
| 70 | Role of the internal Shank-binding segment of glutamate receptor $\hat{2}$ in synaptic localization and cerebellar functions. <i>Neuroscience Letters</i> , 2008, 433, 146-151.  | 2.1 | 10        |
| 71 | Use-dependent amplification of presynaptic Ca <sup>2+</sup> signaling by axonal ryanodine receptors at the hippocampal mossy fiber synapse. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 11998-12003. | 7.1 | 55        |
| 72 | Glutamate Transporters Regulate Lesion-Induced Plasticity in the Developing Somatosensory Cortex. <i>Journal of Neuroscience</i> , 2008, 28, 4995-5006.  | 3.6 | 49        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | Regulation of Long-Term Depression and Climbing Fiber Territory by Glutamate Receptor $\hat{2}$ at Parallel Fiber Synapses through its C-Terminal Domain in Cerebellar Purkinje Cells. <i>Journal of Neuroscience</i> , 2007, 27, 12096-12108.        | 3.6 | 56        |
| 74 | Miniature Synaptic Events Elicited by Presynaptic $Ca^{2+}$ Rise Are Selectively Suppressed by Cannabinoid Receptor Activation in Cerebellar Purkinje Cells. <i>Journal of Neuroscience</i> , 2006, 26, 86-95.  | 3.6 | 64        |
| 75 | 3-Phosphoglycerate Dehydrogenase, a Key Enzyme for Serine Biosynthesis, Is Preferentially Expressed in the Radial Glia/Astrocyte Lineage and Olfactory Ensheathing Glia in the Mouse Brain. <i>Journal of Neuroscience</i> , 2001, 21, 7691-7704.     | 3.6 | 186       |
| 76 | Gq protein $\hat{1}$ subunits $G\hat{1}\pm q$ and $G\hat{1}\pm 11$ are localized at postsynaptic extra-junctional membrane of cerebellar Purkinje cells and hippocampal pyramidal cells. <i>European Journal of Neuroscience</i> , 2000, 12, 781-792. | 2.6 | 118       |
| 77 | Critical Period for Activity-Dependent Synapse Elimination in Developing Cerebellum. <i>Journal of Neuroscience</i> , 2000, 20, 4954-4961.  | 3.6 | 166       |
| 78 | L-Serine and glycine serve as major astroglia-derived trophic factors for cerebellar Purkinje neurons. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 11528-11533.                                | 7.1 | 175       |