

Catharina Lavebratt

List of Publications by Citations

Source: <https://exaly.com/author-pdf/9568791/catharina-lavebratt-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

133
papers

4,276
citations

34
h-index

62
g-index

149
ext. papers

5,667
ext. citations

6.4
avg, IF

5.16
L-index

#	Paper	IF	Citations
133	Genome-wide association study identifies 30 loci associated with bipolar disorder. <i>Nature Genetics</i> , 2019 , 51, 793-803	36.3	662
132	Genetic variants associated with response to lithium treatment in bipolar disorder: a genome-wide association study. <i>Lancet, The</i> , 2016 , 387, 1085-1093	4.0	216
131	Genetic and epigenetic associations of MAOA and NR3C1 with depression and childhood adversities. <i>International Journal of Neuropsychopharmacology</i> , 2013 , 16, 1513-28	5.8	153
130	Epigenetic aberrations in leukocytes of patients with schizophrenia: association of global DNA methylation with antipsychotic drug treatment and disease onset. <i>FASEB Journal</i> , 2012 , 26, 2712-8	0.9	144
129	Genome-wide association study of 40,000 individuals identifies two novel loci associated with bipolar disorder. <i>Human Molecular Genetics</i> , 2016 , 25, 3383-3394	5.6	125
128	Variations in FKBP5 and BDNF genes are suggestively associated with depression in a Swedish population-based cohort. <i>Journal of Affective Disorders</i> , 2010 , 125, 249-55	6.6	120
127	Assessment of Response to Lithium Maintenance Treatment in Bipolar Disorder: A Consortium on Lithium Genetics (ConLiGen) Report. <i>PLoS ONE</i> , 2013 , 8, e65636	3.7	113
126	CRY2 is associated with depression. <i>PLoS ONE</i> , 2010 , 5, e9407	3.7	102
125	Antidepressant treatment is associated with epigenetic alterations in the promoter of P11 in a genetic model of depression. <i>International Journal of Neuropsychopharmacology</i> , 2012 , 15, 669-79	5.8	95
124	Kv1.1 channels act as mechanical brake in the senses of touch and pain. <i>Neuron</i> , 2013 , 77, 899-914	13.9	93
123	PER2 variant is associated with depression vulnerability. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2010 , 153B, 570-581	3.5	88
122	Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. <i>Nature Genetics</i> , 2021 , 53, 817-829	36.3	83
121	Association of Polygenic Score for Schizophrenia and HLA Antigen and Inflammation Genes With Response to Lithium in Bipolar Affective Disorder: A Genome-Wide Association Study. <i>JAMA Psychiatry</i> , 2018 , 75, 65-74	14.5	75
120	Multigenic control of disease severity after virulent Mycobacterium tuberculosis infection in mice. <i>Infection and Immunity</i> , 2003 , 71, 126-31	3.7	74
119	Antidepressant-like effect of sodium butyrate is associated with an increase in TET1 and in 5-hydroxymethylation levels in the Bdnf gene. <i>International Journal of Neuropsychopharmacology</i> , 2014 , 18,	5.8	68
118	CLOCK is suggested to associate with comorbid alcohol use and depressive disorders. <i>Journal of Circadian Rhythms</i> , 2010 , 8, 1	2.5	67
117	Severity of tuberculosis in mice is linked to distal chromosome 3 and proximal chromosome 9. <i>Journal of Infectious Diseases</i> , 1999 , 180, 150-5	7	67

116	Large-scale genotyping of single nucleotide polymorphisms by Pyrosequencing trade mark and validation against the 5'Rnuclease (Taqman((R))) assay. <i>Human Mutation</i> , 2002 , 19, 395-401	4.7	63
115	Epigenetic regulation in obesity. <i>International Journal of Obesity</i> , 2012 , 36, 757-65	5.5	59
114	CRY2 is associated with rapid cycling in bipolar disorder patients. <i>PLoS ONE</i> , 2010 , 5, e12632	3.7	58
113	The functional Val158Met polymorphism in catechol-O-methyltransferase (COMT) is associated with depression and motivation in men from a Swedish population-based study. <i>Journal of Affective Disorders</i> , 2011 , 129, 158-66	6.6	57
112	Hypothalamic CART and serum leptin levels are reduced in the anorectic (anx/anx) mouse. <i>Molecular Brain Research</i> , 2000 , 84, 97-105		57
111	Adenovirus-36 is associated with obesity in children and adults in Sweden as determined by rapid ELISA. <i>PLoS ONE</i> , 2012 , 7, e41652	3.7	57
110	The Genetics of the Mood Disorder Spectrum: Genome-wide Association Analyses of More Than 185,000 Cases and 439,000 Controls. <i>Biological Psychiatry</i> , 2020 , 88, 169-184	7.9	57
109	The Risk of Offspring Psychiatric Disorders in the Setting of Maternal Obesity and Diabetes. <i>Pediatrics</i> , 2018 , 142,	7.4	55
108	Telomerase dysregulation in the hippocampus of a rat model of depression: normalization by lithium. <i>International Journal of Neuropsychopharmacology</i> , 2015 , 18, pyv002	5.8	55
107	Examining the public refusal to consent to DNA biobanking: empirical data from a Swedish population-based study. <i>Journal of Medical Ethics</i> , 2010 , 36, 93-8	2.5	54
106	Truncation of the Shaker-like voltage-gated potassium channel, Kv1.1, causes megencephaly. <i>European Journal of Neuroscience</i> , 2003 , 18, 3231-40	3.5	49
105	Associations of Maternal Diabetes and Body Mass Index With Offspring Birth Weight and Prematurity. <i>JAMA Pediatrics</i> , 2019 , 173, 371-378	8.3	48
104	Pyrosequencing-based SNP allele frequency estimation in DNA pools. <i>Human Mutation</i> , 2004 , 23, 92-7	4.7	48
103	Population-based study of antiepileptic drug exposure in utero--influence on head circumference in newborns. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2009 , 18, 672-5	3.2	44
102	Interleukin-6 and depressive symptom severity in response to physical exercise. <i>Psychiatry Research</i> , 2017 , 252, 270-276	9.9	41
101	AHSG gene variant is associated with leanness among Swedish men. <i>Human Genetics</i> , 2005 , 117, 54-60	6.3	41
100	Single nucleotide polymorphism (SNP) allele frequency estimation in DNA pools using Pyrosequencing. <i>Nature Protocols</i> , 2006 , 1, 2573-82	18.8	39
99	NR3C1 hypermethylation in depressed and bullied adolescents. <i>Translational Psychiatry</i> , 2018 , 8, 121	8.6	32

98	CCDC26, CDKN2BAS, RTEL1 and TERT Polymorphisms in pediatric brain tumor susceptibility. <i>Carcinogenesis</i> , 2015 , 36, 876-82	4.6	31
97	School environment and mental health in early adolescence - a longitudinal study in Sweden (KUPOL). <i>BMC Psychiatry</i> , 2016 , 16, 243	4.2	28
96	A multifactorial developmental model for the etiology of major depression in a population-based sample. <i>Journal of Affective Disorders</i> , 2009 , 113, 66-76	6.6	28
95	PreproNPY Pro7 protects against depression despite exposure to environmental risk factors. <i>Journal of Affective Disorders</i> , 2009 , 118, 124-30	6.6	28
94	Genome-wide association study identifies 30 Loci Associated with Bipolar Disorder		28
93	Genetic polymorphisms in monoamine systems and outcome of cognitive behavior therapy for social anxiety disorder. <i>PLoS ONE</i> , 2013 , 8, e79015	3.7	27
92	MRI and in situ hybridization reveal early disturbances in brain size and gene expression in the megencephalic (mceph/mceph) mouse. <i>European Journal of Neuroscience</i> , 2003 , 18, 3218-30	3.5	27
91	P2RX7: expression responds to sleep deprivation and associates with rapid cycling in bipolar disorder type 1. <i>PLoS ONE</i> , 2012 , 7, e43057	3.7	26
90	Enteric short-chain fatty acids promote proliferation of human neural progenitor cells. <i>Journal of Neurochemistry</i> , 2020 , 154, 635-646	6	26
89	A genome-wide association meta-analysis of prognostic outcomes following cognitive behavioural therapy in individuals with anxiety and depressive disorders. <i>Translational Psychiatry</i> , 2019 , 9, 150	8.6	24
88	Early exposure to antibiotic drugs and risk for psychiatric disorders: a population-based study. <i>Translational Psychiatry</i> , 2019 , 9, 317	8.6	24
87	hTERT genetic variation in depression. <i>Journal of Affective Disorders</i> , 2016 , 189, 62-9	6.6	23
86	Mitochondrial DNA copy number is associated with psychosis severity and anti-psychotic treatment. <i>Scientific Reports</i> , 2018 , 8, 12743	4.9	23
85	Carbamazepine protects against megencephaly and abnormal expression of BDNF and Nogo signaling components in the mceph/mceph mouse. <i>Neurobiology of Disease</i> , 2006 , 24, 374-83	7.5	23
84	A truncated Kv1.1 protein in the brain of the megencephaly mouse: expression and interaction. <i>BMC Neuroscience</i> , 2005 , 6, 65	3.2	22
83	Association of polycystic ovary syndrome or anovulatory infertility with offspring psychiatric and mild neurodevelopmental disorders: a Finnish population-based cohort study. <i>Human Reproduction</i> , 2020 , 35, 2336-2347	5.7	22
82	Relationship of prenatal maternal obesity and diabetes to offspring neurodevelopmental and psychiatric disorders: a narrative review. <i>International Journal of Obesity</i> , 2020 , 44, 1981-2000	5.5	21
81	Impact of Childhood Adversity and Vasopressin receptor 1a Variation on Social Interaction in Adulthood: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015 , 10, e0136436	3.7	20

80	TERT rs2736100 genotypes are associated with differential risk of myeloproliferative neoplasms in Swedish and Chinese male patient populations. <i>Annals of Hematology</i> , 2016 , 95, 1825-32	3	20
79	Polymorphism of the AHSB gene is associated with increased adipocyte beta2-adrenoceptor function. <i>Journal of Lipid Research</i> , 2005 , 46, 2278-81	6.3	19
78	MicroRNA 101b Is Downregulated in the Prefrontal Cortex of a Genetic Model of Depression and Targets the Glutamate Transporter SLC1A1 (EAAT3) in Vitro. <i>International Journal of Neuropsychopharmacology</i> , 2016 , 19,	5.8	18
77	Association of polygenic score for major depression with response to lithium in patients with bipolar disorder. <i>Molecular Psychiatry</i> , 2021 , 26, 2457-2470	15.1	17
76	Twelve-week physical exercise does not have a long-lasting effect on kynurenines in plasma of depressed patients. <i>Neuropsychiatric Disease and Treatment</i> , 2017 , 13, 967-972	3.1	17
75	Neuropeptide Y: identification of a novel rat mRNA splice-variant that is downregulated in the hippocampus and the prefrontal cortex of a depression-like model. <i>Peptides</i> , 2012 , 35, 49-55	3.8	17
74	Effects of internet-based cognitive behavioural therapy and physical exercise on sick leave and employment in primary care patients with depression: two subgroup analyses. <i>Occupational and Environmental Medicine</i> , 2018 , 75, 52-58	2.1	16
73	Mood Stabilizers and the Influence on Global Leukocyte DNA Methylation in Bipolar Disorder. <i>Molecular Neuropsychiatry</i> , 2015 , 1, 76-81	4.9	16
72	Lack of potassium channel induces proliferation and survival causing increased neurogenesis and two-fold hippocampus enlargement. <i>Hippocampus</i> , 2007 , 17, 292-304	3.5	16
71	Kv1.1 null mice have enlarged hippocampus and ventral cortex. <i>BMC Neuroscience</i> , 2007 , 8, 10	3.2	16
70	Carbamazepine treatment recovered low N-acetylaspartate+N-acetylaspartylglutamate (tNAA) levels in the megencephaly mouse BALB/cByJ-Kv1.1(mceph/mceph). <i>Neurobiology of Disease</i> , 2007 , 26, 221-8	7.5	16
69	Analysis of the Influence of microRNAs in Lithium Response in Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2018 , 9, 207	5	15
68	Investigating polygenic burden in age at disease onset in bipolar disorder: Findings from an international multicentric study. <i>Bipolar Disorders</i> , 2019 , 21, 68-75	3.8	15
67	Association of brain-derived neurotrophic factor (BDNF) Val66Met polymorphism with early-onset bipolar disorder. <i>Bipolar Disorders</i> , 2015 , 17, 645-52	3.8	15
66	Human adenovirus-36 is uncommon in type 2 diabetes and is associated with increased insulin sensitivity in adults in Sweden. <i>Annals of Medicine</i> , 2014 , 46, 539-46	1.5	15
65	Associations of Different Types of Maternal Diabetes and Body Mass Index With Offspring Psychiatric Disorders. <i>JAMA Network Open</i> , 2020 , 3, e1920787	10.4	14
64	Genome-wide DNA methylomic differences between dorsolateral prefrontal and temporal pole cortices of bipolar disorder. <i>Journal of Psychiatric Research</i> , 2019 , 117, 45-54	5.2	13
63	The megencephaly mouse has disturbances in the insulin-like growth factor (IGF) system. <i>Molecular Brain Research</i> , 1999 , 72, 80-8		13

62	Stress, depressive status and telomere length: Does social interaction and coping strategy play a mediating role?. <i>Journal of Affective Disorders</i> , 2017 , 222, 138-145	6.6	12
61	Prenatal exposure to carbamazepine reduces hippocampal and cortical neuronal cell population in new-born and young mice without detectable effects on learning and memory. <i>PLoS ONE</i> , 2013 , 8, e80497	3.7	11
60	Idiopathic megalencephaly-possible cause and treatment opportunities: from patient to lab. <i>European Journal of Paediatric Neurology</i> , 2008 , 12, 438-45	3.8	10
59	Genetic and Clinical Factors Affecting Plasma Clozapine Concentration. <i>primary care companion for CNS disorders, The</i> , 2015 , 17,	1.2	10
58	Genetic variant in SLC1A2 is associated with elevated anterior cingulate cortex glutamate and lifetime history of rapid cycling. <i>Translational Psychiatry</i> , 2019 , 9, 149	8.6	9
57	Depression-associated ARNTL and PER2 genetic variants in psychotic disorders. <i>Chronobiology International</i> , 2015 , 32, 579-84	3.6	9
56	The serotonin transporter promoter variant (5-HTTLPR) and childhood adversity are associated with the personality trait openness to experience. <i>Psychiatry Research</i> , 2017 , 257, 322-326	9.9	9
55	Improvement in indices of cellular protection after psychological treatment for social anxiety disorder. <i>Translational Psychiatry</i> , 2019 , 9, 340	8.6	9
54	Genome-wide association study of panic disorder reveals genetic overlap with neuroticism and depression. <i>Molecular Psychiatry</i> , 2021 , 26, 4179-4190	15.1	8
53	Common genetic variations in cell cycle and DNA repair pathways associated with pediatric brain tumor susceptibility. <i>Oncotarget</i> , 2016 , 7, 63640-63650	3.3	8
52	Effects of a synbiotic on symptoms, and daily functioning in attention deficit hyperactivity disorder -A double-blind randomized controlled trial. <i>Brain, Behavior, and Immunity</i> , 2020 , 89, 9-19	16.6	7
51	Treating impulsivity with probiotics in adults (PROBIA): study protocol of a multicenter, double-blind, randomized, placebo-controlled trial. <i>Trials</i> , 2020 , 21, 161	2.8	7
50	Expression of telomerase reverse transcriptase positively correlates with duration of lithium treatment in bipolar disorder. <i>Psychiatry Research</i> , 2020 , 286, 112865	9.9	7
49	Cognitive manic symptoms in bipolar disorder associated with polymorphisms in the DAOA and COMT genes. <i>PLoS ONE</i> , 2013 , 8, e67450	3.7	7
48	Carbamazepine protects against neuronal hyperplasia and abnormal gene expression in the megalencephaly mouse. <i>Neurobiology of Disease</i> , 2008 , 32, 364-76	7.5	7
47	Childhood adversity increases methylation in the GRIN2B gene. <i>Journal of Psychiatric Research</i> , 2021 , 132, 38-43	5.2	7
46	KIBRA genetic polymorphism and cognitive dysfunction in depression. <i>Psychiatry Research</i> , 2015 , 226, 405-6	9.9	6
45	Lithium and the Interplay Between Telomeres and Mitochondria in Bipolar Disorder. <i>Frontiers in Psychiatry</i> , 2020 , 11, 586083	5	6

44	Neuropeptide Y, stressful life events and personality trait conscientiousness: Preliminary associations from a Swedish longitudinal study. <i>Psychiatry Research</i> , 2018 , 263, 48-53	9.9	6
43	Evidence for presence and functional effects of Kv1.1 channels in Ectels: general survey and results from mceph/mceph mice. <i>PLoS ONE</i> , 2011 , 6, e18213	3.7	6
42	The Effect of Smartphone Apps Versus Supervised Exercise on Physical Activity, Cardiorespiratory Fitness, and Body Composition Among Individuals With Mild-to-Moderate Mobility Disability: Randomized Controlled Trial. <i>JMIR MHealth and UHealth</i> , 2020 , 8, e14615	5.5	6
41	Association of maternal polycystic ovary syndrome or anovulatory infertility with obesity and diabetes in offspring: a population-based cohort study. <i>Human Reproduction</i> , 2021 , 36, 2345-2357	5.7	6
40	BDNF Val66Met and childhood adversity on response to physical exercise and internet-based cognitive behavioural therapy in depressed Swedish adults. <i>Journal of Psychiatric Research</i> , 2017 , 93, 50-58	5.2	5
39	Path analysis of the chronicity of depression using the comprehensive developmental model framework. <i>Nordic Journal of Psychiatry</i> , 2016 , 70, 380-91	2.3	5
38	5-HTTLPR, victimization and ecological executive function of adolescents. <i>Psychiatry Research</i> , 2016 , 237, 55-9	9.9	5
37	Plasma GDF15 level is elevated in psychosis and inversely correlated with severity. <i>Scientific Reports</i> , 2017 , 7, 7906	4.9	5
36	Proinflammatory mediators and their associations with medication and comorbid traits in children and adults with ADHD. <i>European Neuropsychopharmacology</i> , 2020 , 41, 118-131	1.2	5
35	Association of Catechol-O-methyltransferase (COMT ValMet) with future risk of cardiovascular disease in depressed individuals - a Swedish population-based cohort study. <i>BMC Medical Genetics</i> , 2018 , 19, 126	2.1	4
34	Influence of serotonin transporter promoter variation on the effects of separation from parent/partner on depression. <i>Journal of Affective Disorders</i> , 2013 , 144, 216-24	6.6	4
33	Acoustic startle hypersensitivity in Mceph mice and its effect on hippocampal excitability. <i>European Journal of Neuroscience</i> , 2011 , 34, 1121-30	3.5	4
32	Pulse Pressure Magnifies the Effect of COMT Val(158)Met on 15 Years Episodic Memory Trajectories. <i>Frontiers in Aging Neuroscience</i> , 2016 , 8, 34	5.3	4
31	Sex-specific effects of gain-of-function P2RX7 variation on bipolar disorder. <i>Journal of Affective Disorders</i> , 2019 , 245, 597-601	6.6	4
30	Working conditions, serotonin transporter gene polymorphism (5-HTTLPR) and anxiety disorders: a prospective cohort study. <i>Journal of Affective Disorders</i> , 2013 , 151, 652-659	6.6	3
29	Genetic variants of increased waist circumference in psychosis. <i>Psychiatric Genetics</i> , 2017 , 27, 210-218	2.9	3
28	Evaluation of serological assays for diagnosis of onchocercosis. <i>Scandinavian Journal of Infectious Diseases</i> , 1997 , 29, 65-70		3
27	A Weighted Genetic Risk Score of Adult Glioma Susceptibility Loci Associated with Pediatric Brain Tumor Risk. <i>Scientific Reports</i> , 2019 , 9, 18142	4.9	3

26	Cortisol Concentration as Predictor of Tobacco Initiation in Adolescents: Results From a Population-Based Swedish Cohort. <i>Journal of Adolescent Health</i> , 2021 , 68, 758-764	5.8	3
25	Diagnosis of onchocerciasis using highly specific and sensitive native proteins. <i>Scandinavian Journal of Infectious Diseases</i> , 2002 , 34, 583-90		2
24	Reply: Association of maternal polycystic ovary syndrome or anovulatory infertility with obesity and diabetes in offspring: a population-based cohort study. <i>Human Reproduction</i> , 2021 ,	5.7	2
23	Melatonin receptor 1B gene associated with hyperglycemia in bipolar disorder. <i>Psychiatric Genetics</i> , 2016 , 26, 136-9	2.9	2
22	Exercise Reduces Salivary Morning Cortisol Levels in Patients with Depression. <i>Molecular Neuropsychiatry</i> , 2019 , 4, 196-203	4.9	2
21	Parental age and risk of genetic syndromes predisposing to nervous system tumors: nested case-control study. <i>Clinical Epidemiology</i> , 2018 , 10, 729-738	5.9	2
20	Troponin T levels associated with genetic variants in NOTCH2 and MTNR1B in women with psychosis. <i>Psychiatry Research</i> , 2017 , 250, 217-220	9.9	1
19	Implications of Gestational Weight Gain in Studies of Gestational Diabetes-Reply. <i>JAMA Pediatrics</i> , 2019 , 173, 889-890	8.3	1
18	FitForLife: study protocol for a randomized controlled trial. <i>Trials</i> , 2015 , 16, 553	2.8	1
17	Field diagnosis of onchocerciasis in an area of high versus low endemicity: evaluation of the Dot Blot Assay. <i>Scandinavian Journal of Infectious Diseases</i> , 1996 , 28, 75-81		1
16	Combining schizophrenia and depression polygenic risk scores improves the genetic prediction of lithium response in bipolar disorder patients. <i>Translational Psychiatry</i> , 2021 , 11, 606	8.6	1
15	Polygenic scores for major depressive disorder and depressive symptoms predict response to lithium in patients with bipolar disorder		1
14	AKT1 and genetic vulnerability to bipolar disorder. <i>Psychiatry Research</i> , 2020 , 284, 112677	9.9	1
13	Single-nucleotide polymorphism in the human TIA1 gene interacts with stressful life events to predict the development of pathological anxiety symptoms in a Swedish population. <i>Journal of Affective Disorders</i> , 2020 , 260, 597-603	6.6	1
12	Prediction of lithium response using genomic data. <i>Scientific Reports</i> , 2021 , 11, 1155	4.9	1
11	No association of cigarette smoking and depressive symptoms with cortisol concentration in adolescents. Results from a population-based Swedish cohort. <i>Psychiatry Research</i> , 2021 , 301, 113968	9.9	1
10	DNA methylation of the glucocorticoid receptor gene predicts substance use in adolescence: longitudinal data from over 1000 young individuals. <i>Translational Psychiatry</i> , 2021 , 11, 477	8.6	1
9	HLA-DRB1 and HLA-DQB1 genetic diversity modulates response to lithium in bipolar affective disorders. <i>Scientific Reports</i> , 2021 , 11, 17823	4.9	1

8	Using polygenic scores and clinical data for bipolar disorder patient stratification and lithium response prediction: machine learning approach.. <i>British Journal of Psychiatry</i> , 2022 , 1-10	5.4	1
7	Association of Preeclampsia and Perinatal Complications With Offspring Neurodevelopmental and Psychiatric Disorders.. <i>JAMA Network Open</i> , 2022 , 5, e2145719	10.4	0
6	The cannabinoid receptor-1 gene interacts with stressful life events to increase the risk for problematic alcohol use.. <i>Scientific Reports</i> , 2022 , 12, 4963	4.9	0
5	Improving lithium dose prediction using population pharmacokinetics and pharmacogenomics: a cohort genome-wide association study in Sweden.. <i>Lancet Psychiatry</i> , 2022 , 9, 447-457	23.3	0
4	Genetic variations of NPY and AGRP in body fatness regulation. <i>Future Lipidology</i> , 2007 , 2, 147-151		
3	Physical exercise is associated with a reduction in plasma levels of fractalkine, TGF- β , eotaxin-1 and IL-6 in younger adults with mobility disability.. <i>PLoS ONE</i> , 2022 , 17, e0263173	3.7	
2	and downregulation in Flinders Sensitive Line Rat Model of Depression. <i>Acta Neuropsychiatrica</i> , 2021 , 1-19	3.9	
1	Physical exercise is associated with a reduction in inflammatory biomarkers in first-episode psychosis: A pilot study of CRP, SAA, sICAM-1 and sVCAM-1. <i>Schizophrenia Research</i> , 2021 , 228, 316-318	3.6	