

Inge S Fomsgaard

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9569074/publications.pdf>

Version: 2024-02-01

103
papers

3,054
citations

126708

33
h-index

189595

50
g-index

105
all docs

105
docs citations

105
times ranked

3288
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Data-dependent acquisition-mass spectrometry guided isolation of new benzoxazinoids from the roots of <i>Acanthus mollis</i> L. <i>International Journal of Mass Spectrometry</i> , 2022, 474, 116815. | 0.7 | 4 |
| 2 | An inverse association between plasma benzoxazinoid metabolites and PSA after rye intake in men with prostate cancer revealed with a new method. <i>Scientific Reports</i> , 2022, 12, 5260. | 1.6 | 3 |
| 3 | Integrated LC-MS and GC-MS-Based Metabolomics Reveal the Effects of Plant Competition on the Rye Metabolome. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 3056-3066. | 2.4 | 13 |
| 4 | Optimised extraction and LC-MS/MS analysis of flavonoids reveal large field variation in exudation into <i>Lupinus Angustifolius</i> L. rhizosphere soil. <i>Rhizosphere</i> , 2022, 22, 100516. | 1.4 | 5 |
| 5 | <i>Fusarium oxysporum</i> Disrupts Microbiome-Metabolome Networks in <i>Arabidopsis thaliana</i> Roots. <i>Microbiology Spectrum</i> , 2022, 10, . | 1.2 | 8 |
| 6 | Dietary quercetin impacts the concentration of pesticides in honey bees. <i>Chemosphere</i> , 2021, 262, 127848. | 4.2 | 24 |
| 7 | Stepwise mass spectrometry-based approach for confirming the presence of benzoxazinoids in herbs and vegetables. <i>Phytochemical Analysis</i> , 2021, 32, 283-297. | 1.2 | 4 |
| 8 | Seed inoculations with entomopathogenic fungi affect aphid populations coinciding with modulation of plant secondary metabolite profiles across plant families. <i>New Phytologist</i> , 2021, 229, 1715-1727. | 3.5 | 38 |
| 9 | Targeted metabolomics unveil alteration in accumulation and root exudation of flavonoids as a response to interspecific competition. <i>Journal of Plant Interactions</i> , 2021, 16, 53-63. | 1.0 | 14 |
| 10 | LC-MS/MS Quantification Reveals Ample Gut Uptake and Metabolization of Dietary Phytochemicals in Honey Bees (<i>Apis mellifera</i>). <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 627-637. | 2.4 | 7 |
| 11 | Benzoxazinoids selectively affect maize root-associated nematode taxa. <i>Journal of Experimental Botany</i> , 2021, 72, 3835-3845. | 2.4 | 15 |
| 12 | Benzoxazinoids Are Inversely Associated With Prostate-Specific Antigen Levels- a Whole Grain Rye vs Refined Wheat Randomized Cross-Over Trial in Men With Prostate Cancer. <i>Current Developments in Nutrition</i> , 2021, 5, 482. | 0.1 | 0 |
| 13 | Metabolomics unveils the influence of dietary phytochemicals on residual pesticide concentrations in honey bees. <i>Environment International</i> , 2021, 152, 106503. | 4.8 | 32 |
| 14 | Metabolic profiling of benzoxazinoids in the roots and rhizosphere of commercial winter wheat genotypes. <i>Plant and Soil</i> , 2021, 466, 467-489. | 1.8 | 15 |
| 15 | Barley Nepenthesin-Like Aspartic Protease HvNEP-1 Degrades <i>Fusarium</i> Phytase, Impairs Toxin Production, and Suppresses the Fungal Growth. <i>Frontiers in Plant Science</i> , 2021, 12, 702557. | 1.7 | 0 |
| 16 | Determination of the Effect of Co-cultivation on the Production and Root Exudation of Flavonoids in Four Legume Species Using LC-MS/MS Analysis. <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 9208-9219. | 2.4 | 17 |
| 17 | Mass spectrometry-based metabolomics unravel the transfer of bioactive compounds between rye and neighbouring plants. <i>Plant, Cell and Environment</i> , 2021, 44, 3492-3501. | 2.8 | 5 |
| 18 | Analytical Methods for Quantification and Identification of Intact Glucosinolates in <i>Arabidopsis</i> Roots Using LC-QQ(LIT)-MS/MS. <i>Metabolites</i> , 2021, 11, 47. | 1.3 | 11 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Root-Exuded Benzoxazinoids: Uptake and Translocation in Neighboring Plants. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 10609-10617. | 2.4 | 25 |
| 20 | Mass Spectrometry-Based Metabolomics Reveals a Concurrent Action of Several Chemical Mechanisms in <i>Arabidopsis-Fusarium oxysporum</i> Compatible and Incompatible Interactions. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 15335-15344. | 2.4 | 6 |
| 21 | Overexpression of Nepenthesin HvNEP-1 in Barley Endosperm Reduces Fusarium Head Blight and Mycotoxin Accumulation. <i>Agronomy</i> , 2020, 10, 203. | 1.3 | 11 |
| 22 | Identification of Azoxystrobin Glutathione Conjugate Metabolites in Maize Roots by LC-MS. <i>Molecules</i> , 2019, 24, 2473. | 1.7 | 5 |
| 23 | Maize synthesized benzoxazinoids affect the host associated microbiome. <i>Microbiome</i> , 2019, 7, 59. | 4.9 | 185 |
| 24 | Influence of the growing conditions on the flavonoids and phenolic acids accumulation in amaranth (<i>Amaranthus hypochondriacus</i> L.) leaves.. <i>Terra Latinoamericana</i> , 2019, 37, 449. | 0.3 | 6 |
| 25 | Weed suppressive traits of winter cereals: Allelopathy and competition. <i>Biochemical Systematics and Ecology</i> , 2018, 76, 35-41. | 0.6 | 23 |
| 26 | Maize root culture as a model system for studying azoxystrobin biotransformation in plants. <i>Chemosphere</i> , 2018, 195, 624-631. | 4.2 | 7 |
| 27 | Effect of Tillage Systems on the Dissipation of Prosulfocarb Herbicide. <i>Weed Technology</i> , 2018, 32, 195-204. | 0.4 | 2 |
| 28 | Weed suppression by Canadian spring cereals: relative contribution of competition for resources and allelopathy. <i>Chemoecology</i> , 2018, 28, 183-187. | 0.6 | 5 |
| 29 | Sorption and degradation of neonicotinoid insecticides in tropical soils. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2018, 53, 587-594. | 0.7 | 37 |
| 30 | Multiple effects of secondary metabolites on amino acid cycling in white clover rhizosphere. <i>Soil Biology and Biochemistry</i> , 2018, 123, 54-63. | 4.2 | 30 |
| 31 | Weed suppression by winter cereals: relative contribution of competition for resources and allelopathy. <i>Chemoecology</i> , 2018, 28, 109-121. | 0.6 | 18 |
| 32 | Quantitative analysis of absorption, metabolism, and excretion of benzoxazinoids in humans after the consumption of high- and low-benzoxazinoid diets with similar contents of cereal dietary fibres: a crossover study. <i>European Journal of Nutrition</i> , 2017, 56, 387-397. | 4.6 | 14 |
| 33 | Biosynthesis and chemical transformation of benzoxazinoids in rye during seed germination and the identification of a rye Bx6-like gene. <i>Phytochemistry</i> , 2017, 140, 95-107. | 1.4 | 33 |
| 34 | Alterations of the Benzoxazinoid Profiles of Uninjured Maize Seedlings During Freezing, Storage, and Lyophilization. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 4103-4110. | 2.4 | 8 |
| 35 | Quantification of azoxystrobin and identification of two novel metabolites in lettuce via liquid chromatography-quadrupole-linear ion trap (QTRAP) mass spectrometry. <i>International Journal of Environmental Analytical Chemistry</i> , 2017, 97, 419-430. | 1.8 | 12 |
| 36 | Liquid chromatography-tandem mass spectrometry method for simultaneous quantification of azoxystrobin and its metabolites, azoxystrobin free acid and 2-hydroxybenzoxazinone, in greenhouse-grown lettuce. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2017, 34, 2173-2180. | 1.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Biphenyl Columns Provide Good Separation of the Glucosides of DIMBOA and DIM2BOA. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200. | 0.2 | 6 |
| 38 | Bioactive small molecules in commercially available cereal food: Benzoxazinoids. <i>Journal of Food Composition and Analysis</i> , 2017, 64, 213-222. | 1.9 | 9 |
| 39 | Profiling and Metabolism of Sterols in the Weaver Ant Genus <i>Oecophylla</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100. | 0.2 | 0 |
| 40 | Dissipation kinetics of asparagine in soil measured by compound-specific analysis with metabolite tracking. <i>Biology and Fertility of Soils</i> , 2016, 52, 911-916. | 2.3 | 5 |
| 41 | Identification and Quantification of Loline-Type Alkaloids in Endophyte-Infected Grasses by LC-MS/MS. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 6212-6218. | 2.4 | 15 |
| 42 | Benzoxazinoids in Prostate Cancer Patients after a Rye-Intensive Diet: Methods and Initial Results. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 8235-8245. | 2.4 | 16 |
| 43 | Direct acquisition of organic N by white clover even in the presence of inorganic N. <i>Plant and Soil</i> , 2016, 407, 91-107. | 1.8 | 31 |
| 44 | Correlation of Deoxynivalenol Accumulation in <i>Fusarium</i> -Infected Winter and Spring Wheat Cultivars with Secondary Metabolites at Different Growth Stages. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 4545-4555. | 2.4 | 21 |
| 45 | Urea in Weaver Ant Feces: Quantification and Investigation of the Uptake and Translocation of Urea in <i>Coffea arabica</i> . <i>Journal of Plant Growth Regulation</i> , 2016, 35, 803-814. | 2.8 | 8 |
| 46 | Dietary exposure to benzoxazinoids enhances bacteria-induced monokine responses by peripheral blood mononuclear cells. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 2190-2198. | 1.5 | 2 |
| 47 | Metabolic Profiling of <i>Arabidopsis thaliana</i> Reveals Herbicide- and Allelochemical-Dependent Alterations Before They Become Apparent in Plant Growth. <i>Journal of Plant Growth Regulation</i> , 2015, 34, 96-107. | 2.8 | 5 |
| 48 | Benzoxazinoids: Cereal phytochemicals with putative therapeutic and health-protecting properties. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1324-1338. | 1.5 | 71 |
| 49 | Are ant feces nutrients for plants? A metabolomics approach to elucidate the nutritional effects on plants hosting weaver ants. <i>Metabolomics</i> , 2015, 11, 1013-1028. | 1.4 | 10 |
| 50 | 2,4-Dihydroxy-7-methoxy-2 H -1,4-benzoxazin-3(4 H)-one (DIMBOA) inhibits trichothecene production by <i>Fusarium graminearum</i> through suppression of Tri6 expression. <i>International Journal of Food Microbiology</i> , 2015, 214, 123-128. | 2.1 | 34 |
| 51 | Application of the QuEChERS procedure and LC-MS/MS for the assessment of neonicotinoid insecticide residues in cocoa beans and shells. <i>Journal of Food Composition and Analysis</i> , 2015, 44, 149-157. | 1.9 | 43 |
| 52 | Threshold response of stomatal closing ability to leaf abscisic acid concentration during growth. <i>Journal of Experimental Botany</i> , 2014, 65, 4361-4370. | 2.4 | 61 |
| 53 | Quantification of neonicotinoid insecticide residues in soils from cocoa plantations using a QuEChERS extraction procedure and LC-MS/MS. <i>Science of the Total Environment</i> , 2014, 499, 276-283. | 3.9 | 83 |
| 54 | Foliar abscisic acid content underlies genotypic variation in stomatal responsiveness after growth at high relative air humidity. <i>Annals of Botany</i> , 2013, 112, 1857-1867. | 1.4 | 45 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Nutritional composition of minor indigenous fruits: Cheapest nutritional source for the rural people of Bangladesh. <i>Food Chemistry</i> , 2013, 140, 466-470. | 4.2 | 42 |
| 56 | The Response of <i>Arabidopsis</i> to Co-cultivation with Clover. <i>ACS Symposium Series</i> , 2013, , 189-201. | 0.5 | 4 |
| 57 | Phenolic Acids from Wheat Show Different Absorption Profiles in Plasma: A Model Experiment with Catheterized Pigs. <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 8842-8850. | 2.4 | 12 |
| 58 | Comparison of the levels of bioactive benzoxazinoids in different wheat and rye fractions and the transformation of these compounds in homemade foods. <i>Food Chemistry</i> , 2013, 141, 444-450. | 4.2 | 51 |
| 59 | Differences among five amaranth varieties (<i>Amaranthus</i> spp.) regarding secondary metabolites and foliar herbivory by chewing insects in the field. <i>Arthropod-Plant Interactions</i> , 2013, 7, 235-245. | 0.5 | 26 |
| 60 | Absorption and metabolic fate of bioactive dietary benzoxazinoids in humans. <i>Molecular Nutrition and Food Research</i> , 2013, 57, 1847-1858. | 1.5 | 37 |
| 61 | Plasma and Urine Concentrations of Bioactive Dietary Benzoxazinoids and Their Glucuronidated Conjugates in Rats Fed a Rye Bread-Based Diet. <i>Journal of Agricultural and Food Chemistry</i> , 2012, 60, 11518-11524. | 2.4 | 24 |
| 62 | Bioactive Benzoxazinoids in Rye Bread Are Absorbed and Metabolized in Pigs. <i>Journal of Agricultural and Food Chemistry</i> , 2012, 60, 2497-2506. | 2.4 | 33 |
| 63 | Phytotoxic Effect, Uptake, and Transformation of Biochanin A in Selected Weed Species. <i>Journal of Agricultural and Food Chemistry</i> , 2012, 60, 10715-10722. | 2.4 | 19 |
| 64 | Fate in Soil of Flavonoids Released from White Clover (<i>Trifolium repens</i> L.). <i>Applied and Environmental Soil Science</i> , 2012, 2012, 1-10. | 0.8 | 39 |
| 65 | Variation of Polyphenols and Betaines in Aerial Parts of Young, Field-Grown <i>Amaranthus</i> Genotypes. <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 12073-12082. | 2.4 | 29 |
| 66 | Degradation of biochanin A in soil. <i>Chemoecology</i> , 2011, 21, 59-66. | 0.6 | 6 |
| 67 | Bread from common cereal cultivars contains an important array of neglected bioactive benzoxazinoids. <i>Food Chemistry</i> , 2011, 127, 1814-1820. | 4.2 | 65 |
| 68 | Variations in the polyphenol content of seeds of field grown <i>Amaranthus</i> genotypes. <i>Food Chemistry</i> , 2011, 129, 131-138. | 4.2 | 57 |
| 69 | Proximate composition, phenolic acids, and flavonoids characterization of commercial and wild nopal (<i>Opuntia</i> spp.). <i>Journal of Food Composition and Analysis</i> , 2010, 23, 525-532. | 1.9 | 121 |
| 70 | Phenolic Compounds in Different Barley Varieties: Identification by Tandem Mass Spectrometry (QStar) and NMR; Quantification by Liquid Chromatography Triple Quadrupole-Linear Ion Trap Mass Spectrometry (Q-Trap). <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500. | 0.2 | 15 |
| 71 | Synthesis and Quantitation of Six Phenolic Amides in <i>Amaranthus</i> spp.. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 6306-6311. | 2.4 | 32 |
| 72 | Allelochemicals in Rye (<i>Secale Cereale</i> L.): Cultivar and Tissue Differences in the Production of Benzoxazinoids and Phenolic Acids. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900400. | 0.2 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Amaranth (<i>Amaranthus hypochondriacus</i>) as an alternative crop for sustainable food production: Phenolic acids and flavonoids with potential impact on its nutraceutical quality. <i>Journal of Cereal Science</i> , 2009, 49, 117-121. | 1.8 | 144 |
| 74 | Allelochemicals in rye (<i>Secale cereale</i> L.): cultivar and tissue differences in the production of benzoxazinoids and phenolic acids. <i>Natural Product Communications</i> , 2009, 4, 199-208. | 0.2 | 32 |
| 75 | Flavonoids in roots of white clover: interaction of arbuscular mycorrhizal fungi and a pathogenic fungus. <i>Plant and Soil</i> , 2008, 302, 33-43. | 1.8 | 72 |
| 76 | Biologically active secondary metabolites in white clover (<i>Trifolium repens</i> L.) – a review focusing on contents in the plant, plant–pest interactions and transformation. <i>Chemoecology</i> , 2008, 18, 129-170. | 0.6 | 53 |
| 77 | Benzoxazinoid concentrations show correlation with Fusarium Head Blight resistance in Danish wheat varieties. <i>Biochemical Systematics and Ecology</i> , 2008, 36, 245-259. | 0.6 | 47 |
| 78 | Fate and availability of glyphosate and AMPA in agricultural soil. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2008, 43, 365-375. | 0.7 | 104 |
| 79 | Transformation kinetics of 6-methoxybenzoxazolin-2-one in soil. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2008, 43, 1-7. | 0.7 | 38 |
| 80 | Variation in Flavonoids in Leaves, Stems and Flowers of White Clover Cultivars. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300. | 0.2 | 6 |
| 81 | Rebuttal on Results from the FATEALLCHEM Project. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 1645-1647. | 2.4 | 0 |
| 82 | Chemical Ecology in Wheat Plant–Pest Interactions. How the Use of Modern Techniques and a Multidisciplinary Approach Can Throw New Light on a Well-known Phenomenon: Allelopathy. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 987-990. | 2.4 | 42 |
| 83 | Fate of Benzoxazinone Allelochemicals in Soil after Incorporation of Wheat and Rye Sprouts. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1064-1074. | 2.4 | 78 |
| 84 | Transformation of Benzoxazinones and Derivatives and Microbial Activity in the Test Environment of Soil Ecotoxicological Tests on <i>Poecilus cupreus</i> and <i>Folsomia candida</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1086-1092. | 2.4 | 43 |
| 85 | Elucidating the Transformation Pattern of the Cereal Allelochemical 6-Methoxy-2-benzoxazolinone (MBOA) and the Trideuteriomethoxy Analogue [D3]-MBOA in Soil. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1075-1085. | 2.4 | 32 |
| 86 | Biotransformation of 2-Benzoxazolinone to 2-Amino-(3H)-Phenoxazin-3-one and 2-Acetylamino-(3H)-Phenoxazin-3-one in Soil. <i>Journal of Chemical Ecology</i> , 2005, 31, 1205-1222. | 0.9 | 52 |
| 87 | Transformation products of 2-benzoxazolinone (BOA) in soil. <i>Chemosphere</i> , 2005, 61, 74-84. | 4.2 | 38 |
| 88 | First European interlaboratory study of the analysis of benzoxazinone derivatives in plants by liquid chromatography. <i>Journal of Chromatography A</i> , 2004, 1047, 69-76. | 1.8 | 33 |
| 89 | Microbial transformation products of benzoxazolinone and benzoxazinone allelochemicals – a review. <i>Chemosphere</i> , 2004, 54, 1025-1038. | 4.2 | 106 |
| 90 | Leaching of Pesticides Through Normal–Tillage and Low–Tillage Soil – A Lysimeter Study. I. Isoproturon. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2003, 38, 1-18. | 0.7 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Leaching of Pesticides Through Normal Tillage and Low Tillage Soil – A Lysimeter Study. II. Glyphosate. Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes, 2003, 38, 19-35. | 0.7 | 40 |
| 92 | Comparison and evaluation of eight pesticide environmental risk indicators developed in Europe and recommendations for future use. Agriculture, Ecosystems and Environment, 2002, 90, 177-187. | 2.5 | 161 |
| 93 | Influence of microbial activity, organic carbon content, soil texture and soil depth on mineralisation rates of low concentrations of ¹⁴ C-mecoprop – development of a predictive model. Ecological Modelling, 1999, 122, 45-68. | 1.2 | 20 |
| 94 | Sampling and Substrate Application Methods for Pesticide Mineralization Experiments in Undisturbed Soil Samples. International Journal of Environmental Analytical Chemistry, 1998, 70, 121-132. | 1.8 | 3 |
| 95 | Degradation of Mecoprop and Isoproturon in Soil Influence of Initial Concentration. International Journal of Environmental Analytical Chemistry, 1998, 70, 133-148. | 1.8 | 23 |
| 96 | Degradation of ¹⁴ C-maneb in sediment from a Nicaraguan estuary. International Journal of Environmental Studies, 1998, 55, 175-198. | 0.7 | 2 |
| 97 | Modelling the mineralization kinetics for low concentrations of pesticides in surface and subsurface soil. Ecological Modelling, 1997, 102, 175-208. | 1.2 | 44 |
| 98 | Degradation of [¹⁴ C]ethylenethiourea in surface and subsurface soil. Science of the Total Environment, 1996, 191, 271-276. | 3.9 | 6 |
| 99 | Degradation of Pesticides in Subsurface Soils, Unsaturated Zone – a Review Of Methods and Results. International Journal of Environmental Analytical Chemistry, 1995, 58, 231-245. | 1.8 | 53 |
| 100 | Lead, arsenic, cadmium and copper in Lake Asososca, Nicaragua. Science of the Total Environment, 1994, 155, 229-236. | 3.9 | 5 |
| 101 | Toxaphene and Other Organochlorine Pesticides in Fish and Sediment from Lake Xolotlán, Nicaragua. International Journal of Environmental Analytical Chemistry, 1993, 53, 297-305. | 1.8 | 14 |
| 102 | Preliminary study of 15 organochlorine pesticides in Lake Xolotlan, Nicaragua. Chemosphere, 1992, 24, 1413-1419. | 4.2 | 7 |
| 103 | Mercury contamination in Lake Xolotlán (Managua). Hydrobiological Bulletin, 1991, 25, 173-176. | 0.5 | 5 |