

# Inge S Fomsgaard

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9569074/publications.pdf>

Version: 2024-02-01

103  
papers

3,054  
citations

126907

33  
h-index

189892

50  
g-index

105  
all docs

105  
docs citations

105  
times ranked

3288  
citing authors

#	ARTICLE	IF	CITATIONS
1	Data-dependent acquisition-mass spectrometry guided isolation of new benzoxazinoids from the roots of <i>Acanthus mollis</i> L. <i>International Journal of Mass Spectrometry</i> , 2022, 474, 116815.	1.5	4
2	An inverse association between plasma benzoxazinoid metabolites and PSA after rye intake in men with prostate cancer revealed with a new method. <i>Scientific Reports</i> , 2022, 12, 5260.	3.3	3
3	Integrated LC-MS and GC-MS-Based Metabolomics Reveal the Effects of Plant Competition on the Rye Metabolome. <i>Journal of Agricultural and Food Chemistry</i> , 2022, 70, 3056-3066.	5.2	13
4	Optimised extraction and LC-MS/MS analysis of flavonoids reveal large field variation in exudation into <i>Lupinus Angustifolius</i> L. rhizosphere soil. <i>Rhizosphere</i> , 2022, 22, 100516.	3.0	5
5	<i>Fusarium oxysporum</i> Disrupts Microbiome-Metabolome Networks in <i>Arabidopsis thaliana</i> Roots. <i>Microbiology Spectrum</i> , 2022, 10, .	3.0	8
6	Dietary quercetin impacts the concentration of pesticides in honey bees. <i>Chemosphere</i> , 2021, 262, 127848.	8.2	24
7	Stepwise mass spectrometry-based approach for confirming the presence of benzoxazinoids in herbs and vegetables. <i>Phytochemical Analysis</i> , 2021, 32, 283-297.	2.4	4
8	Seed inoculations with entomopathogenic fungi affect aphid populations coinciding with modulation of plant secondary metabolite profiles across plant families. <i>New Phytologist</i> , 2021, 229, 1715-1727.	7.3	38
9	Targeted metabolomics unveil alteration in accumulation and root exudation of flavonoids as a response to interspecific competition. <i>Journal of Plant Interactions</i> , 2021, 16, 53-63.	2.1	14
10	LC-MS/MS Quantification Reveals Ample Gut Uptake and Metabolization of Dietary Phytochemicals in Honey Bees ( <i>Apis mellifera</i> ). <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 627-637.	5.2	7
11	Benzoxazinoids selectively affect maize root-associated nematode taxa. <i>Journal of Experimental Botany</i> , 2021, 72, 3835-3845.	4.8	15
12	Benzoxazinoids Are Inversely Associated With Prostate-Specific Antigen Levels- a Whole Grain Rye vs Refined Wheat Randomized Cross-Over Trial in Men With Prostate Cancer. <i>Current Developments in Nutrition</i> , 2021, 5, 482.	0.3	0
13	Metabolomics unveils the influence of dietary phytochemicals on residual pesticide concentrations in honey bees. <i>Environment International</i> , 2021, 152, 106503.	10.0	32
14	Metabolic profiling of benzoxazinoids in the roots and rhizosphere of commercial winter wheat genotypes. <i>Plant and Soil</i> , 2021, 466, 467-489.	3.7	15
15	Barley Nepenthesin-Like Aspartic Protease HvNEP-1 Degrades <i>Fusarium</i> Phytase, Impairs Toxin Production, and Suppresses the Fungal Growth. <i>Frontiers in Plant Science</i> , 2021, 12, 702557.	3.6	0
16	Determination of the Effect of Co-cultivation on the Production and Root Exudation of Flavonoids in Four Legume Species Using LC-MS/MS Analysis. <i>Journal of Agricultural and Food Chemistry</i> , 2021, 69, 9208-9219.	5.2	17
17	Mass spectrometry-based metabolomics unravel the transfer of bioactive compounds between rye and neighbouring plants. <i>Plant, Cell and Environment</i> , 2021, 44, 3492-3501.	5.7	5
18	Analytical Methods for Quantification and Identification of Intact Glucosinolates in <i>Arabidopsis</i> Roots Using LC-QqQ(LIT)-MS/MS. <i>Metabolites</i> , 2021, 11, 47.	2.9	11

#	ARTICLE	IF	CITATIONS
19	Root-Exuded Benzoxazinoids: Uptake and Translocation in Neighboring Plants. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 10609-10617.	5.2	25
20	Mass Spectrometry-Based Metabolomics Reveals a Concurrent Action of Several Chemical Mechanisms in <i>Arabidopsis-Fusarium oxysporum</i> Compatible and Incompatible Interactions. <i>Journal of Agricultural and Food Chemistry</i> , 2020, 68, 15335-15344.	5.2	6
21	Overexpression of Nepenthesin HvNEP-1 in Barley Endosperm Reduces Fusarium Head Blight and Mycotoxin Accumulation. <i>Agronomy</i> , 2020, 10, 203.	3.0	11
22	Identification of Azoxystrobin Glutathione Conjugate Metabolites in Maize Roots by LC-MS. <i>Molecules</i> , 2019, 24, 2473.	3.8	5
23	Maize synthesized benzoxazinoids affect the host associated microbiome. <i>Microbiome</i> , 2019, 7, 59.	11.1	185
24	Influence of the growing conditions on the flavonoids and phenolic acids accumulation in amaranth ( <i>Amaranthus hypochondriacus</i> L.) leaves.. <i>Terra Latinoamericana</i> , 2019, 37, 449.	0.3	6
25	Weed suppressive traits of winter cereals: Allelopathy and competition. <i>Biochemical Systematics and Ecology</i> , 2018, 76, 35-41.	1.3	23
26	Maize root culture as a model system for studying azoxystrobin biotransformation in plants. <i>Chemosphere</i> , 2018, 195, 624-631.	8.2	7
27	Effect of Tillage Systems on the Dissipation of Prosulfocarb Herbicide. <i>Weed Technology</i> , 2018, 32, 195-204.	0.9	2
28	Weed suppression by Canadian spring cereals: relative contribution of competition for resources and allelopathy. <i>Chemoecology</i> , 2018, 28, 183-187.	1.1	5
29	Sorption and degradation of neonicotinoid insecticides in tropical soils. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2018, 53, 587-594.	1.5	37
30	Multiple effects of secondary metabolites on amino acid cycling in white clover rhizosphere. <i>Soil Biology and Biochemistry</i> , 2018, 123, 54-63.	8.8	30
31	Weed suppression by winter cereals: relative contribution of competition for resources and allelopathy. <i>Chemoecology</i> , 2018, 28, 109-121.	1.1	18
32	Quantitative analysis of absorption, metabolism, and excretion of benzoxazinoids in humans after the consumption of high- and low-benzoxazinoid diets with similar contents of cereal dietary fibres: a crossover study. <i>European Journal of Nutrition</i> , 2017, 56, 387-397.	4.6	14
33	Biosynthesis and chemical transformation of benzoxazinoids in rye during seed germination and the identification of a rye Bx6-like gene. <i>Phytochemistry</i> , 2017, 140, 95-107.	2.9	33
34	Alterations of the Benzoxazinoid Profiles of Uninjured Maize Seedlings During Freezing, Storage, and Lyophilization. <i>Journal of Agricultural and Food Chemistry</i> , 2017, 65, 4103-4110.	5.2	8
35	Quantification of azoxystrobin and identification of two novel metabolites in lettuce via liquid chromatography–quadrupole-linear ion trap (QTRAP) mass spectrometry. <i>International Journal of Environmental Analytical Chemistry</i> , 2017, 97, 419-430.	3.3	12
36	Liquid chromatography-tandem mass spectrometry method for simultaneous quantification of azoxystrobin and its metabolites, azoxystrobin free acid and 2-hydroxybenzonitrile, in greenhouse-grown lettuce. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2017, 34, 2173-2180.	2.3	5

#	ARTICLE	IF	CITATIONS
37	Biphenyl Columns Provide Good Separation of the Glucosides of DIMBOA and DIM2BOA. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	6
38	Bioactive small molecules in commercially available cereal food: Benzoxazinoids. <i>Journal of Food Composition and Analysis</i> , 2017, 64, 213-222.	3.9	9
39	Profiling and Metabolism of Sterols in the Weaver Ant Genus <i>Oecophylla</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	0
40	Dissipation kinetics of asparagine in soil measured by compound-specific analysis with metabolite tracking. <i>Biology and Fertility of Soils</i> , 2016, 52, 911-916.	4.3	5
41	Identification and Quantification of Loline-Type Alkaloids in Endophyte-Infected Grasses by LC-MS/MS. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 6212-6218.	5.2	15
42	Benzoxazinoids in Prostate Cancer Patients after a Rye-Intensive Diet: Methods and Initial Results. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 8235-8245.	5.2	16
43	Direct acquisition of organic N by white clover even in the presence of inorganic N. <i>Plant and Soil</i> , 2016, 407, 91-107.	3.7	31
44	Correlation of Deoxynivalenol Accumulation in Fusarium-Infected Winter and Spring Wheat Cultivars with Secondary Metabolites at Different Growth Stages. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 4545-4555.	5.2	21
45	Urea in Weaver Ant Feces: Quantification and Investigation of the Uptake and Translocation of Urea in <i>Coffea arabica</i> . <i>Journal of Plant Growth Regulation</i> , 2016, 35, 803-814.	5.1	8
46	Dietary exposure to benzoxazinoids enhances bacteria-induced monokine responses by peripheral blood mononuclear cells. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 2190-2198.	3.3	2
47	Metabolic Profiling of <i>Arabidopsis thaliana</i> Reveals Herbicide- and Allelochemical-Dependent Alterations Before They Become Apparent in Plant Growth. <i>Journal of Plant Growth Regulation</i> , 2015, 34, 96-107.	5.1	5
48	Benzoxazinoids: Cereal phytochemicals with putative therapeutic and health-protecting properties. <i>Molecular Nutrition and Food Research</i> , 2015, 59, 1324-1338.	3.3	71
49	Are ant feces nutrients for plants? A metabolomics approach to elucidate the nutritional effects on plants hosting weaver ants. <i>Metabolomics</i> , 2015, 11, 1013-1028.	3.0	10
50	2,4-Dihydroxy-7-methoxy-2 H -1,4-benzoxazin-3(4 H )-one (DIMBOA) inhibits trichothecene production by <i>Fusarium graminearum</i> through suppression of Tri6 expression. <i>International Journal of Food Microbiology</i> , 2015, 214, 123-128.	4.7	34
51	Application of the QuEChERS procedure and LC-MS/MS for the assessment of neonicotinoid insecticide residues in cocoa beans and shells. <i>Journal of Food Composition and Analysis</i> , 2015, 44, 149-157.	3.9	43
52	Threshold response of stomatal closing ability to leaf abscisic acid concentration during growth. <i>Journal of Experimental Botany</i> , 2014, 65, 4361-4370.	4.8	61
53	Quantification of neonicotinoid insecticide residues in soils from cocoa plantations using a QuEChERS extraction procedure and LC-MS/MS. <i>Science of the Total Environment</i> , 2014, 499, 276-283.	8.0	83
54	Foliar abscisic acid content underlies genotypic variation in stomatal responsiveness after growth at high relative air humidity. <i>Annals of Botany</i> , 2013, 112, 1857-1867.	2.9	45

#	ARTICLE	IF	CITATIONS
55	Nutritional composition of minor indigenous fruits: Cheapest nutritional source for the rural people of Bangladesh. Food Chemistry, 2013, 140, 466-470.	8.2	42
56	The Response of <i>Arabidopsis</i> to Co-cultivation with Clover. ACS Symposium Series, 2013, , 189-201.	0.5	4
57	Phenolic Acids from Wheat Show Different Absorption Profiles in Plasma: A Model Experiment with Catheterized Pigs. Journal of Agricultural and Food Chemistry, 2013, 61, 8842-8850.	5.2	12
58	Comparison of the levels of bioactive benzoxazinoids in different wheat and rye fractions and the transformation of these compounds in homemade foods. Food Chemistry, 2013, 141, 444-450.	8.2	51
59	Differences among five amaranth varieties ( <i>Amaranthus</i> spp.) regarding secondary metabolites and foliar herbivory by chewing insects in the field. Arthropod-Plant Interactions, 2013, 7, 235-245.	1.1	26
60	Absorption and metabolic fate of bioactive dietary benzoxazinoids in humans. Molecular Nutrition and Food Research, 2013, 57, 1847-1858.	3.3	37
61	Plasma and Urine Concentrations of Bioactive Dietary Benzoxazinoids and Their Glucuronidated Conjugates in Rats Fed a Rye Bread-Based Diet. Journal of Agricultural and Food Chemistry, 2012, 60, 11518-11524.	5.2	24
62	Bioactive Benzoxazinoids in Rye Bread Are Absorbed and Metabolized in Pigs. Journal of Agricultural and Food Chemistry, 2012, 60, 2497-2506.	5.2	33
63	Phytotoxic Effect, Uptake, and Transformation of Biochanin A in Selected Weed Species. Journal of Agricultural and Food Chemistry, 2012, 60, 10715-10722.	5.2	19
64	Fate in Soil of Flavonoids Released from White Clover ( <i>Trifolium repens</i> L.). Applied and Environmental Soil Science, 2012, 2012, 1-10.	1.7	39
65	Variation of Polyphenols and Betaines in Aerial Parts of Young, Field-Grown <i>Amaranthus</i> Genotypes. Journal of Agricultural and Food Chemistry, 2011, 59, 12073-12082.	5.2	29
66	Degradation of biochanin A in soil. Chemoecology, 2011, 21, 59-66.	1.1	6
67	Bread from common cereal cultivars contains an important array of neglected bioactive benzoxazinoids. Food Chemistry, 2011, 127, 1814-1820.	8.2	65
68	Variations in the polyphenol content of seeds of field grown <i>Amaranthus</i> genotypes. Food Chemistry, 2011, 129, 131-138.	8.2	57
69	Proximate composition, phenolic acids, and flavonoids characterization of commercial and wild nopal ( <i>Opuntia</i> spp.). Journal of Food Composition and Analysis, 2010, 23, 525-532.	3.9	121
70	Phenolic Compounds in Different Barley Varieties: Identification by Tandem Mass Spectrometry (QStar) and NMR; Quantification by Liquid Chromatography Triple Quadrupole-Linear Ion Trap Mass Spectrometry (Q-Trap). Natural Product Communications, 2010, 5, 1934578X1000500.	0.5	15
71	Synthesis and Quantitation of Six Phenolic Amides in <i>Amaranthus</i> spp.. Journal of Agricultural and Food Chemistry, 2010, 58, 6306-6311.	5.2	32
72	Allelochemicals in Rye ( <i>Secale Cereale</i> L.): Cultivar and Tissue Differences in the Production of Benzoxazinoids and Phenolic Acids. Natural Product Communications, 2009, 4, 1934578X0900400.	0.5	17

#	ARTICLE	IF	CITATIONS
73	Amaranth ( <i>Amaranthus hypochondriacus</i> ) as an alternative crop for sustainable food production: Phenolic acids and flavonoids with potential impact on its nutraceutical quality. <i>Journal of Cereal Science</i> , 2009, 49, 117-121.	3.7	144
74	Allelochemicals in rye ( <i>Secale cereale</i> L.): cultivar and tissue differences in the production of benzoxazinoids and phenolic acids. <i>Natural Product Communications</i> , 2009, 4, 199-208.	0.5	32
75	Flavonoids in roots of white clover: interaction of arbuscular mycorrhizal fungi and a pathogenic fungus. <i>Plant and Soil</i> , 2008, 302, 33-43.	3.7	72
76	Biologically active secondary metabolites in white clover ( <i>Trifolium repens</i> L.) – a review focusing on contents in the plant, plant–pest interactions and transformation. <i>Chemoecology</i> , 2008, 18, 129-170.	1.1	53
77	Benzoxazinoid concentrations show correlation with Fusarium Head Blight resistance in Danish wheat varieties. <i>Biochemical Systematics and Ecology</i> , 2008, 36, 245-259.	1.3	47
78	Fate and availability of glyphosate and AMPA in agricultural soil. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2008, 43, 365-375.	1.5	104
79	Transformation kinetics of 6-methoxybenzoxazolin-2-one in soil. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2008, 43, 1-7.	1.5	38
80	Variation in Flavonoids in Leaves, Stems and Flowers of White Clover Cultivars. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.5	6
81	Rebuttal on Results from the FATEALLCHEM Project. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 1645-1647.	5.2	0
82	Chemical Ecology in Wheat Plant–Pest Interactions. How the Use of Modern Techniques and a Multidisciplinary Approach Can Throw New Light on a Well-known Phenomenon: Allelopathy. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 987-990.	5.2	42
83	Fate of Benzoxazinone Allelochemicals in Soil after Incorporation of Wheat and Rye Sprouts. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1064-1074.	5.2	78
84	Transformation of Benzoxazinones and Derivatives and Microbial Activity in the Test Environment of Soil Ecotoxicological Tests on <i>Poecilus cupreus</i> and <i>Folsomia candida</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1086-1092.	5.2	43
85	Elucidating the Transformation Pattern of the Cereal Allelochemical 6-Methoxy-2-benzoxazolinone (MBOA) and the Trideuteriomethoxy Analogue [D3]-MBOA in Soil. <i>Journal of Agricultural and Food Chemistry</i> , 2006, 54, 1075-1085.	5.2	32
86	Biotransformation of 2-Benzoxazolinone to 2-Amino-(3H)-Phenoxazin-3-one and 2-Acetylamino-(3H)-Phenoxazin-3-one in Soil. <i>Journal of Chemical Ecology</i> , 2005, 31, 1205-1222.	1.8	52
87	Transformation products of 2-benzoxazolinone (BOA) in soil. <i>Chemosphere</i> , 2005, 61, 74-84.	8.2	38
88	First European interlaboratory study of the analysis of benzoxazinone derivatives in plants by liquid chromatography. <i>Journal of Chromatography A</i> , 2004, 1047, 69-76.	3.7	33
89	Microbial transformation products of benzoxazolinone and benzoxazinone allelochemicals – a review. <i>Chemosphere</i> , 2004, 54, 1025-1038.	8.2	106
90	Leaching of Pesticides Through Normal–Tillage and Low–Tillage Soil – A Lysimeter Study. I. Isoproturon. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2003, 38, 1-18.	1.5	9

#	ARTICLE	IF	CITATIONS
91	Leaching of Pesticides Through Normalâ€Tillegemeinde and Lowâ€Tillegemeinde Soilâ€Tillegemeinde”A Lysimeter Study. II. Glyphosate. Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes, 2003, 38, 19-35.	1.5	40
92	Comparison and evaluation of eight pesticide environmental risk indicators developed in Europe and recommendations for future use. Agriculture, Ecosystems and Environment, 2002, 90, 177-187.	5.3	161
93	Influence of microbial activity, organic carbon content, soil texture and soil depth on mineralisation rates of low concentrations of <sup>14</sup> C-mecopropâ€Tillegemeinde”development of a predictive model. Ecological Modelling, 1999, 122, 45-68.	2.5	20
94	Sampling and Substrate Application Methods for Pesticide Mineralization Experiments in Undisturbed Soil Samples. International Journal of Environmental Analytical Chemistry, 1998, 70, 121-132.	3.3	3
95	Degradation of Mecoprop and Isoproturon in Soil Influence of Initial Concentration. International Journal of Environmental Analytical Chemistry, 1998, 70, 133-148.	3.3	23
96	Degradation of <sup>14</sup> C-maneb in sediment from a Nicaraguan estuary. International Journal of Environmental Studies, 1998, 55, 175-198.	1.6	2
97	Modelling the mineralization kinetics for low concentrations of pesticides in surface and subsurface soil. Ecological Modelling, 1997, 102, 175-208.	2.5	44
98	Degradation of [ <sup>14</sup> C]ethylenethiourea in surface and subsurface soil. Science of the Total Environment, 1996, 191, 271-276.	8.0	6
99	Degradation of Pesticides in Subsurface Soils, Unsaturated Zoneâ€Tillegemeinde”a Review Of Methods and Results. International Journal of Environmental Analytical Chemistry, 1995, 58, 231-245.	3.3	53
100	Lead, arsenic, cadmium and copper in Lake Asososca, Nicaragua. Science of the Total Environment, 1994, 155, 229-236.	8.0	5
101	Toxaphene and Other Organochlorine Pesticides in Fish and Sediment from Lake XolotlÃ¡n, Nicaragua. International Journal of Environmental Analytical Chemistry, 1993, 53, 297-305.	3.3	14
102	Preliminary study of 15 organochlorine pesticides in Lake Xolotlan, Nicaragua. Chemosphere, 1992, 24, 1413-1419.	8.2	7
103	Mercury contamination in Lake XolotlÃ¡n (Managua). Hydrobiological Bulletin, 1991, 25, 173-176.	0.5	5