

# Celine A Beamer

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/9572776/celine-a-beamer-publications-by-year.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

17  
papers

547  
citations

12  
h-index

18  
g-index

18  
ext. papers

611  
ext. citations

5  
avg, IF

3.86  
L-index

#	Paper	IF	Citations
17	Nanoparticle-Induced Airway Eosinophilia Is Independent of ILC2 Signaling but Associated With Sex Differences in Macrophage Phenotype Development. <i>Journal of Immunology</i> , <b>2021</b> ,	5.3	1
16	Targeted deletion of the aryl hydrocarbon receptor in dendritic cells prevents thymic atrophy in response to dioxin. <i>Archives of Toxicology</i> , <b>2019</b> , 93, 355-368	5.8	10
15	Isolation and Identification of Innate Lymphoid Cells (ILCs) for Immunotoxicity Testing. <i>Methods in Molecular Biology</i> , <b>2018</b> , 1803, 353-370	1.4	8
14	Acute Exposure to Crystalline Silica Reduces Macrophage Activation in Response to Bacterial Lipoproteins. <i>Frontiers in Immunology</i> , <b>2016</b> , 7, 49	8.4	11
13	Environmental Immunology: Lessons Learned from Exposure to a Select Panel of Immunotoxicants. <i>Journal of Immunology</i> , <b>2016</b> , 196, 3217-25	5.3	39
12	IL-1R signalling is critical for regulation of multi-walled carbon nanotubes-induced acute lung inflammation in C57Bl/6 mice. <i>Nanotoxicology</i> , <b>2014</b> , 8, 17-27	5.3	42
11	Role of the aryl hydrocarbon receptor (AhR) in lung inflammation. <i>Seminars in Immunopathology</i> , <b>2013</b> , 35, 693-704	12	54
10	IL-33 mediates multi-walled carbon nanotube (MWCNT)-induced airway hyper-reactivity via the mobilization of innate helper cells in the lung. <i>Nanotoxicology</i> , <b>2013</b> , 7, 1070-81	5.3	64
9	Aryl hydrocarbon receptor (AhR) regulates silica-induced inflammation but not fibrosis. <i>Toxicological Sciences</i> , <b>2012</b> , 126, 554-68	4.4	32
8	COPD and Other Inflammatory Diseases of the Lung: Focus on AhR Signaling. <i>Molecular and Integrative Toxicology</i> , <b>2012</b> , 313-343	0.5	1
7	Inhibition of TLR ligand- and interferon gamma-induced murine microglial activation by Panax notoginseng. <i>Journal of NeuroImmune Pharmacology</i> , <b>2012</b> , 7, 465-76	6.9	16
6	Innate immune processes are sufficient for driving silicosis in mice. <i>Journal of Leukocyte Biology</i> , <b>2010</b> , 88, 547-57	6.5	66
5	Critical role of MARCO in crystalline silica-induced pulmonary inflammation. <i>Toxicological Sciences</i> , <b>2009</b> , 108, 462-71	4.4	74
4	Silica suppresses Toll-like receptor ligand-induced dendritic cell activation. <i>FASEB Journal</i> , <b>2008</b> , 22, 2053-63	5.9	21
3	Antigen-presenting cell population dynamics during murine silicosis. <i>American Journal of Respiratory Cell and Molecular Biology</i> , <b>2007</b> , 37, 729-38	5.7	31
2	Motheaten (me/me) mice deficient in SHP-1 are less susceptible to focal cerebral ischemia. <i>Journal of Neuroscience Research</i> , <b>2006</b> , 83, 1220-30	4.4	13
1	Scavenger receptor class A type I/II (CD204) null mice fail to develop fibrosis following silica exposure. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , <b>2005</b> , 289, L186-95	5.8	64

