

Ying-Ying Shen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9577976/publications.pdf>

Version: 2024-02-01

17
papers

214
citations

1163117

8
h-index

1058476

14
g-index

17
all docs

17
docs citations

17
times ranked

199
citing authors

#	ARTICLE	IF	CITATIONS
1	Physiological clearance of amyloid-beta by the kidney and its therapeutic potential for Alzheimer's disease. <i>Molecular Psychiatry</i> , 2021, 26, 6074-6082.	7.9	39
2	Amyloid-beta uptake by blood monocytes is reduced with ageing and Alzheimer's disease. <i>Translational Psychiatry</i> , 2020, 10, 423.	4.8	35
3	Brain Amyloid- β^2 Deposition and Blood Biomarkers in Patients with Clinically Diagnosed Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2019, 69, 169-178.	2.6	28
4	Dynamic changes of CSF sPDGFR β^2 during ageing and AD progression and associations with CSF ATN biomarkers. <i>Molecular Neurodegeneration</i> , 2022, 17, 9.	10.8	24
5	Association of Polygenic Risk Score with Age at Onset and Cerebrospinal Fluid Biomarkers of Alzheimer's Disease in a Chinese Cohort. <i>Neuroscience Bulletin</i> , 2020, 36, 696-704.	2.9	19
6	The Correlations Between Plasma Fibrinogen With Amyloid-Beta and Tau Levels in Patients With Alzheimer's Disease. <i>Frontiers in Neuroscience</i> , 2020, 14, 625844.	2.8	11
7	Polysaccharide Krestin Prevents Alzheimer's Disease-type Pathology and Cognitive Deficits by Enhancing Monocyte Amyloid- β^2 Processing. <i>Neuroscience Bulletin</i> , 2022, 38, 290-302.	2.9	11
8	Spicy food consumption is associated with cognition and cerebrospinal fluid biomarkers of Alzheimer disease. <i>Chinese Medical Journal</i> , 2021, 134, 173-177.	2.3	9
9	Association of the Polygenic Risk Score with the Incidence Risk of Parkinson's Disease and Cerebrospinal Fluid α -Synuclein in a Chinese Cohort. <i>Neurotoxicity Research</i> , 2019, 36, 515-522.	2.7	8
10	Genetic Association Between NGFR, ADAM17 Gene Polymorphism, and Parkinson's Disease in the Chinese Han Population. <i>Neurotoxicity Research</i> , 2019, 36, 463-471.	2.7	7
11	Associations of plasma soluble CD22 levels with brain amyloid burden and cognitive decline in Alzheimer's disease. <i>Science Advances</i> , 2022, 8, eabm5667.	10.3	6
12	The Correlations of Plasma Liver-Type Fatty Acid-Binding Protein with Amyloid- β^2 and Tau Levels in Patients with Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2022, 88, 375-383.	2.6	4
13	Establishment of combined diagnostic models of Alzheimer's disease in a Chinese cohort: the Chongqing Ageing & Dementia Study (CADS). <i>Translational Psychiatry</i> , 2022, 12, .	4.8	4
14	Neurotrophin Receptor p75 mRNA Level in Peripheral Blood Cells of Patients with Alzheimer's Disease. <i>Neurotoxicity Research</i> , 2019, 36, 101-107.	2.7	3
15	Circulating Naturally Occurring Antibodies to P2RY2 Are Decreased in Alzheimer's Disease. <i>Journal of Alzheimer's Disease</i> , 2022, 87, 711-719.	2.6	3
16	Associations of plasma angiostatin and amyloid- β^2 and tau levels in Alzheimer's disease. <i>Translational Psychiatry</i> , 2022, 12, 194.	4.8	2
17	The Association of Serum Neurofilament Light Chain and Acute Ischaemic Stroke Is Influenced by Effective Revascularization. <i>Disease Markers</i> , 2022, 2022, 1-6.	1.3	1